

**Link full download:  
Solution Manual:**

<https://testbankpack.com/p/solution-manual-for-calculus-single-variable-canadian-9th-edition-by-adams-isbn-9780134579801-0134579801/>

**Test Bank:**

<https://testbankpack.com/p/test-bank-for-calculus-single-variable-canadian-9th-edition-by-adams-isbn-9780134579801-0134579801/>

**CHAPTER 2. DIFFERENTIATION**

**Section 2.1 Tangent Lines and Their Slopes (page 100)**

1. Slope of y = 3x - 1 at x = 1/2 is

$$m = \lim_{h \rightarrow 0} \frac{3(1/2 + h) - 1 - (3(1/2) - 1)}{h} = \lim_{h \rightarrow 0} \frac{3h}{h} = 3$$

The tangent line is y = 3x - 1/2, or y = 3x - 1. (The tangent to a straight line at any point on it is the same straight line.)

2. Since y = x^2 is a straight line, its tangent at any point (a, a^2) on it is the same line y = x^2.

3. Slope of y = 2x^2 at x = 2 is

$$m = \lim_{h \rightarrow 0} \frac{2(2+h)^2 - 2(2)^2}{h} = \lim_{h \rightarrow 0} \frac{8h + 4h^2}{h} = 8$$

$$\lim_{h \rightarrow 0} \frac{8h + 4h^2}{h} = 8$$

Tangent line is y = 8x - 8 or y = 8x - 16.

4. The slope of y = 6x^2 at x = 2 is

$$m = \lim_{h \rightarrow 0} \frac{6(2+h)^2 - 6(2)^2}{h} = \lim_{h \rightarrow 0} \frac{24h + 6h^2}{h} = 24$$

The tangent line at (2, 24) is y = 24x - 24.

5. Slope of y = x^3 at x = 2 is

7. Slope of y = x^3 at x = 3 is

$$m = \lim_{h \rightarrow 0} \frac{(3+h)^3 - 3^3}{h} = \lim_{h \rightarrow 0} \frac{27 + 27h + 9h^2 + h^3 - 27}{h} = \lim_{h \rightarrow 0} \frac{27h + 9h^2 + h^3}{h} = 27$$

$$\lim_{h \rightarrow 0} \frac{27h + 9h^2 + h^3}{h} = 27$$

$$= 27$$

Tangent line is y = 27x - 54, or y = 27x - 54.

8. The slope of y = x^3 at x = 9 is

$$m = \lim_{h \rightarrow 0} \frac{(9+h)^3 - 9^3}{h} = \lim_{h \rightarrow 0} \frac{27h^2 + 27h + h^3}{h} = 27$$

The tangent line at (9, 729) is y = 27x - 189, or

y = 27x - 189.

$$m = \lim_{h \rightarrow 0} \frac{(9+h)^3 - 9^3}{h} = 27$$

The tangent line at (9, 729) is y = 27x - 189.

$$m = \lim_{h \rightarrow 0} \frac{2Ch^3 + C^2 - (2C^2)}{h} = \lim_{h \rightarrow 0} \frac{2Ch^3 + C^2 - 2C^2}{h} = \lim_{h \rightarrow 0} \frac{2Ch^3 - C^2}{h}$$

Tangent line is  $y = 2Cx + C^2$  or  $y = 2Cx - C^2$ .

6. The slope of  $y = x^2 + 1$  at  $x = 0$ ;  $1/$  is

9. Slope of  $y = \frac{2x}{x^2 - 2}$  at  $x = 2$  is

$$m = \lim_{h \rightarrow 0} \frac{\frac{2(2+h)}{(2+h)^2 - 2} - \frac{2(2)}{2^2 - 2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{2(2+h)}{4+h^2} - \frac{4}{2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{2(2+h) - 4(2)}{4+h^2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2(2+h) - 8}{h(4+h^2)}$$

$$= \lim_{h \rightarrow 0} \frac{4+2h-8}{h(4+h^2)}$$

$$= \lim_{h \rightarrow 0} \frac{-4+2h}{h(4+h^2)}$$

$$= \lim_{h \rightarrow 0} \frac{-4+2h}{h(4+h^2)}$$

Tangent line is  $y = \frac{1}{4}x - \frac{1}{2}$ ,

10. The slope of  $y = \frac{1}{x^2}$  at  $x = 1$  is

$$m = \lim_{h \rightarrow 0} \frac{\frac{1}{(1+h)^2} - \frac{1}{1^2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{1}{1+2h+h^2} - 1}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\frac{1 - (1+2h+h^2)}{1+2h+h^2}}{h}$$

$$= \lim_{h \rightarrow 0} \frac{-2h-h^2}{h(1+2h+h^2)}$$

$$= \lim_{h \rightarrow 0} \frac{-2-h}{1+2h+h^2}$$

$$= \frac{-2-1}{1+2(0)+0^2}$$

$$= \frac{-3}{1}$$

$$= -3$$

$$= \lim_{h \rightarrow 0} \frac{-2-h}{1+2h+h^2}$$

The tangent line at  $(1, \frac{1}{2})$  is  $y = -\frac{3}{2}x + \frac{5}{2}$ , or

$$y = -\frac{3}{2}x + \frac{5}{2}$$

11. Slope of  $y = x^2$  at  $x = 0$  is

$$m = \lim_{h \rightarrow 0} \frac{(0+h)^2 - 0^2}{h} = \lim_{h \rightarrow 0} \frac{h^2}{h} = \lim_{h \rightarrow 0} h = 0$$

Tangent line is  $y = 2x_0 + x_0^2$   
 or  $y = 2x_0 + x_0^2$

12. The slope of  $y = x^2$  at  $x = a$  is  $2a$   
 $m = \lim_{h \rightarrow 0} \frac{(a+h)^2 - a^2}{h} = \lim_{h \rightarrow 0} \frac{a^2 + 2ah + h^2 - a^2}{h} = \lim_{h \rightarrow 0} \frac{2ah + h^2}{h} = \lim_{h \rightarrow 0} (2a + h) = 2a$

The tangent line at  $x = a$  is  $y = 2ax - a^2$   
 $y = 2ax - a^2$

13. Since  $\lim_{h \rightarrow 0} \frac{1}{h}$  does not exist (and is not  $\infty$  or  $-\infty$ ), the graph of  $f(x) = \frac{1}{x}$  has no tangent at  $x = 0$ .

14. The slope of  $f(x) = x^{1/4}$  at  $x = 1$  is  $\frac{1}{4}$   
 $m = \lim_{h \rightarrow 0} \frac{(1+h)^{1/4} - 1}{h} = 0$

The graph of  $f$  has a tangent line with slope 0 at  $x = 1$ . Since  $f(1) = 1$ , the tangent has equation  $y = 1$ .

15. The slope of  $f(x) = 2x^{3/5}$  at  $x = 2$  is  $\frac{6}{5}$   
 $m = \lim_{h \rightarrow 0} \frac{2(2+h)^{3/5} - 2 \cdot 2^{3/5}}{h} = \frac{6}{5}$

The graph of  $f$  has vertical tangent  $x = 2$  at  $x = 2$ .

16. The slope of  $f(x) = x^2 - 1$  at  $x = 1$  is  $2$   
 $m = \lim_{h \rightarrow 0} \frac{(1+h)^2 - 1}{h} = 2$   
 which does not exist, and is not  $\infty$  or  $-\infty$ . The graph of  $f$  has no tangent at  $x = 1$ .

17. If  $f(x) = \frac{1}{x}$ , then  
 $\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{h} - \frac{1}{0}}{h}$   
 $\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{h} - \frac{1}{0}}{h}$

19. a) Slope of  $y = x^3$  at  $x = a$  is

$$m = \lim_{h \rightarrow 0} \frac{(a+h)^3 - a^3}{h} = \lim_{h \rightarrow 0} \frac{a^3 + 3a^2h + 3ah^2 + h^3 - a^3}{h} = \lim_{h \rightarrow 0} \frac{3a^2h + 3ah^2 + h^3}{h} = \lim_{h \rightarrow 0} (3a^2 + 3ah + h^2) = 3a^2$$

b) We have  $m = 3$  if  $3a^2 = 3$ , i.e., if  $a = \pm 1$ .

Lines of slope 3 tangent to  $y = x^3$  are  $y = 3x - 2$  and  $y = 3x + 2$ .

20. The slope of  $y = x^3$  at  $x = a$  is  $3a^2$   
 $m = \lim_{h \rightarrow 0} \frac{(a+h)^3 - a^3}{h} = \lim_{h \rightarrow 0} \frac{a^3 + 3a^2h + 3ah^2 + h^3 - a^3}{h} = \lim_{h \rightarrow 0} (3a^2 + 3ah + h^2) = 3a^2$

At points where the tangent line is parallel to the x-axis, the slope is zero, so such points must satisfy  $3a^2 = 0$ . Thus,  $a = 0$ . Hence, the tangent line is parallel to the x-axis at the points  $(0, 0)$  and  $(0, 0)$ .

21. The slope of the curve  $y = x^3 + 1$  at  $x = a$  is

$$m = \lim_{h \rightarrow 0} \frac{(a+h)^3 + 1 - (a^3 + 1)}{h} = \lim_{h \rightarrow 0} \frac{a^3 + 3a^2h + 3ah^2 + h^3 + 1 - a^3 - 1}{h} = \lim_{h \rightarrow 0} (3a^2 + 3ah + h^2) = 3a^2$$

The tangent at  $x = a$  is parallel to the line  $y = 2x + 5$  if  $3a^2 = 2$ , that is, if  $a = \pm \sqrt{2/3}$ . The corresponding points on the curve are  $(\sqrt{2/3}, 1 + 2\sqrt{2/3})$  and  $(-\sqrt{2/3}, 1 - 2\sqrt{2/3})$ .

22. The slope of the curve  $y = 1/x$  at  $x = a$  is

$$m = \lim_{h \rightarrow 0} \frac{\frac{1}{a+h} - \frac{1}{a}}{h} = \lim_{h \rightarrow 0} \frac{\frac{a - (a+h)}{(a+h)a}}{h} = \lim_{h \rightarrow 0} \frac{-h}{h(a+h)a} = \lim_{h \rightarrow 0} \frac{-1}{(a+h)a} = -\frac{1}{a^2}$$

Thus the graph of  $f$  has a vertical tangent at  $x = 0$ .

18. The slope of  $y = x^2 - 1$  at  $x = x_0$  is

$$m = \lim_{h \rightarrow 0} \frac{(x_0 + h)^2 - 1 - (x_0^2 - 1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2x_0 h + h^2}{h} = 2x_0$$

If  $m = 3$ , then  $x_0 = \frac{3}{2}$ . The tangent line with slope

$m = 3$  at  $(\frac{3}{2}, \frac{5}{4})$  is  $y - \frac{5}{4} = 3(x - \frac{3}{2})$ , that is,

$$y = 3x - \frac{1}{4}.$$

The tangent at  $x = a$  is perpendicular to the line  $y = 4x - 3$  if  $1 = a' = 4$ , that is, if  $a = \frac{1}{4}$ . The corresponding points on the curve are  $(\frac{1}{2}, \frac{1}{4})$  and  $(\frac{1}{2}, \frac{1}{4})$ .

23. The slope of the curve  $y = x^2$  at  $x = a$  is

$$m = \lim_{h \rightarrow 0} \frac{(a + h)^2 - a^2}{h} = \lim_{h \rightarrow 0} \frac{2ah + h^2}{h} = 2a$$

The normal at  $x = a$  has slope  $m = -\frac{1}{2a}$ , and has equation

$$y - a^2 = -\frac{1}{2a}(x - a); \text{ or } \frac{x}{2a} + y = \frac{1}{2} + a^2.$$

This is the line  $x = 2$  if  $2a = 1$ , and so  $k = 1/2$ .  $k = 1/2$  if  $2 = 1/k$ .

24. The curves  $y = kx^2$  and  $y = k/x^2$  intersect at  $(1, k)$ .

The slope of  $y = kx^2$  at  $x = 1$  is

$$m_1 = \lim_{h \rightarrow 0} \frac{k(1+h)^2 - k}{h} = \lim_{h \rightarrow 0} \frac{2kh + kh^2}{h} = 2k$$

The slope of  $y = k/x^2$  at  $x = 1$  is

$$m_2 = \lim_{h \rightarrow 0} \frac{k/(1+h)^2 - k}{h} = \lim_{h \rightarrow 0} \frac{-2kh - kh^2}{h(1+h)^2} = -2k$$

The two curves intersect at right angles if  $2k = -2k$ , that is, if  $k = 0$ , which is satisfied if  $k = 0$ .

25. Horizontal tangents at  $(0, 0)$ ,  $(3, 108)$ , and  $(5, 0)$ .




Fig. 2.1-25

26. Horizontal tangent at  $(-1, 8)$  and  $(2, -19)$ .




Fig. 2.1-26




Fig. 2.1-27

28. Horizontal tangent at  $(a, 2/a)$  and  $(-a, -2/a)$  for all  $a > 1$ . No tangents at  $(1, 2)$  and  $(-1, -2)$ .




Fig. 2.1-28

29. Horizontal tangent at  $(0, 1/3)$ . The tangents at  $(-1, 0)$  are vertical.




Fig. 2.1-29

27. Horizontal tangent at  $x = 2; y = 5/4$ . No tangents at  $x = 1; y = 1/4$  and  $x = 1; y = 1$ .
30. Horizontal tangent at  $x = 0; y = 1/4$ . No tangents at  $x = 1; y = 0$  and  $x = 1; y = 0$ .





Fig. 2.1-30

2.



31. The graph of the function  $f(x) = x^{2/3}$  (see Figure 2.1.7 in the text) has a cusp at the origin  $O$ , so does not have a tangent line there. However, the angle between  $OP$  and the positive  $y$ -axis does  $\rightarrow 0$  as  $P$  approaches  $O$  along the graph. Thus the answer is NO.

32. The slope of  $f(x)$  at  $x = a$  is

$$m = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

Since  $f(a+h) = a_0 + a_1 h + a_2 h^2 + \dots + a_n h^n$  and  $f(a) = a_0$ , the slope is

$$m = \lim_{h \rightarrow 0} \frac{a_0 + a_1 h + a_2 h^2 + \dots + a_n h^n - a_0}{h} = \lim_{h \rightarrow 0} (a_1 + a_2 h + \dots + a_n h^{n-1}) = a_1$$

Thus the line  $y - f(a) = m(x - a)$  is tangent to  $y = f(x)$  at  $x = a$  if and only if  $m = a_1$  and  $b = f(a)$ , that is, if and only if


$$f(x) = a_0 + a_1(x-a) + a_2(x-a)^2 + a_3(x-a)^3 + \dots + a_n(x-a)^n$$

$$D_x f = a_1 + 2a_2(x-a) + \dots + na_n(x-a)^{n-1}$$


where  $Q$  is a polynomial.

**Section 2.2 The Derivative (page 107)**


1.



3.



4.



5. Assuming the tick marks are spaced 1 unit apart, the function  $f$  is differentiable on the intervals  $(-2, -1)$ ,  $(-1, 1)$ , and  $(1, 2)$ .

6. Assuming the tick marks are spaced 1 unit apart, the function  $g$  is differentiable on the intervals  $(-2, -1)$ ,  $(-1, 0)$ ,  $(0, 1)$ , and  $(1, 2)$ .



7.  $y = f(x)$  has its minimum at  $x = 3$  where  $f'(x) = 0$




Fig. 2.2-7




Fig. 2.2-9

8.  $y = Df(x)/Dx$  has horizontal tangents at the points near  $x=1=2$  and  $x=3=2$  where  $f^0(x)/Dx = 0$




Fig. 2.2-8

10.  $y = Df(x)/Dx$  is constant on the intervals  $[-1, -2/3]$ ;  $[-1/3, 1/3]$ ; and  $[2/3, 1]$ . It is not differentiable at  $x = D^{-2}$  and  $x = D^{-1}$ .




Fig. 2.2-10

9.  $y = Df(x)/Dx$  fails to be differentiable at  $x = D^{-1}$ ,  $x = D^0$ , and  $x = D^1$

11.  $y = D^2 x^2 = 3x$

$$y^0 = D \lim_{h \rightarrow 0} \frac{x + Ch^2 - 3xCh^2 - (x^2 - 3x)}{h} = \lim_{h \rightarrow 0} \frac{2xh + Ch^2 - 3h}{h} = 2x - 3$$

1. It has horizontal tangents at two points, one between 1 and 0 and the other between 0 and 1.

$$D_x \left( \frac{1}{2}x^2 - \frac{3}{2}x \right) = x - \frac{3}{2}$$

$$D_x \left( \frac{1}{2}x^2 - \frac{3}{2}x \right) = 0 \implies x - \frac{3}{2} = 0 \implies x = \frac{3}{2}$$

12. f(x) = 4x<sup>2</sup> - 5x + 1

$$f'(x) = \lim_{h \rightarrow 0} \frac{4(x+h)^2 - 5(x+h) + 1 - (4x^2 - 5x + 1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{4x^2 + 8xh + 4h^2 - 5x - 5h + 1 - 4x^2 + 5x - 1}{h}$$

f'(x) = 8x - 5

13. f(x) = x<sup>3</sup>

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^3 - x^3}{h}$$

$$= \lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2 + h^3}{h}$$

f'(x) = 3x<sup>2</sup>

14. s(t) = 3t<sup>4</sup> - 4t<sup>3</sup> + 5t<sup>2</sup> - 6t + 7

$$s'(t) = \lim_{h \rightarrow 0} \frac{3(t+h)^4 - 4(t+h)^3 + 5(t+h)^2 - 6(t+h) + 7 - (3t^4 - 4t^3 + 5t^2 - 6t + 7)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{12t^3h + 6t^2h^2 + 2th^3 + 12t^2h - 12th^2 - 6h^3 + 10th - 6h^2 - 6}{h}$$

s'(t) = 12t<sup>3</sup> + 6t<sup>2</sup> + 2t - 6

s'(2) = 22

15. g(x) = 2x<sup>2</sup> - 3x + 1

$$g'(x) = \lim_{h \rightarrow 0} \frac{2(x+h)^2 - 3(x+h) + 1 - (2x^2 - 3x + 1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{4xh + 2h^2 - 3h}{h} = 4x + 2h - 3$$

g'(2) = 5

17. f(t) = 2t<sup>3</sup> - 5t + 1

$$f'(t) = \lim_{h \rightarrow 0} \frac{2(t+h)^3 - 5(t+h) + 1 - (2t^3 - 5t + 1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{6t^2h + 6th^2 + 2h^3 - 5h}{h}$$

f'(t) = 6t<sup>2</sup> - 5

$$f'(1) = 6(1)^2 - 5 = 1$$

f'(1) = 1

18. f(x) = 4x<sup>3</sup> - 3x<sup>2</sup> + 2x - 1

$$f'(x) = \lim_{h \rightarrow 0} \frac{4(x+h)^3 - 3(x+h)^2 + 2(x+h) - 1 - (4x^3 - 3x^2 + 2x - 1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{12x^2h + 6xh^2 + 2h^3 - 6xh - 3h^2 + 2h}{h}$$

$$= 12x^2 + 6xh + 2h^2 - 6x - 3h + 2$$

f'(x) = 12x<sup>2</sup> - 6x + 2

19. y = x<sup>2</sup> - 3x + 1

$$y'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^2 - 3(x+h) + 1 - (x^2 - 3x + 1)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2 - 3h}{h} = 2x + h - 3$$

$$D \frac{4}{x^2} = \frac{4}{x^2}$$

$$D \frac{4}{x^2} dx$$

$$D \lim_{x \rightarrow 1} \frac{1}{x^2} = \lim_{x \rightarrow 1} \frac{-2}{x^3} = -2$$

16.  $y = x^3$

$$D \frac{1}{3} x^3 = x^2$$

20.  $z = \frac{1}{s}$

$$D \frac{1}{s} = -\frac{1}{s^2}$$

$$D \lim_{h \rightarrow 0} \frac{1}{h^3} = -\frac{3}{h^4}$$

$$\lim_{h \rightarrow 0} \frac{1}{h^3} = \infty$$

$$D \lim_{h \rightarrow 0} \frac{x+h}{h} = \frac{1}{h^2}$$

$$D \lim_{h \rightarrow 0} \frac{1}{h^2} = \infty$$

$$D \lim_{h \rightarrow 0} \frac{1}{h^2} = \infty$$

$$D \frac{1}{s^2} = -\frac{2}{s^3}$$

$$D \frac{1}{x^2} = -\frac{2}{x^3}$$

21. 
$$f(x) = \frac{1}{\sqrt{1-x^2}}$$

$$f'(x) = \frac{1}{2} (1-x^2)^{-3/2} \cdot (-2x) = \frac{-x}{(1-x^2)^{3/2}}$$

$$f''(x) = \frac{-(1-x^2)^{3/2} - (-x) \cdot 3(1-x^2)^{1/2} \cdot (-2x)}{(1-x^2)^3}$$

$$= \frac{-(1-x^2)^{3/2} - 6x^2(1-x^2)^{1/2}}{(1-x^2)^3}$$

$$= \frac{-(1-x^2) - 6x^2}{(1-x^2)^{5/2}} = \frac{-1+x^2-6x^2}{(1-x^2)^{5/2}} = \frac{-1-5x^2}{(1-x^2)^{5/2}}$$

$$dF(x)/dx = \frac{1}{2} \frac{d}{dx} (1-x^2)^{-3/2}$$

22. 
$$y = x^2$$

$$\lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{x^2 + 2xh + h^2 - x^2}{h} = \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = \lim_{h \rightarrow 0} (2x + h) = 2x$$

$$dy = 2x dx$$

23. 
$$y = \frac{1}{x}$$

$$y'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h} = \lim_{h \rightarrow 0} \frac{\frac{x - (x+h)}{(x+h)x}}{h} = \lim_{h \rightarrow 0} \frac{-h}{h(x+h)x} = \lim_{h \rightarrow 0} \frac{-1}{(x+h)x} = \frac{-1}{x^2}$$

25. Since  $f(x) = \frac{1}{|x|}$ , for  $x \neq 0$ ,  $f$  will become continuous at  $x = 0$  if we define  $f(0) = \frac{1}{0}$ . However,  $f$  will still not be differentiable at  $x = 0$  since  $|x|$  is not differentiable at  $x = 0$ .

$$x^2 \quad \text{if } x > 0$$

become continuous and differentiable at  $x = 0$  if we define  $f(0) = \frac{1}{0}$ .

27.  $f(x) = x^2 + 3x + 2$  fails to be differentiable where

$x^2 + 3x + 2 = 0$ , that is, at  $x = -2$  and  $x = -1$ . Note: both of these are single zeros of  $x^2 + 3x + 2$ .

were higher order zeros.

a factor of  $(x + 2)^n$  for some integer  $n \geq 2$  then  $f$  would be differentiable at the corresponding point.

28.  $y = x^3 + 2x$

$x$	$f(x) = x^3 + 2x$	$f'(x) = 3x^2 + 2$
1:1	1:31000	1:31000
1:01	1:03010	1:03010
1:001	1:00300	1:00300
1:0001	1:00030	1:00030
0:9999	0:99970	0:99970

$$d(x^3 + 2x) = (3x^2 + 2) dx$$

$$D_x (x^3 + 2x) = 3x^2 + 2$$

29.  $f(x) = \frac{1}{x^2}$

$x$	$f(x) = \frac{1}{x^2}$	$f'(x) = -\frac{2}{x^3}$
1:9	0:26316	0:23810
1:99	0:25126	0:24876
1:999	0:25013	0:24988
1:9999	0:25001	0:24999

dy  $D$   $2.1 C x^{3-2} dx$

24.  $f(t) = C t^3$   
 $\lim_{h \rightarrow 0} \frac{f(t+h) - f(t)}{h} = \lim_{h \rightarrow 0} \frac{C(t+h)^3 - C t^3}{h}$

$$\lim_{h \rightarrow 0} \frac{C(t^3 + 3t^2h + 3t h^2 + h^3) - C t^3}{h} = \lim_{h \rightarrow 0} \frac{3C t^2 h + 3C t h^2 + C h^3}{h}$$

$$\lim_{h \rightarrow 0} \frac{3C t^2 h + 3C t h^2 + C h^3}{h} = \lim_{h \rightarrow 0} (3C t^2 + 3C t h + C h^2) = 3C t^2$$

$D_{h \rightarrow 0} h \cdot t^2 C 3 / C h^2 C 3$   
 $D_{t^2} C 3/2$

$\frac{12t}{12t}$   
 $df(t) = D t^2 C 3/2 dt$

$f(x) = C x^2$   
 $\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{C(x+h)^2 - C x^2}{h}$

30. The slope of  $y = 5 C 4x - x^2$  at  $x = 2$  is

$$\frac{dy}{dx} = 5 C 4 - 2x$$

$$\lim_{h \rightarrow 0} \frac{5 C 4(x+h) - 2(x+h)^2 - (5 C 4x - 2x^2)}{h} = \lim_{h \rightarrow 0} \frac{5 C 4h - 2(x^2 + 2xh + h^2) + 2x^2}{h}$$

$$= \lim_{h \rightarrow 0} (5 C 4 - 2x - 2h) = 5 C 4 - 2x$$

Thus, the tangent line at  $x = 2$  has the equation  $y = 9$ .

31.  $y = \frac{1}{x^3}$ . Slope at  $x = 3$ ;  $\frac{3}{16}$  is  

$$m = \lim_{h \rightarrow 0} \frac{\frac{1}{(3+h)^3} - \frac{1}{3^3}}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{27+h^3} - \frac{1}{27}}{h} = \frac{1}{27}$$
 Tangent line is  $y = \frac{1}{27}x - \frac{2}{9}$ , or  $x - 6y = 15$ .

32. The slope of  $y = t^2 - 2$  at  $t = 2$  and  $y = 1$  is  

$$\frac{dy}{dt} = \lim_{h \rightarrow 0} \frac{(2+h)^2 - 2 - (2^2 - 2)}{h} = \lim_{h \rightarrow 0} \frac{2h + 2 - 2}{h} = 2$$
 Thus, the tangent line has the equation  
 $y - 1 = 2(t - 2)$ , that is,  $y = 2t - 3$ .

33.  $y = t^2 - 2t$ . Slope at  $t = a$  is  

$$m = \lim_{h \rightarrow 0} \frac{(a+h)^2 - 2(a+h) - (a^2 - 2a)}{h} = \lim_{h \rightarrow 0} \frac{a^2 + 2ah + h^2 - 2a - 2h - a^2 + 2a}{h} = \lim_{h \rightarrow 0} \frac{2ah - 2h + h^2}{h} = 2a - 2$$
 Tangent line is  $y = (2a - 2)t - a^2 + 2a$ .

34.  $f(x) = 17x^{18}$  for  $x \neq 0$

35.  $g(t) = 22t^{21}$  for all  $t$

36.  $\frac{dy}{dx} = \frac{1}{3}x^{2-3}$  for  $x \neq 0$   

$$\frac{dy}{dx} = \frac{1}{3}x^{-1}$$

44. The slope of  $y = \frac{1}{x^2}$  at  $x = x_0$  is  

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{\frac{1}{(x_0+h)^2} - \frac{1}{x_0^2}}{h} = \lim_{h \rightarrow 0} \frac{\frac{1}{x_0^2+h^2} - \frac{1}{x_0^2}}{h} = -\frac{2}{x_0^3}$$

Thus, the equation of the tangent line is

$y - \frac{1}{x_0^2} = -\frac{2}{x_0^3}(x - x_0)$ , that is,  $y = \frac{x - x_0}{x_0^2}$ .

45. Slope of  $y = x^2$  at  $x = a$  is  

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{(a+h)^2 - a^2}{h} = \lim_{h \rightarrow 0} \frac{2ah + h^2}{h} = 2a$$

Normal has slope  $-\frac{1}{2a}$ , and equation  $y - a^2 = -\frac{1}{2a}(x - a)$

or  $y = a^2 - \frac{1}{2a}(x - a)$

46. The intersection points of  $y = x^2$  and  $x^2 + 4y = 18$  satisfy

$$\begin{aligned} 4x^2 + 4x^2 &= 18 \\ 4x^2 &= 9/x \end{aligned}$$

Therefore  $x = \frac{9}{4}$  or  $x = -\frac{9}{4}$ .

The slope of  $y = x^2$  is  $m_1 = 2x$ .

At  $x = \frac{9}{4}$ ,  $m_1 = \frac{9}{2}$ . At  $x = -\frac{9}{4}$ ,  $m_1 = -\frac{9}{2}$ .

The slope of  $x^2 + 4y = 18$ , i.e.  $y = \frac{18 - x^2}{4}$ , is  $m_2 = -\frac{x}{2}$ .

Thus, at  $x = \frac{9}{4}$ , the product of these slopes is

$-\frac{9}{2} \cdot \frac{1}{2} = -\frac{9}{4} \neq -1$ . So, the curve and line intersect at right angles at that point.



37.  $\frac{dx}{dt} = 3x^{4-3}$  for  $x \neq 0$   
 $\frac{dx}{dt} = 3x$

38.  $\frac{d}{dt} t^{2.25} = 2.25 t^{0.25}$  for  $t > 0$

39.  $\frac{ds}{ds} = \frac{119}{4s^{115-4}}$  for  $s > 0$

40.  $\frac{d}{dt} t^{2.25} = 2.25 t^{0.25}$

41.  $F(x) = x^4$ ;  $F'(x) = 4x^3$ ;  $F''(x) = 12x^2$ ;  $F'''(x) = 24x$

42.  $\frac{d}{dx} x^{2.5} = 2.5 x^{1.5}$

43.  $\frac{dy}{dt} = \frac{1}{t^4} = t^{-4}$

47. Let the point of tangency be  $(a, a^2)$ . Slope of tangent is

$\frac{dy}{dx} = 2x$

$2a$

is the slope from

$\frac{a^2 - C}{a - 2a}$ , and

$\frac{a^2 - C}{-a} = 2a$

$a^2 - C = 2a^2$

$-C = a^2$

$C = -a^2$

$C = -1$  or  $C = 1$

The two tangent lines are

(for  $C = 3$ ):  $y = 9 - 6x$  or  $6x = 9 - y$

(for  $C = 1$ ):  $y = 1 - 2x$  or  $y = 2x - 1$




Fig. 2.2-47

If  $b < a^2$ , i.e.  $a^2 - b > 0$ , then there are two real solutions. Therefore, there will be two distinct tangent lines passing through  $(a, a^2)$  and  $(b, b^2)$  with equations

$y = 2ax - a^2$  and  $y = 2bx - b^2$ . If  $b = a^2$ , then there will be only one tangent line with slope  $2a$  and equation  $y = 2ax - a^2$ .

If  $b > a^2$ , then  $a^2 - b < 0$ . There will be no real solution for  $t$ . Thus, there will be no tangent line.

48. The slope of  $y = x^2$  at  $(a, a^2)$  is

$$\frac{dy}{dx} = 2x \Big|_{x=a} = 2a$$

If the slope is  $2$ , then  $2a = 2$ , or  $a = 1$ . Therefore, the equations of the two straight lines are

$$y = 2x - 1 \quad \text{and} \quad y = 2x - 1$$


or  $y = 2x - 1$ .

49. Let the point of tangency be  $(a, a^2)$ .

Slope of tangent is  $2a$ .

Thus  $\frac{1}{2a} = \frac{1}{2a}$ , so  $a = 1$ .

The required slope is  $2$ .



51. Suppose  $f$  is odd:  $f(-x) = -f(x)$ . Then

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0-h)}{h - (-h)} = \lim_{h \rightarrow 0} \frac{f(h) - (-f(h))}{2h} = \lim_{h \rightarrow 0} \frac{2f(h)}{2h} = \lim_{h \rightarrow 0} \frac{f(h)}{h}$$

Let  $h = k$ . Then  $\lim_{k \rightarrow 0} \frac{f(k)}{k} = \lim_{k \rightarrow 0} \frac{f(-k)}{-k} = -\lim_{k \rightarrow 0} \frac{f(k)}{k}$ .

Thus  $f'(0)$  is even.

Now suppose  $f$  is even:  $f(-x) = f(x)$ . Then

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0-h)}{h - (-h)} = \lim_{h \rightarrow 0} \frac{f(h) - f(h)}{2h} = \lim_{h \rightarrow 0} \frac{0}{2h} = 0$$

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(-h)}{h - (-h)} = \lim_{h \rightarrow 0} \frac{f(h) - f(h)}{2h} = \lim_{h \rightarrow 0} \frac{0}{2h} = 0$$

so  $f'(0)$  is odd.

52. Let  $f(x) = x^n$ . Then

$$f'(0) = \lim_{h \rightarrow 0} \frac{(0+h)^n - (0-h)^n}{h - (-h)} = \lim_{h \rightarrow 0} \frac{h^n - (-1)^n h^n}{2h} = \lim_{h \rightarrow 0} \frac{h^n(1 - (-1)^n)}{2h} = \lim_{h \rightarrow 0} \frac{h^{n-1}(1 - (-1)^n)}{2}$$

Fig. 2.2-49

50. If a line is tangent to  $y = x^2$  at  $(t, t^2)$ , then its slope is

$\frac{dy}{dx} = 2t$ . If this line also passes through  $(a, b)$ , then

its

$$\frac{t^2 - b}{t - a} = 2t; \text{ that is } t^2 - 2at + b = 0:$$

lim

$$\lim_{h \rightarrow 0} \frac{x^n - (x+h)^n}{h} = \frac{0}{0}$$

$$\lim_{h \rightarrow 0} \frac{x^n - (x+h)^n}{h} = \frac{0}{0}$$

$$x^{n-1} - nx^{n-2}(x+h) = x^{n-1} - nx^{n-1} - nh^{n-1}$$

$$\lim_{h \rightarrow 0} \frac{-nh^{n-1}}{h} = \lim_{h \rightarrow 0} -nh^{n-2} = 0$$

53.  $f(x) = x^3$

$$f'(x) = \lim_{h \rightarrow 0} \frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{h} = \lim_{h \rightarrow 0} \frac{3x^2h + 3xh^2 + h^3}{h} = \lim_{h \rightarrow 0} (3x^2 + 3xh + h^2) = 3x^2$$

54. Let  $f(x) = x^{1-n}$ . Then

$$f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^{1-n} - x^{1-n}}{h} = \lim_{h \rightarrow 0} \frac{x^{1-n} + (1-n)x^{-n}h + \dots - x^{1-n}}{h} = \lim_{h \rightarrow 0} \frac{(1-n)x^{-n}h + \dots}{h} = (1-n)x^{-n} = -nx^{1-n-1}$$

55.  $\frac{d}{dx} x^n = nx^{n-1}$

$$\lim_{h \rightarrow 0} \frac{(x+h)^n - x^n}{h} = \lim_{h \rightarrow 0} \frac{x^n + nx^{n-1}h + \dots - x^n}{h} = \lim_{h \rightarrow 0} (nx^{n-1} + \dots) = nx^{n-1}$$

If  $f'(a)$  is finite, call the half-line with equation  $y = f'(a)(x-a)$ , the right tangent line to the graph of  $f$  at  $x = a$ . Similarly, if  $f'(a)$  is finite, call the half-line  $y = f'(a)(x-a)$ , the left tangent line to the graph of  $f$  at  $x = a$ . The graph has a tangent line at  $x = a$  if and only if both quantities may be  $\pm\infty$  or both may be finite. In this case the right and left tangents are two opposite halves of the same straight line. For  $f(x) = x^2$ ,  $f'(0) = 0$ . In this case both left and right tangents are the positive  $y$ -axis, and the curve does not have a tangent line at the origin.

For  $f(x) = |x|$ , we have  $f'(x) = 1$  if  $x > 0$  and  $f'(x) = -1$  if  $x < 0$ . The right tangent is  $y = x$ , and the left tangent is  $y = -x$ . There is no tangent line at  $x = 0$ .

**Section 2.3 Differentiation Rules (page 115)**

- $y = 3x^2 - 5x + 7$ ;  $y' = 6x - 5$
- $y = 4x^{1/2} - x^2$ ;  $y' = 2x^{-1/2} - 2x = \frac{2}{\sqrt{x}} - 2x$
- $f(x) = \frac{Ax^6 + Bx^2 + C}{x^2}$ ;  $f'(x) = \frac{6Ax^5 + 2Bx - 2(Ax^6 + Bx^2 + C)}{x^4}$
- $f(x) = x^3 + x^2 - 2$ ;  $f'(x) = 3x^2 + 2x$
- $z = 15t^{-3}$ ;  $\frac{dz}{dt} = -45t^{-4} = -\frac{45}{t^4}$
- $y = x^{45}$ ;  $y' = 45x^{44}$
- $g(t) = 2t^{1/3} + 3t^{1/5}$ ;  $g'(t) = \frac{2}{3}t^{-2/3} + \frac{3}{5}t^{-4/5}$

$$c^{n \cdot n} \frac{1}{1} \frac{2}{2} \frac{3}{3} x^{n^3} h^2 c c h^n \frac{1}{1}$$


---


$$1/n \ 2/$$

$$D n x^{n-1}$$

56. Let

$$f^0 \cdot a / \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

$$f^0 \cdot a / \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

$$g^0 \cdot t / D 3t \quad C 2t^{3=4} C 5t$$

$$d \frac{t^3 - 2}{t^3} \frac{2}{5=2} \quad 2=3 \quad 3=2$$

$$d t D 2t^{1=3} C 3t$$

9.  $\frac{d}{du} D \frac{3}{5} x^{5=3} \frac{5}{3} x^{3=5}$

$$dx D x^{2=3} C x^{8=5}$$

10.  $F \cdot x / D \cdot 3x \quad 2/1 \quad 5x/$

$$F^0 \cdot x / D 3 \cdot 1 \quad 5x / C \cdot 3x \quad 2/ \cdot 5 / D 13 \quad 30x$$

$$11. \quad y^0 \frac{D}{Dx} x^5 - x^3 = 5x^4 - 3x^2 = 2x^3(5x^2 - 3x^0) = 2x^3(5x^2 - 3)$$

$$12. \quad g(t) = \frac{1}{2t^3}; \quad g'(t) = \frac{D}{Dt} \frac{1}{2t^3} = \frac{1}{2} \cdot \frac{-3}{t^4} = -\frac{3}{2t^4}$$

$$13. \quad y^0 \frac{D}{Dx} x^2 = 2x$$

$$14. \quad y^0 \frac{D}{Dx} x^{-3} = -3x^{-4} = -\frac{3}{x^4}$$

$$15. \quad f(t) = \frac{1}{2}t^2$$

$$f'(t) = \frac{D}{Dt} \frac{1}{2}t^2 = \frac{1}{2} \cdot 2t = t$$

$$16. \quad g(y) = 1 - y^2; \quad g'(y) = -2y$$

$$17. \quad f(x) = \frac{4x^2}{x^3} = \frac{4}{x}$$

$$f'(x) = \frac{D}{Dx} \frac{4}{x} = -\frac{4}{x^2}$$

$$f^0(x) = \frac{D}{Dx} 3x^4 = 12x^3$$

$$18. \quad g(u) = \frac{1}{3}u^3 = \frac{1}{3}u^3$$

$$g'(u) = \frac{D}{Du} \frac{1}{3}u^3 = \frac{1}{3} \cdot 3u^2 = u^2$$

$$19. \quad y = 2t^2 - t^3$$

$$\frac{dy}{dt} = \frac{D}{Dt} (2t^2 - t^3) = 4t - 3t^2$$

$$23. \quad s = \frac{1}{t^2}$$

$$\frac{ds}{dt} = \frac{D}{Dt} t^{-2} = -2t^{-3} = -\frac{2}{t^3}$$

$$\frac{D}{Dt} \frac{1}{x^3} = -\frac{3}{x^4}$$

$$24. \quad f(x) = x^{-1}$$

$$f'(x) = \frac{D}{Dx} x^{-1} = -x^{-2} = -\frac{1}{x^2}$$

$$25. \quad f(x) = \frac{ax + b}{cx + d}$$

$$f'(x) = \frac{D}{Dx} \frac{ax + b}{cx + d} = \frac{a(cx + d) - c(ax + b)}{(cx + d)^2} = \frac{ad - bc}{(cx + d)^2}$$

$$26. \quad F(t) = \frac{t^2 - 7t + 8}{t}$$

$$F'(t) = \frac{D}{Dt} \frac{t^2 - 7t + 8}{t} = \frac{t(t - 7) + (t^2 - 7t + 8)(-1)}{t^2} = \frac{t^2 - 7t - t^2 + 7t - 8}{t^2} = \frac{-8}{t^2}$$

$$27. \quad f(x) = \frac{1}{x} + \frac{1}{2x} + \frac{1}{3x} + \frac{1}{4x} = \frac{1}{x} \left( 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} \right) = \frac{1}{x} \left( \frac{12}{12} + \frac{6}{12} + \frac{4}{12} + \frac{3}{12} \right) = \frac{25}{12x}$$

OR

$$f(x) = \frac{1}{x} + \frac{1}{2x} + \frac{1}{3x} + \frac{1}{4x} = \frac{1}{x} \left( 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} \right) = \frac{1}{x} \left( \frac{12}{12} + \frac{6}{12} + \frac{4}{12} + \frac{3}{12} \right) = \frac{25}{12x}$$

$$dt \quad D \quad C 2 p t \quad C 2 \quad D \quad 2 t \quad p \quad t$$

$$20. \quad z \quad D \quad \frac{x-1}{x^2=3} \quad D \quad X^{1=3} \quad x \quad 2=3$$

$$\frac{dz}{dx} = \frac{1}{x^2=3} \quad 2_x \quad 5=3 \quad D \quad x \quad C \quad 2$$

$$D \quad 3 \quad C \quad 3 \quad 3x^{5=3}$$

$$21. \quad f \quad .x / D \quad \frac{4x}{3 C 4x}$$

$$f^0 \quad .x / \quad \frac{.3 C 4x \cdot 4 / \quad .3 \quad 4x / .4 /}{D \quad .3 \quad 4x^2}$$

$$D \quad \frac{24 \quad C}{.3 \quad C \quad 4x^2}$$

$$22. \quad z \quad \frac{t^2 C 2t}{D \quad t^2 - 1}$$

$$z^0 \quad \frac{t^2 - 1 \cdot 2t C 2 / \quad - t^2 C 2t / .2t /}{D \quad t^2 \quad 1^2}$$

$$D \quad t^2 \quad 1^2$$

$$D \quad t^2 \quad 1^2$$

$$D \quad 1 \quad C \quad 10x \quad C \quad 25x^2 \quad C \quad 10x^2 \quad .1 \quad C \quad 5x / \quad C \quad 24x^4$$

$$D \quad 1 \quad C \quad 10x \quad C \quad 35x^2 \quad C \quad 50x^3 \quad C \quad 24x^4$$

$$f^0 \quad .x / \quad D \quad 10 \quad C \quad 70x \quad C \quad 150x^2 \quad C \quad 96x^3$$

$$28. \quad f \quad .r / D \quad r^2 \quad C \quad r^3 \quad 4 / .r^2 \quad C \quad r^3 \quad C \quad 1 /$$

$$f^0 \quad .r / D \quad . \quad 2r^3 \quad 3r^4 \quad / .r^2 \quad C \quad r^3 \quad C \quad 1 /$$

$$C \quad .r^2 \quad C \quad r^3 \quad 4 / .2r \quad C \quad 3r^2 /$$

$$\text{or}$$

$$1 \quad 2 \quad 3 \quad 2 \quad 3$$

$$f \quad .r / D \quad 2 \quad C \quad r \quad C \quad r \quad C \quad r \quad C \quad r \quad 4r \quad 4r$$

$$29. \quad f^0 \quad .r / D \quad r^2 \quad 2r^3 \quad 3r^4 \quad C \quad 1 \quad 8r \quad 12r^2$$

$$y \quad D \quad .x^2 \quad C \quad 4 / \quad p \quad \bar{x} \quad C \quad 1 / .5x^{2=3} \quad 2 /$$

$$y^0 \quad D \quad 2x \quad p \quad x \quad C \quad 1 / .5x^{2=3} \quad 2 /$$

$$\frac{-1}{-}$$

$$C \quad 2p \quad x \quad .x^2 \quad C \quad 4 / .5x^{2=3} \quad 2 /$$

$$\frac{10}{C \quad 3x \quad .x \quad C \quad 4 / \quad x \quad C \quad 1 /}$$

30.  $y' = \frac{x^2 C_1 + x^3 C_2}{x^2 C_2 + x^3 C_1 + x^5}$

$D \frac{C_1 x^3 + C_2 x^2 - C_2}{x^5 C_2 x^3 + x^2 C_2}$

$y_0 = \frac{x^5 C_2 x^3 + x^2 C_2 - 2x^4 C_2 + 3x^2 C_2 + 4x}{x^5 C_2 x^3 + x^2 C_2}$

$\frac{x^5 C_2 x^3 + x^2 C_2 - 2x^4 C_2 + 3x^2 C_2 + 4x}{x^5 C_2 x^3 + x^2 C_2}$

$D \frac{2x^7 - 3x^6 - 3x^4 - 6x^2 C_2 + 4x}{x^5 C_2 x^3 + x^2 C_2}$

$D \frac{2x^7 - 3x^6 - 3x^4 - 6x^2 C_2 + 4x}{x^2 C_2 x^2 + x^3 C_1}$

31.  $y = \frac{x}{2x C_3} + \frac{3x^2 C_1}{6x C_2 x C_1}$

$D \frac{-1}{2x C_3} + \frac{D}{2} \frac{3x^2 C_1}{6x C_2 x C_1}$

$y_0 = \frac{6x^2 C_2 x C_1 + 6x C_1 - 3x^2 C_1 + 12x C_2}{6x^2 C_2 x C_1}$

$D \frac{6x^2 C_2 x C_1 + 6x C_1 - 3x^2 C_1 + 12x C_2}{6x^2 C_2 x C_1}$

$D \frac{6x^2 C_2 x C_1 + 6x C_1 - 3x^2 C_1 + 12x C_2}{6x^2 C_2 x C_1}$

$\frac{p}{x} = \frac{1}{x} - \frac{x}{1-x^2}$

32.  $f(x) = \frac{1}{x} - \frac{x}{1-x^2}$

$\frac{1}{x} - \frac{x}{1-x^2}$

$f'(x) = \frac{1}{x^2} - \frac{1-x^2}{(1-x^2)^2}$

$\frac{1}{x^2} - \frac{1-x^2}{(1-x^2)^2}$

$\frac{3 C_2 x - 1}{7.2} - \frac{4x C_3 x^2 - 2x^2 C_1 x^3}{7.2}$

$\frac{2x - 1}{x^2}$

$D \frac{2x^{3-2} - 3 C_2 x}{3 \cdot 2x^2}$

36.  $\frac{d}{dx} \frac{C}{x^2} = \frac{-2Cx}{x^4} = -\frac{2C}{x^3}$

$D \frac{C}{x^2} = \frac{-2Cx}{x^4} = -\frac{2C}{x^3}$

$\frac{C}{x^2} = \frac{C}{x^2}$

$D \frac{C}{x^2} = \frac{-2Cx}{x^4} = -\frac{2C}{x^3}$

$\frac{d}{dx} \frac{4}{x^2} = \frac{-8}{x^3}$

$\frac{d}{dx} \frac{8}{x^2} = \frac{-16}{x^3}$

$\frac{d}{dx} \frac{8}{x^2} = \frac{-16}{x^3}$

$\frac{d}{dx} \frac{8}{x^2} = \frac{-16}{x^3}$

$D \frac{32}{642}$

38.  $\frac{d}{dt} \frac{1}{t^4} = \frac{-4}{t^5}$

$\frac{d}{dt} \frac{1}{t^4} = \frac{-4}{t^5}$

$\frac{d}{dt} \frac{1}{t^4} = \frac{-4}{t^5}$

$D \frac{1}{16}$

$\frac{1}{4} - \frac{1}{2} = -\frac{1}{4}$

39.  $f(x) = \frac{1}{x^2} = x^{-2}$

$f'(x) = -2x^{-3} = -\frac{2}{x^3}$

$\frac{d}{dx} \frac{1}{x^2} = -\frac{2}{x^3}$



$$1 - \frac{4x^3 + 5x^2 + 12x + 7}{x^2}$$

33.  $\frac{d}{dx} \left( \frac{f(x)}{x^2} \right)$

$$\frac{4f'(x) - 4f(x)}{x^3}$$

34.  $\frac{d}{dx} \left( \frac{f}{x^2} \right)$

$$\frac{4f'(x) - 4f(x)}{16}$$

35.  $\frac{d}{dx} (x^2 f(x))$

$$4f'(x) - 4f(x) + 20$$

$$f'(x) = \frac{d}{dx} (9x^2 + 18x + 2)$$

$$= 18x + 18$$

41.  $y = \frac{3x^4 + 2x^2}{4x^3 + 1}$

$$y' = \frac{12x^3 + 4x}{4x^3 + 1} - \frac{3x^4 + 2x^2}{(4x^3 + 1)^2}$$

Slope of tangent at  $x = 1$ ;  $y' = 2$  is  $m = 2$

Tangent line has the equation  $y = 2x - 4$

42. For  $y = \frac{6}{x - 1}$  we calculate

$$y' = \frac{-6}{(x - 1)^2}$$

At  $x = 2$  we have  $y = 3$  and  $y' = 2$ . Thus, the equation of the tangent line is  $y = 3 + 2(x - 2)$ , or  $y = 2x - 1$ .  
The normal line is  $y = 3 - 2(x - 2)$ , or  $y = -2x + 7$ .




Fig. 2.3-47

43.  $y = \frac{1}{x^2}$ ,  $y' = -\frac{2}{x^3}$   
For horizontal tangent:  $0 = y' = -\frac{2}{x^3}$  so  $x^2 = 1$  and  $x = \pm 1$   
The tangent is horizontal at  $(1, 1)$  and at  $(-1, 1)$

44. If  $y = \frac{4}{x^2}$ , then  
 $y' = -\frac{8}{x^3}$

The slope of a horizontal line must be zero, so  
 $-\frac{8}{x^3} = 0$  which implies that  $x = 0$  or  $x = \pm 2$ .  
At  $x = 0$ ;  $y = 0$  and at  $x = \pm 2$ ;  $y = 1$ .  
Hence, there are two horizontal lines that are tangent to

the curve. Their equations are  $y = 0$  and  $y = 1$ .

45.  $y = \frac{1}{x^2}$ ,  $y' = -\frac{2}{x^3}$

For horizontal tangent we want  $0 = y' = -\frac{2}{x^3}$

Thus  $2x^3 = 0$  and  $x = 0$

The tangent is horizontal only at  $(0, 0)$ .

46. If  $y = \frac{2}{x-1}$ , then

$$y' = -\frac{2}{(x-1)^2}$$

In order to be parallel to  $y = 4x$ , the tangent line must have slope equal to 4, i.e.,

$$-\frac{2}{(x-1)^2} = 4$$

48. Since  $y = \frac{1}{x^2}$ ,  $y' = -\frac{2}{x^3}$   
The slope of  $y = \frac{1}{x^2}$  at  $x = 1$  is  $-\frac{2}{1^3} = -2$

$$\frac{dy}{dx} = -\frac{2}{x^3}$$

The product of the slopes of two curves intersect at right angles is  $-1$ .

49. The tangent to  $y = x^3$  at  $(a, a^3)$  has equation  
 $y - a^3 = 3a^2(x - a)$ , or  $y = 3a^2x - 2a^3$ . This line passes through  $(2, 8)$  if  $8 = 6a^2 - 2a^3$ , or, equivalently, if  $a^3 - 3a^2 + 4 = 0$ . Since  $(2, 8)$  lies on  $y = x^3$ ,  $a = 2$  must be a solution of this equation. In fact it must be a double root;  $(a - 2)^2$  must be a factor of  $a^3 - 3a^2 + 4$ . Dividing by this factor, we find that the other factor is  $a + 1$ , that is,

$$a^3 - 3a^2 + 4 = (a - 2)^2(a + 1)$$

The two tangent lines to  $y = x^3$  passing through  $(2, 8)$

correspond to  $a = 2$  and  $a = -1$ , so their equations are  $y = 12x - 16$  and  $y = 3x - 2$ .

50. The tangent to  $y = x^2 = x$  at  $(a, a^2)$  has slope  $m = 2a$

$$y - a^2 = 2a(x - a)$$

$$\frac{1}{2}x^2 - \frac{1}{5}x + a = a^2 - 2a$$

Hence  $x = 2a - \frac{1}{5}$ , and  $x = \frac{3}{2}$  or  $\frac{5}{2}$ . At  $x = \frac{3}{2}$ ,  $y = 1$ , and at  $x = \frac{5}{2}$ ,  $y = 3$ .

Hence, the tangent is parallel to  $y = 4x$  at the points  $(\frac{3}{2}, 1)$  and  $(\frac{5}{2}, 3)$ .

$(\frac{3}{2}, 1)$  and  $(\frac{5}{2}, 3)$ .

47. Let the point of tangency be  $(a, \frac{1}{a})$ . The slope of the tan-

gent is  $-\frac{1}{a^2}$ . Thus  $b = a$  and  $a = b$ .

$b^2$   
Tangent has slope  $-\frac{1}{a^2}$  so has equation  $y - \frac{1}{a} = -\frac{1}{a^2}(x - a)$ .

The equation of the tangent is  $y - \frac{1}{a} = -\frac{1}{a^2}(x - a)$ .

$$y = -\frac{1}{a^2}x + \frac{1}{a} + \frac{1}{a} = -\frac{1}{a^2}x + \frac{2}{a}$$

This line passes through  $(2, 0)$  provided

$$-\frac{2}{a^2} + \frac{2}{a} = 0$$

or, upon simplification,  $3a^2 - 4a = 0$ . Thus we can have either  $a = 0$  or  $a = \frac{4}{3}$ . There are two tangents through  $(2, 0)$ . Their equations are  $y = 0$  and  $y = 8x - 16$ .

51. 
$$\frac{d}{dx} x^p = \lim_{h \rightarrow 0} \frac{(x+h)^p - x^p}{h}$$

$$= \lim_{h \rightarrow 0} \frac{x^p \left( \left(1 + \frac{h}{x}\right)^p - 1 \right)}{h}$$

$$= x^p \lim_{h \rightarrow 0} \frac{\left(1 + \frac{h}{x}\right)^p - 1}{\frac{h}{x}}$$

$$= x^p \lim_{u \rightarrow 0} \frac{(1+u)^p - 1}{u} = x^p \cdot p = px^{p-1}$$

52.  $f(x) = \begin{cases} x^3 & \text{if } x < 0 \\ x^3 & \text{if } x \geq 0 \end{cases}$ . Therefore  $f$  is differentiable everywhere except possibly at  $x = 0$ . However,

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{h^3 - 0}{h} = \lim_{h \rightarrow 0} h^2 = 0$$

$$\lim_{h \rightarrow 0} \frac{f(0-h) - f(0)}{-h} = \lim_{h \rightarrow 0} \frac{-(0-h)^3 - 0}{-h} = \lim_{h \rightarrow 0} \frac{-(-h^3)}{-h} = \lim_{h \rightarrow 0} \frac{-h^3}{-h} = \lim_{h \rightarrow 0} h^2 = 0$$

Thus  $f'(0)$  exists and equals 0. We have

$$f'(x) = \begin{cases} 3x^2 & \text{if } x < 0 \\ 3x^2 & \text{if } x \geq 0 \end{cases}$$

53. To be proved:  $\frac{d}{dx} x^n = nx^{n-1}$  for  $n = 1, 2, 3, \dots$ .

Proof: It is already known that the case  $n = 1$  is true: the derivative of  $x^1 = x$  is  $1 = 2x^{1-1}$ . Assume that the formula is valid for  $n = k$  for some positive integer  $k$ :

$$\frac{d}{dx} x^k = kx^{k-1}$$

Then, by the Product Rule and this hypothesis,

$$\frac{d}{dx} x^{k+1} = \frac{d}{dx} (x \cdot x^k) = x \cdot \frac{d}{dx} x^k + x^k \cdot \frac{d}{dx} x = x \cdot kx^{k-1} + x^k \cdot 1 = kx^k + x^k = (k+1)x^k = (k+1)x^{(k+1)-1}$$

Thus the formula is also true for  $n = k + 1$ . Therefore it is

Proof: The case  $n = 2$  is just the Product Rule. Assume the formula holds for  $n = k$  for some integer  $k > 2$ . Using the Product Rule and this hypothesis we calculate

$$\frac{d}{dx} (f \cdot g) = f'g + fg'$$

$$\frac{d}{dx} (f \cdot g^k) = f'g^k + kfg^{k-1}g' = f'g^k + kf'fg^{k-1}$$

$$\frac{d}{dx} (f \cdot g^{k+1}) = f'(g^{k+1})' + (f \cdot g^{k+1})' = f'g^{k+1} + (f \cdot g^k)'g + f \cdot (g^k)'$$

$$= f'g^{k+1} + (f'g^k + kf'fg^{k-1})g + f \cdot kfg^{k-1}g' = f'g^{k+1} + f'g^{k+1} + kf'fg^k + kf'fg^k = f'g^{k+1} + 2kf'fg^k$$

so the formula is also true for  $n = k + 1$ . The formula is therefore for all integers  $n \geq 2$  by induction.

**Section 2.4 The Chain Rule (page 120)**

- $y = 2x^3$ ;  $y' = 6x^2$ ;  $y' = 12x^2$
- $y = 1 - 3x^98$ ;  $y' = -3 \cdot 98x^{97} = -294x^{97}$
- $f(x) = 4 - x^{2/10}$ ;  $f'(x) = -\frac{2}{10}x^{-8/10} = -\frac{1}{5}x^{-4/5} = -\frac{1}{5}x^{-0.8}$
- $\frac{dx}{dt} = 2x$ ;  $\frac{d}{dt} x^2 = 2x \cdot \frac{dx}{dt} = 2x \cdot 2x = 4x^2$
- $F(t) = 10 - 2t^3$ ;  $F'(t) = -6t^2$
- $z = 1 - x^{2/3}$ ;  $z' = -\frac{2}{3}x^{-1/3} = -\frac{2}{3}x^{-0.333}$
- $y = 5 - \frac{4x}{3}$ ;  $y' = -\frac{4}{3}$

Thus the formula is also true for  $n = k + 1$ . Therefore it is

true for all positive integers  $n$  by induction. For negative  $n$

$D^m$  (where  $m > 0$ ) we have

$$\frac{d}{dx} x^{n-2} = \frac{d}{dx} x^{m-2} \quad \text{where } m = n-2$$

$$\frac{d}{dx} x^{m-2} = \frac{d}{dx} x^{m-2} \quad \text{where } m = n-2$$

$$\frac{d}{dx} x^{m-2} = \frac{d}{dx} x^{m-2} \quad \text{where } m = n-2$$

54. To be proved:

$$f_1 f_2 \dots f_n = f_n f_1 f_2 \dots f_{n-1}$$

$$y^0 = D^{-1} 4x^{2/3} = 4 \int 4x^{2/3} dx$$

$$8. y = D^{-1} 2t^{2/3-2} = D^{-1} 2t^{-4/3}$$

$$y = D^{-3} 1 = \frac{1}{2} t^{2/5-2} = \frac{1}{2} t^{-8/5} = \frac{1}{2} t^{-1.6}$$

$$9. y = D^{-1} x^2; y^0 = D^{-2} x^2 = \frac{1}{2} x^2 = \frac{1}{2} x^2$$

$$10. f(t) = D^{-2} C t^3 = \frac{1}{2} C t^3 = \frac{3}{2} C t^2 = \frac{3}{2} C t^2$$

$$11. y = D^{-4} C x^4 = \frac{1}{24} C x^4 = \frac{1}{24} C x^4$$

12.  $y = D^{-2} C x^3 / 1=3$   
 $y^0 = D^{-1} \cdot 2 C x^3 / 2=3 = 3x^2 / \text{sgn} \cdot x /$

$D x^2 \cdot 2 C x^3 / 2=3 = \frac{x}{2} D x x^2 \cdot 2 C x^3 / 2=3$

$\frac{2 C}{3x} \cdot \frac{4}{C}$

1  $\frac{C}{C_3} \cdot C$   
 $D \frac{C}{2}$

$2^p \cdot 3x \cdot C \cdot 4 \cdot 2 C \cdot p \cdot 3x \cdot C \cdot 4$

14.  $f(x) = D^{-1} C \cdot \frac{x-2}{3} \cdot 4$   
 $\frac{x-2}{3} \cdot 4$   
 $\frac{2}{3} \cdot \frac{4}{x-2} \cdot \frac{d}{dx}$

15.  $z = D^{-5} C u^{-1} \cdot 5=3$   
 $\frac{du}{dz} = \frac{3}{5} C u^{-1} \cdot 1 = \frac{8=3}{5} \cdot \frac{1}{u^{1/2}}$   
 $D^{-3} \cdot 1 \cdot u^{-1/2} C u^{-1} \cdot 8=3$   
 $\frac{1}{5} \cdot \frac{1}{u^{1/2}} \cdot \frac{1}{u}$

16.  $y = D^{-5} \frac{p}{3} C x^6$   
 $\frac{4}{3} C x^2 / 3$


$y^0 = D^{-1} \cdot \frac{1}{3} \cdot 4 C x^{2/3} \cdot 5x^4 \cdot \frac{1}{3} C x^6 \cdot C x^5$   
 $\frac{4}{3} C x^{2/3} \cdot 5x^4 \cdot \frac{1}{3} C x^6 \cdot C x^5$

$D C h C \cdot 2^4 \cdot i \cdot 6$

$\frac{4}{6x} \cdot \frac{x^2}{5x^4} \cdot \frac{3}{x^6} \cdot \frac{1}{3x^{10}} \cdot \frac{1}{x^5} \cdot \frac{3}{3x^6}$   
 $\frac{4}{6x} \cdot \frac{x^2}{5x^4} \cdot \frac{3}{x^6} \cdot \frac{1}{3x^{10}} \cdot \frac{1}{x^5} \cdot \frac{3}{3x^6}$

$\frac{4}{3} C x / 3 C x$

18.



19.  $\frac{d}{dx} x^{1=4} = \frac{d}{dx} x^3 = 3x^2$   
 $\frac{d}{dx} \frac{p}{x} = \frac{d}{dx} p x^{-1} = -p x^{-2} = -\frac{p}{x^2}$

20.  $\frac{d}{dx} x^{3=4} = \frac{d}{dx} x^{-1} = -x^{-2} = -\frac{1}{x^2}$   
 $\frac{d}{dx} \frac{p}{x} = \frac{d}{dx} p x^{-1} = -p x^{-2} = -\frac{p}{x^2}$

21.  $\frac{d}{dx} x^{3=2} = \frac{d}{dx} x^{1/2} = \frac{1}{2} x^{-1/2} = \frac{1}{2\sqrt{x}}$   
 $\frac{d}{dx} \frac{p}{x} = \frac{d}{dx} p x^{-1} = -p x^{-2} = -\frac{p}{x^2}$

22.  $\frac{d}{dt} t \cdot 2t C / 3 = D^{-1} 2t C / 3$

23.  $\frac{d}{dx} f \cdot 5x \cdot x^2 = D^{-1} 5 \cdot 2x f \cdot 0.5x \cdot x^2$

24.  $\frac{d}{dx} f \cdot x^3 = D^{-3} 3f \cdot x^2 = 2f \cdot 0 \cdot x x^2$   
 $\frac{d}{dx} f \cdot x^2 = D^{-2} 2f \cdot x = 2f \cdot 0 \cdot x x^2$

$D^{-2} x^2 f \cdot 0 \cdot x f \cdot 2$

25.  $\frac{d}{dx} \frac{p}{3 C 2f \cdot x} = D^{-1} \frac{-2f \cdot 0 \cdot x}{3 C 2f \cdot x^2} = D^{-1} \frac{-f \cdot 0 \cdot x}{3 C 2f \cdot x^2}$

$\frac{d}{dx} \frac{p}{3 C 2f \cdot x} = D^{-1} \frac{-2f \cdot 0 \cdot x}{3 C 2f \cdot x^2} = D^{-1} \frac{-f \cdot 0 \cdot x}{3 C 2f \cdot x^2}$

$\frac{d}{dx} \frac{p}{3 C 2f \cdot x} = D^{-1} \frac{-2f \cdot 0 \cdot x}{3 C 2f \cdot x^2} = D^{-1} \frac{-f \cdot 0 \cdot x}{3 C 2f \cdot x^2}$

$\frac{d}{dx} \frac{p}{3 C 2f \cdot x} = D^{-1} \frac{-2f \cdot 0 \cdot x}{3 C 2f \cdot x^2} = D^{-1} \frac{-f \cdot 0 \cdot x}{3 C 2f \cdot x^2}$

27.  $\frac{d}{dx} \frac{p}{3x^6} = \frac{d}{dx} p x^{-6} = -6 p x^{-7} = -\frac{60x^4}{3x^6} C$   
 $\frac{d}{dx} \frac{p}{32x^{10} C 2x^{12}}$


$$\frac{dx}{dt} = 3Cx^2 - x$$

$$0.3Cx^2 - x$$

17.

$$D \left( \frac{1}{4}Cx^{2/4} + \frac{1}{3}Cx^6 \right)$$

$$28. \frac{d}{dt} (2f - 3f \cdot x)$$



$$f(x) = 2f^0 + 3f(x) - 3f^0(x)$$

$$29. \frac{d}{dx} (2f^0 + 3f(x) - 5t) = 2 \cdot 0 + 3 \cdot f'(x) - 0 = 3f'(x)$$

$$D_x (2 + 3f(x) - 5t) = 0 + 3f'(x) - 0 = 3f'(x)$$

$$D_x (15f^0 + 2 \frac{5t}{f} - 3f(x) - 5t) = 0 - 2 \frac{5t}{f^2} f'(x) - 3f'(x) - 0 = -\frac{10t}{f^2} f'(x) - 3f'(x)$$



30. 
$$d \frac{p}{x^2} = -\frac{2p}{x^3}$$

$$= -\frac{2p}{x^3} \cdot \frac{x^2}{x^2} = -\frac{2px}{x^5}$$

$$= -\frac{2px}{x^5}$$

31. 
$$d \frac{t^3}{3} = t^2$$

$$= t^2$$

32. 
$$f(x) = \frac{1}{2x} = \frac{1}{2} x^{-1/2}$$

$$f'(x) = \frac{1}{2} \cdot (-\frac{1}{2}) x^{-3/2} = -\frac{1}{4} x^{-3/2}$$

$$= -\frac{1}{4x^{3/2}}$$

33. 
$$y = x^3 \cdot 9^{17} = 9^{17} x^3$$

$$y' = 9^{17} \cdot 3x^2 = 3 \cdot 9^{17} x^2$$

34. 
$$F(x) = 2x^2 + 3x^3 + 4x^4$$

$$F'(x) = 4x + 9x^2 + 16x^3$$

37. Slope of  $y = \frac{1}{3} x^{2/3}$  at  $x = 1$  is  $\frac{2}{9}$ .  
The tangent line at  $(1, \frac{2}{9})$  has equation  $y - \frac{2}{9} = \frac{2}{9}(x - 1)$ .

38. The slope of  $y = \frac{a}{b} x^8$  at  $x = a$  is  $\frac{8a^7}{b}$ .  
The equation of the tangent line at  $(a, \frac{a^8}{b})$  is  $y - \frac{a^8}{b} = \frac{8a^7}{b}(x - a)$ .

39. Slope of  $y = x^2 + \frac{3}{x}$  at  $x = 2$  is  $\frac{5}{3}$ .  
The tangent line at  $(2, \frac{17}{3})$  has equation  $y - \frac{17}{3} = \frac{5}{3}(x - 2)$ .

40. Given that  $f(x) = \frac{a}{x} + \frac{b}{x^2}$ , then  $f'(x) = -\frac{a}{x^2} - \frac{2b}{x^3}$ .  
If  $x = a$  and  $x = b$ , then  $f'(x) = 0$  if and only if  $mx + mb + nx + na = 0$ , which is equivalent to  $\frac{n}{m} = \frac{a}{b}$ .

41.  $x \cdot x^2 = x^3$ ,  $\frac{d}{dx} x^3 = 3x^2$ .  
42.  $4.7x^4 = 49x^2$ ,  $\frac{d}{dx} 49x^2 = 98x$ .

$$x^2 = 3x^{5/2} \implies 2x = 15x^{3/2} \implies 2 = 15x^{1/2} \implies x = \frac{4}{9}$$

36. The slope of  $y = \frac{1}{2}x^2$  at  $x = 2$  is

$$\frac{dy}{dx} = x \implies \left. \frac{dy}{dx} \right|_{x=2} = 2$$

Thus, the equation of the tangent line at  $(2, 2)$  is  $y - 2 = 2(x - 2)$

$$y = 2x - 2, \text{ or } y = 2x - 2$$

857; 592

44.  $5 = 8$

45. The Chain Rule does not enable you to calculate the derivatives of  $|x|^2$  and  $|x^2|$  at  $x = 0$  directly as a composition of two functions, one of which is  $|x|$ , because  $|x|$  is not differentiable at  $x = 0$ . However,  $|x|^2 = x^2$  and  $|x^2| = x^2$ , so both functions are differentiable at  $x = 0$  and have derivative 0 there.

46. It may happen that  $k \neq g(x)h/g(x) = 0$  for values of  $h$  arbitrarily close to 0 so that the division by  $k$  in the “proof” is not justified.

**Section 2.5 Derivatives of Trigonometric Functions (page 126)**

1.  $\frac{d}{dx} \csc x = -\csc x \cot x$ ;  $\frac{d}{dx} \sin x = \cos x$ ;  $\frac{d}{dx} \sin^2 x = 2 \sin x \cos x$ ;  $\frac{d}{dx} \csc x \cot x = -\csc x \cot^2 x - \csc^2 x$
2.  $\frac{d}{dx} \cot x = -\csc^2 x$ ;  $\frac{d}{dx} \sin x = \cos x$ ;  $\frac{d}{dx} \sin^2 x = 2 \sin x \cos x$ ;  $\frac{d}{dx} \csc^2 x = -2 \csc x \cot x$
3.  $y = \cos 3x$ ;  $y' = -3 \sin 3x$
4.  $y = \sin \frac{x}{5}$ ;  $y' = \frac{1}{5} \cos \frac{x}{5}$
5.  $y = \tan x$ ;  $y' = \sec^2 x$
6.  $y = \sec ax$ ;  $y' = a \sec ax \tan ax$
7.  $y = \cot 4 - 3x$ ;  $y' = -3 \csc^2 4 - 3$
8.  $\frac{d}{dx} \sin^3 x = 3 \sin^2 x \cos x$
9.  $f(x) = \cos \sin^{-1} x$ ;  $f'(x) = -\sin \sin^{-1} x \cdot \frac{1}{\sqrt{1-x^2}} = -\frac{\sin \sin^{-1} x}{\sqrt{1-x^2}}$
10.  $y = \sin Ax \cos Bx$ ;  $y' = A \cos Ax \cos Bx - B \sin Ax \sin Bx$
11.  $\frac{d}{dx} \sin x^2 = 2x \cos x^2$
12.  $\frac{d}{dx} \cos^{-1} x = \frac{-1}{\sqrt{1-x^2}}$ ;  $\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}}$
13.  $y = \cos^{-1} \cos x$ ;  $y' = -\frac{1}{\sqrt{1-\cos^2 x}} \cdot (-\sin x) = \frac{\sin x}{\sin x} = 1$
14.  $\frac{d}{dx} \sin 2 \cos x = 2 \cos x \cos x - \sin 2 \sin x = 2 \cos^2 x - \sin 2 \sin x$
15.  $f(x) = \cos x \sin x$ ;  $f'(x) = -\sin x \sin x + \cos x \cos x = \cos^2 x - \sin^2 x$
16.  $f(x) = \frac{1}{\tan x} = \cot x$ ;  $f'(x) = -\csc^2 x$

23.  $\frac{d}{dx} \tan x = \sec^2 x$ ;  $\frac{d}{dx} \cot x = -\csc^2 x$
24.  $\frac{d}{dx} \sec x = \sec x \tan x$ ;  $\frac{d}{dx} \csc x = -\csc x \cot x$
25.  $\frac{d}{dx} \tan x^2 = 2x \sec^2 x^2$
26.  $\frac{d}{dx} \tan 3x = 3 \sec^2 3x$ ;  $\frac{d}{dx} \cot 3x = -3 \csc^2 3x$
27.  $\frac{d}{dt} \cos t = -\sin t$ ;  $\frac{d}{dt} \sin t = \cos t$
28.  $\frac{d}{dt} \sin t \cos t = \cos t \cos t - \sin t \sin t = \cos^2 t - \sin^2 t$
29.  $\frac{d}{dx} \frac{\sin x}{\cos x} = \frac{\cos x \cos x - \sin x (-\sin x)}{\cos^2 x} = \frac{\cos^2 x + \sin^2 x}{\cos^2 x} = \frac{1}{\cos^2 x} = \sec^2 x$
30.  $\frac{d}{dx} \frac{\cos x}{\sin x} = \frac{-\sin x \sin x - \cos x \cos x}{\sin^2 x} = \frac{-\sin^2 x - \cos^2 x}{\sin^2 x} = -\frac{1}{\sin^2 x} = -\csc^2 x$
31.  $\frac{d}{dx} x^2 \cos 3x = 2x \cos 3x - 3x^2 \sin 3x$
32.  $\frac{d}{dt} \sin t \cos t = \cos t \cos t - \sin t \sin t = \cos^2 t - \sin^2 t$
33.  $\frac{d}{dx} \sec x \tan x = \sec x \sec^2 x + \tan x \sec^2 x = \sec^3 x (1 + \tan^2 x) = \sec^3 x \sec^2 x = \sec^5 x$
34.  $\frac{d}{dx} \frac{1}{\cos x} = \frac{0 \cdot \cos x - 1 \cdot (-\sin x)}{\cos^2 x} = \frac{\sin x}{\cos^2 x} = \sec x \tan x$

17.  $u = \frac{1}{\sin^3 x} = \cos^3 x$ ;  $u' = -2 \cos x \sin^2 x = -2 \cos x \sin^2 x$

18.  $y = \sec x$ ;  $y' = \sec x \tan x$

19.  $F = \sin at \cos at$ ;  $F' = 2 \sin at \cos at$

$F' = 2a \sin at \cos at = a \sin 2at$

20.  $G = \frac{\sin a}{\cos b}$   
 $G' = \frac{a \cos b \cos a - b \sin a \sin b}{\cos^2 b}$

21.  $\frac{d}{dx} \sin 2x = 2 \cos 2x$ ;  $\frac{d}{dx} \cos 2x = -2 \sin 2x$

22.  $\frac{d}{dx} \cos^2 x = 2 \cos x (-\sin x) = -2 \sin x \cos x$

$\frac{d}{dx} \cos^{-1} x = \frac{-1}{\sqrt{1-x^2}}$

$\frac{d}{dx} \cos^{-1} x^2 = 2x \cdot \frac{-1}{\sqrt{1-x^2}} = \frac{-2x}{\sqrt{1-x^2}}$

$\frac{d}{dt}$

35.  $\frac{d}{dt} \sin \cos \tan t = \cos^2 t \sec^2 t = \frac{1}{\cos^2 t}$

36.  $f = \cos s$ ;  $f' = -\sin s$

$f' = -\sin s$

37. Differentiate both sides of  $\sin 2x = 2 \sin x \cos x$  and divide by 2 to get  $\cos 2x = \cos^2 x - \sin^2 x$ .

38. Differentiate both sides of  $\cos 2x = \cos^2 x \sin^2 x$  and divide by 2 to get  $\sin 2x = 2 \sin x \cos x$ .

39. Slope of  $y = \sin x$  at  $0$  is  $\cos 0 = 1$ . Therefore the tangent and normal lines to  $y = \sin x$  at  $0$  have equations  $y = x$  and  $y = -x$ , respectively.

40. The slope of  $y = \tan 2x$  at  $x = 0$  is  $2 \sec^2 0 = 2$ .  
Therefore the tangent and normal lines to  $y = \tan 2x$  at  $x = 0$  have equations  $y = 2x$  and  $y = -x/2$ , respectively.

41. The slope of  $y = \frac{1}{2} \cos x = 4$  at  $x = \pi/4$  is  $-\frac{1}{2} \sin x = -\frac{1}{2} \sin(\pi/4) = -\frac{1}{2}(\frac{\sqrt{2}}{2}) = -\frac{\sqrt{2}}{4}$ . Therefore the tangent and normal lines to  $y = \frac{1}{2} \cos x$  at  $x = \pi/4$  have equations  $y - \frac{1}{2} = -\frac{\sqrt{2}}{4}(x - \frac{\pi}{4})$  and  $y - \frac{1}{2} = 4(x - \frac{\pi}{4})$ , respectively.

42. The slope of  $y = \cos^2 x$  at  $x = \pi/3$  is  $2 \cos x (-\sin x) = 2(\frac{1}{2})(-\frac{\sqrt{3}}{2}) = -\sqrt{3}$ . Therefore the tangent and normal lines to  $y = \cos^2 x$  at  $x = \pi/3$  have equations  $y - \frac{1}{4} = -\sqrt{3}(x - \frac{\pi}{3})$  and  $y - \frac{1}{4} = \frac{1}{\sqrt{3}}(x - \frac{\pi}{3})$ , respectively.

43. Slope of  $y = \sin x$  at  $x = \pi/4$  is  $\cos x = \frac{\sqrt{2}}{2}$ .  
At  $x = \pi/4$  the tangent line is  $y - \frac{\sqrt{2}}{2} = \frac{\sqrt{2}}{2}(x - \frac{\pi}{4})$ .

44. For  $y = \sec x$  we have  $\frac{dy}{dx} = \sec x \tan x$ . At  $x = \pi/2$ ,  $\frac{dy}{dx}$  is undefined.

At  $x = \pi/2$  the tangent line is  $x = \pi/2$ .

45. The slope of  $y = \tan x$  at  $x = \pi/3$  is  $\sec^2 x = 4$ .  
The tangent line is  $y - \frac{\sqrt{3}}{3} = 4(x - \frac{\pi}{3})$ .

46. The slope of  $y = \tan 2x$  at  $x = \pi/4$  is  $2 \sec^2 x = 4$ .  
The tangent line is  $y - 1 = 4(x - \frac{\pi}{4})$ .

47. The slope of  $y = \sin x$  at  $x = \pi/2$  is  $\cos x = 0$ .

Thus, the normal line has slope  $\frac{90}{\pi}$  and has equation  $y - \frac{90}{\pi} = \frac{90}{\pi}(x - \frac{\pi}{2})$ .

48. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

49. The slope of  $y = \tan x$  at  $x = \pi/2$  is undefined.  
The normal line is  $x = \pi/2$ .

50. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

51. The slope of  $y = \sin x$  at  $x = \pi/2$  is  $\cos x = 0$ .  
The normal line is  $x = \pi/2$ .

52. The slope of  $y = \cos x$  at  $x = \pi/2$  is  $-\sin x = -1$ .  
The tangent line is  $y - 0 = -1(x - \frac{\pi}{2})$ .

53. The slope of  $y = \cos x$  at  $x = \pi/4$  is  $-\sin x = -\frac{\sqrt{2}}{2}$ .  
The tangent line is  $y - \frac{\sqrt{2}}{2} = -\frac{\sqrt{2}}{2}(x - \frac{\pi}{4})$ .

54. The slope of  $y = \sec x$  at  $x = \pi/4$  is  $\sec x \tan x = \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{2}}{2} = 1$ .  
The tangent line is  $y - \sqrt{2} = 1(x - \frac{\pi}{4})$ .

55. The slope of  $y = \csc x$  at  $x = \pi/2$  is  $-\csc x \cot x = -1 \cdot 0 = 0$ .  
The tangent line is  $y = 2$ .

56. The slope of  $y = \sec x$  at  $x = \pi/4$  is  $\sec x \tan x = \frac{\sqrt{2}}{2} \cdot \frac{\sqrt{2}}{2} = 1$ .  
The tangent line is  $y - \sqrt{2} = 1(x - \frac{\pi}{4})$ .

57. The slope of  $y = \csc x$  at  $x = \pi/2$  is  $-\csc x \cot x = -1 \cdot 0 = 0$ .  
The tangent line is  $y = 2$ .

58. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

59. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

60. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

61. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

62. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

63. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .


64. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

65. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

66. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

67. The slope of  $y = \tan x$  at  $x = \pi/4$  is  $\sec^2 x = 2$ .  
The tangent line is  $y - 1 = 2(x - \frac{\pi}{4})$ .

59. There are infinitely many lines through the origin that are tangent to  $y = \cos x$ . The two with largest slope are shown in the figure.



Thus neither of these functions has a horizontal tangent.

Fig. 2.5-59

The tangent to  $y = \cos x$  at  $x = a$  has equation  $y = \cos a - \sin a(x - a)$ . This line passes through the origin if  $\cos a = a \sin a$ . We use a calculator with a "solve" function to find solutions of this equation near  $a = 2$  and  $a = 6$  as suggested in the figure. The solutions are  $a = 2.798386$  and  $a = 6.121250$ . The slopes of the corresponding tangents are given by  $-\sin a$ , so they are  $0.336508$  and  $0.161228$  to six decimal places.

60.  $\frac{d}{dx} \cot x = -\csc^2 x$

61.  $\frac{d}{dx} \csc x = -\csc x \cot x$

62. a) As suggested by the figure in the problem, the square of the length of chord AP is  $1 - \cos \theta$  and the square of the length of arc AP is  $\theta^2$ . Hence

$$1 - \cos \theta < \theta^2 < 1 + \cos \theta$$

and, since squares cannot be negative, each term in the sum on the left is less than  $\frac{\theta^2}{2}$ . Therefore

$$0 < \sum_{j=1}^n \cos \theta_j < \sum_{j=1}^n \theta_j; \quad 0 < \sum_{j=1}^n \sin \theta_j < \sum_{j=1}^n \theta_j$$

Since  $\lim_{n \rightarrow \infty} \sum_{j=1}^n \theta_j = 0$ , the squeeze theorem implies that

$$\lim_{h \rightarrow 0} \frac{1 - \cos h}{h} = 0; \quad \lim_{h \rightarrow 0} \frac{\sin h}{h} = 1$$

From the first of these,  $\lim_{h \rightarrow 0} \cos h = 1$ .

- b) Using the result of (a) and the addition formulas for cosine and sine we obtain

$$\lim_{h \rightarrow 0} \frac{\cos h - \cos 0}{h - 0} = \lim_{h \rightarrow 0} \frac{-\sin h}{h} = -\lim_{h \rightarrow 0} \frac{\sin h}{h} = -1$$

$$\lim_{h \rightarrow 0} \frac{\sin h - \sin 0}{h - 0} = \lim_{h \rightarrow 0} \frac{\cos h}{1} = \lim_{h \rightarrow 0} \cos h = 1$$

This says that cosine and sine are continuous at any point 0.

### Section 2.6 Higher-Order Derivatives (page 131)

1.  $y = 2x^7 \implies y' = 14x^6$

$y'' = 84x^5$   
 $y''' = 420x^4$

$y^{(4)} = 1680x^3$

2.  $y = x^2 \implies y' = 2x$   
 $y'' = 2$

5.  $y = x^{1/3} \implies y' = \frac{1}{3}x^{-2/3}$

$y'' = -\frac{2}{9}x^{-5/3}$

$y''' = \frac{10}{27}x^{-8/3}$

$y^{(4)} = -\frac{28}{27}x^{-11/3}$

6.  $y = 2x^8 \implies y' = 16x^7$

$y'' = 112x^6$

$y''' = 672x^5$

$y^{(4)} = 3360x^4$

$y^{(5)} = 13440x^3$

$y^{(6)} = 40320x^2$

8.  $y = \frac{1}{x} \implies y' = -\frac{1}{x^2}$

$y'' = \frac{2}{x^3}$

9.  $y = \tan x \implies y' = \sec^2 x$

$y'' = 2 \sec^2 x \tan x$

10.  $y = \sec x \implies y' = \sec x \tan x$

$y'' = \sec x \tan^2 x + \sec^3 x$

11.  $y = \cos x^2 \implies y' = -2x \sin x^2$

$y'' = 2 \cos x^2 - 4x^2 \sin x^2$

12.  $y = \frac{\sin x}{\cos x} \implies y' = \frac{\cos^2 x + \sin^2 x}{\cos^2 x}$

$y'' = \frac{2 \cos x \sin x}{\cos^3 x} = \frac{2 \sin x}{\cos^2 x}$

$y''' = \frac{2 \cos x - 4 \sin^2 x}{\cos^3 x} = \frac{2(1 - 2 \sin^2 x)}{\cos^3 x}$

$y^{(4)} = \frac{2(2 \sin x \cos x + 6 \sin^2 x \cos x)}{\cos^4 x}$

$$y^{(4)} = x^4$$

$$y^{(4)} = \dots$$

$$3. \quad y = \frac{6}{x^{1/2}} \implies y' = -\frac{3}{x^{3/2}}$$

$$y^{(3)} = \frac{12}{x^{5/2}}$$

$$y^{(4)} = \frac{36}{x^{7/2}}$$

$$y^{(5)} = \frac{144}{x^{9/2}}$$

$$4. \quad y = \frac{p}{ax^2 + b} \implies y' = \frac{-2axp}{(ax^2 + b)^2}$$

$$y^{(2)} = \frac{3a^2 p}{(ax^2 + b)^3}$$

$$y^{(3)} = \frac{6a^3 p}{(ax^2 + b)^4}$$

$$x^3$$

$$13. \quad f(x) = x^2 \implies f'(x) = 2x$$

$$f''(x) = 2$$

$$f^{(3)}(x) = 0$$

Proof: (\*) is valid for  $n = 1$  (and 2, 3, 4).  
Assume  $f^{(k)}(x) = k! C x^{k-1}$  for some  $k \geq 1$ .

$$\text{Then } f^{(k+1)}(x) = k! C k x^{k-1} = (k+1)! C x^{k-1}$$

$f^{(k+1)}(x) = (k+1)! C x^{k-1}$  which is (\*) for  $n = k+1$ .  
Therefore, (\*) holds for  $n = 1, 2, 3, \dots$  by induction.



14.  $f(x) = x^{-2} = x^{-2}$

$f'(x) = -2x^{-3} = -\frac{2}{x^3}$   
 $f''(x) = \frac{6}{x^4} = \frac{6x^{-4}}$

$f'''(x) = -\frac{24}{x^5} = -24x^{-5}$

Conjecture:

$f^{(n)}(x) = (-1)^n \frac{n!}{x^{n+2}}$  for  $n = 1, 2, 3, \dots$

Proof: Evidently, the above formula holds for  $n = 1, 2$

and 3. Assume it holds for  $n = k$ ,

i.e.,  $f^{(k)}(x) = (-1)^k \frac{k!}{x^{k+2}}$ . Then

$$f^{(k+1)}(x) = \frac{d}{dx} f^{(k)}(x)$$

$$= (-1)^k \frac{k!}{x^{k+2}} \cdot (-k-2) x^{-k-3}$$

$$= (-1)^{k+1} \frac{(k+1)!}{x^{k+3}}$$

Thus, the formula is also true for  $n = k + 1$ . Hence it is true for  $n = 1, 2, 3, \dots$  by induction.

$$\frac{1}{x^2}$$

15.  $f(x) = x^{-2} = x^{-2}$   
 $f'(x) = -2x^{-3} = -\frac{2}{x^3}$

$f''(x) = \frac{6}{x^4} = \frac{6x^{-4}}$

$f'''(x) = -\frac{24}{x^5} = -24x^{-5}$

$f^{(n)}(x) = (-1)^n \frac{n!}{x^{n+2}}$

Proof: (\*) holds for  $n = 1, 2, 3$ .

Assume  $f^{(k)}(x) = (-1)^k \frac{k!}{x^{k+2}}$  (i.e., (\*) holds for  $n = k$ )

Then  $f^{(k+1)}(x) = \frac{d}{dx} f^{(k)}(x)$   
 $= (-1)^k \frac{k!}{x^{k+2}} \cdot (-k-2) x^{-k-3}$   
 $= (-1)^{k+1} \frac{(k+1)!}{x^{k+3}}$

Thus (\*) holds for  $n = k + 1$  if it holds for  $k$ .

Therefore, (\*) holds for  $n = 1, 2, 3, \dots$  by induction.

16.  $f(x) = x^{-1} = x^{-1}$

$f'(x) = -x^{-2} = -\frac{1}{x^2}$

$f''(x) = \frac{2}{x^3} = \frac{2x^{-3}}$   
 $f'''(x) = -\frac{6}{x^4} = -6x^{-4}$

Thus, the formula is also true for  $n = k + 1$ . Hence, it is true for  $n = 2$  by induction.

17.  $f(x) = \frac{1}{a + bx} = (a + bx)^{-1}$

$f'(x) = -b(a + bx)^{-2} = -\frac{b}{(a + bx)^2}$

$f''(x) = \frac{2b^2}{(a + bx)^3} = \frac{2b^2}{(a + bx)^3}$   
 $f'''(x) = -\frac{6b^3}{(a + bx)^4} = -\frac{6b^3}{(a + bx)^4}$

Guess:  $f^{(n)}(x) = (-1)^n \frac{n! b^n}{(a + bx)^{n+1}}$

Proof: (\*) holds for  $n = 1, 2, 3$

Assume (\*) holds for  $n = k$ :

$f^{(k)}(x) = (-1)^k \frac{k! b^k}{(a + bx)^{k+1}}$

Then

$f^{(k+1)}(x) = \frac{d}{dx} f^{(k)}(x)$

$= (-1)^k \frac{k! b^k}{(a + bx)^{k+1}} \cdot (-k-1) b (a + bx)^{-k-2}$   
 $= (-1)^{k+1} \frac{(k+1)! b^{k+1}}{(a + bx)^{k+2}}$

$= (-1)^{k+1} \frac{(k+1)! b^{k+1}}{(a + bx)^{k+2}}$

So (\*) holds for  $n = k + 1$  if it holds for  $n = k$ . Therefore,

(\*) holds for  $n = 1, 2, 3, 4, \dots$  by induction.

18.  $f(x) = x^{-3} = x^{-3}$

$f'(x) = -3x^{-4} = -\frac{3}{x^4}$

$f''(x) = \frac{12}{x^5} = \frac{12x^{-5}}$   
 $f'''(x) = -\frac{60}{x^6} = -60x^{-6}$

$f^{(4)}(x) = \frac{720}{x^7} = \frac{720x^{-7}}$

Conjecture:

$f^{(n)}(x) = (-1)^n \frac{n! \cdot 3^n}{x^{n+3}}$  for  $n = 1, 2, 3, \dots$

$n = 2$ .

Proof: Evidently, the above formula holds for  $n = 2$  and 3.

Assume that it holds for  $n = k$ , i.e.

$f^{(k)}(x) = (-1)^k \frac{k! \cdot 3^k}{x^{k+3}}$

Then,

$\frac{d}{dx}$

$f^{(k+1)}(x) = \frac{d}{dx} f^{(k)}(x)$



Differentiating any of these four formulas produces the one for the next higher value of  $n$ , so induction confirms the overall formula.

$$\begin{aligned} 20. \quad & f^{(n)}(x) = D^n x \cos x \\ & f^{(0)}(x) = D^0 \cos x = x \sin x \\ & f^{(1)}(x) = D^1 \cos x = -x \cos x \end{aligned}$$

$$\begin{aligned} & f^{(2)}(x) = D^2 \cos x = -\cos x \\ & f^{(3)}(x) = D^3 \cos x = x \sin x \end{aligned}$$

This suggests the formula (for  $k \geq 0; 1; 2; \dots$ )

$$\begin{aligned} f^{(n)}(x) &= D^n x \cos x && \text{if } n = 4k \\ &= D^n x \sin x && \text{if } n = 4k + 1 \\ &= -D^n x \cos x && \text{if } n = 4k + 2 \\ &= -D^n x \sin x && \text{if } n = 4k + 3 \end{aligned}$$

Differentiating any of these four formulas produces the one for the next higher value of  $n$ , so induction confirms the overall formula.

$$\begin{aligned} 21. \quad & f(x) = D x \sin ax \\ & f^{(0)}(x) = D \sin ax = a \cos ax \\ & f^{(1)}(x) = D^2 \sin ax = -a^2 \sin ax \\ & f^{(2)}(x) = D^3 \sin ax = -a^3 \cos ax \\ & f^{(3)}(x) = D^4 \sin ax = a^4 \sin ax \end{aligned}$$

This suggests the formula

$$\begin{aligned} f^{(n)}(x) &= a^n \cos ax && \text{if } n = 4k \\ &= a^n \sin ax && \text{if } n = 4k + 1 \\ &= -a^n \cos ax && \text{if } n = 4k + 2 \\ &= -a^n \sin ax && \text{if } n = 4k + 3 \end{aligned}$$

for  $k \geq 0; 1; 2; \dots$ . Differentiating any of these four formulas produces the one for the next higher value of  $n$ , so induction confirms the overall formula.

$$22. f(x) = D^n x \sin x. \text{ Recall that } dx \sin x = \cos x, \text{ so}$$

The pattern suggests that

$$f^{(n)}(x) = D^n x \sin x = (-1)^{n/2} x \sin x \text{ if } n \text{ is even}$$

Differentiating this formula leads to the same formula with  $n$  replaced by  $n - 1$  so the formula is valid for all  $n \geq 1$  by induction.

$$\begin{aligned} 23. \quad & f(x) = D^n \frac{1}{3x} \\ & f^{(0)}(x) = D^0 \frac{1}{3x} = \frac{1}{3x} \\ & f^{(1)}(x) = D^1 \frac{1}{3x} = -\frac{1}{3x^2} \\ & f^{(2)}(x) = D^2 \frac{1}{3x} = \frac{2}{3x^3} \\ & f^{(3)}(x) = D^3 \frac{1}{3x} = -\frac{6}{3x^4} \\ & f^{(4)}(x) = D^4 \frac{1}{3x} = \frac{24}{3x^5} \end{aligned}$$

$$\text{Guess: } f^{(n)}(x) = D^n \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2n-1)}{2^n} \frac{1}{3x^{2n}}$$

Proof: (\*) is valid for  $n = 2; 3; 4$ ; (but not  $n = 1$ ) Assume (\*) holds for  $n = k$  for some integer  $k \geq 2$

$$\begin{aligned} \text{i.e., } f^{(k)}(x) &= D^k \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k-1)}{2^k} \frac{1}{3x^{2k}} \\ \text{Then } f^{(k+1)}(x) &= D^{k+1} \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k-1)}{2^k} \frac{1}{3x^{2k}} \end{aligned}$$

$$\begin{aligned} &= \frac{2k-1}{2} \cdot \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k-1)}{2^k} \frac{1}{3x^{2k}} - \frac{2k}{3x^{2k+1}} \\ &= \frac{2k-1}{2} \cdot \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k-1)}{2^k} \frac{1}{3x^{2k}} - \frac{2k}{3x^{2k+1}} \\ &= \frac{2k-1}{2} \cdot \frac{1 \cdot 3 \cdot 5 \cdot \dots \cdot (2k-1)}{2^k} \frac{1}{3x^{2k}} - \frac{2k}{3x^{2k+1}} \end{aligned}$$

Thus (\*) holds for  $n = k + 1$  if it holds for  $n = k$ . Therefore, (\*) holds for  $n = 2; 3; 4; \dots$  by induction.

$$\begin{aligned} 24. \quad & \text{If } y = D^n \tan kx, \text{ then } y^{(0)} = D^n \tan kx \text{ and} \\ & y^{(1)} = D^{n+1} \tan kx = D^n \sec^2 kx \\ & y^{(2)} = D^{n+2} \tan kx = D^n \sec^2 kx \tan kx \\ & y^{(3)} = D^{n+3} \tan kx = D^n \sec^2 kx \tan^2 kx \end{aligned}$$

$$f^{(0)}(x) = |x|^2 \operatorname{sgn} x:$$

If  $x \neq 0$  we have

$$\frac{d}{dx} |x|^2 \operatorname{sgn} x = 2|x| \operatorname{sgn} x = 2|x|$$

Thus we can calculate successive derivatives of  $f$  using

the product rule where necessary, but will get only one nonzero term in each case:

$$\begin{aligned} f^{(0)}(x) &= |x|^2 \operatorname{sgn} x \\ f^{(1)}(x) &= 2|x| \operatorname{sgn} x \\ f^{(2)}(x) &= 2 \operatorname{sgn} x \\ f^{(3)}(x) &= 2 \delta(x) \\ f^{(4)}(x) &= 0 \end{aligned}$$

25. If  $y = \sec kx$ , then  $y' = k \sec kx \tan kx$  and

$$\begin{aligned} y'' &= k^2 \sec^2 kx \tan^2 kx + k^3 \sec^3 kx \\ &= k^2 y^2 \sec^2 kx + k^3 y^3 \end{aligned}$$

$$f^{(n)}(x) = \frac{1}{k^n} a^n \cos ax \quad \text{if } n = 2k + 1$$

$$= \frac{1}{k^n} a^n \sin ax \quad \text{if } n = 2k$$

for  $k = 0, 1, 2, \dots$ : Proof: The formula works for  $k = 0$  ( $n = 0, 1$ ) and  $n = 2, 3$  ( $k = 1, 2$ ):

$$\begin{aligned} f^{(0)}(x) &= f(x) = a^0 \cos ax \\ f^{(1)}(x) &= f'(x) = -a^1 \sin ax \\ f^{(2)}(x) &= f''(x) = -a^2 \cos ax \\ f^{(3)}(x) &= f'''(x) = a^3 \sin ax \end{aligned}$$

Now assume the formula holds for some  $k \geq 0$ .  
If  $n = 2k + 1$ , then

$$f^{(n)}(x) = \frac{d}{dx} f^{(n-1)}(x) = \frac{d}{dx} [(-1)^k (2k)! a^{2k} \sin(ax) + b] \\ = (-1)^{k+1} (2k+1)! a^{2k+1} \cos(ax) + 0$$

and if  $n = 2k + 3$ , then

$$f^{(n)}(x) = \frac{d}{dx} f^{(n-1)}(x) = \frac{d}{dx} [(-1)^{k+1} (2k+2)! a^{2k+2} \sin(ax) + b] \\ = (-1)^{k+2} (2k+3)! a^{2k+3} \cos(ax) + 0$$

Thus the formula also holds for  $k \geq 1$ . Therefore it holds for all positive integers  $k$  by induction.

27. If  $y = \tan x$ , then

$$y' = \sec^2 x = 1 + \tan^2 x = 1 + y^2 = P_2(y);$$

where  $P_2$  is a polynomial of degree 2. Assume that  $y^{(n)}$

$y^{(n)} = P_n(y)$  where  $P_n$  is a polynomial of degree

$n - 1$ . The derivative of any polynomial is a polynomial of one lower degree, so

$$y^{(n+1)} = \frac{d}{dx} P_n(y) = P_n'(y) y' = P_n'(y) (1 + y^2) = P_{n+1}(y)$$

$y'$ ; a polynomial of degree  $n - 2$ . By induction,

$y^{(n)} = dx^n \tan x = P_n(y)$ , a polynomial of degree

$n - 1$  in  $\tan x$ .

$$f^{(n)}(g) = f^{(n-1)}(g) f'(g) = f^{(n-1)}(g) f'(g) = f^{(n-1)}(g) f'(g) = \dots$$

29.  $f^{(3)}(g) = \frac{d}{dx} f^{(2)}(g)$

$$= \frac{d}{dx} [2f''(g)g' + f'(g)g'']$$

$$= 2f'''(g)g'^2 + 2f''(g)g''g' + f''(g)g''' + f'(g)g''^2$$

### Section 2.7 Using Differentials and Derivatives (page 137)

1.  $y = \frac{1}{x^2} \Rightarrow \frac{dy}{dx} = -\frac{2}{x^3} = -\frac{2}{2^3} = -\frac{2}{8} = -\frac{1}{4}$

If  $x = 2$ , then  $y = \frac{1}{4}$

2.  $f(x) = \frac{3}{2}x^2 - \frac{1}{3}x^3 \Rightarrow f'(x) = 3x - x^2$

$f'(1) = 3(1) - (1)^2 = 2$

3.  $h(t) = 4 \sin^{-1} t \Rightarrow \frac{dh}{dt} = \frac{4}{\sqrt{1-t^2}}$

4.  $u = \frac{1}{4} \sec^2 s \Rightarrow \frac{du}{ds} = \frac{1}{2} \sec^2 s$

If  $s = \frac{\pi}{4}$ , then  $u = \frac{1}{2}$

5. If  $y = x^2$ , then  $dy = 2x dx$ . If  $dx = 2 = 100/x$ , then  $y = 4 = 100/x^2$

6. If  $y = 1/x$ , then  $dy = -1/x^2 dx$

$dx = 2 = 100/x$ , then  $y = 2 = 100/x^2$

7. If  $y = 1/x^2$ , then  $dy = -2/x^3 dx$

$dx = 2 = 100/x$ , then  $y = 4 = 100/x^2$

8. If  $y = x^3$ , then  $dy = 3x^2 dx$

$dx = 6 = 100/x$ , then  $y = 6 = 100/x^2$

9. If  $y = x^p$ , then  $dy = px^{p-1} dx$

10. If  $y = x^2$ , then  $dy = 2x dx$

11. If  $V = \frac{4}{3}r^3$ , then  $dV = 4r^2 dr$



15. The diameter  $D$  and area  $A$  of a circle are related by  $A = \frac{\pi}{4} D^2$ . The rate of change of diameter with respect to area is  $\frac{dD}{dA} = \frac{1}{2\pi A}$  units per square unit.

16. Since  $A = \frac{\pi}{4} D^2 = 4$ , the rate of change of area with respect to diameter is  $\frac{dA}{dD} = \frac{\pi}{2} D = 2\pi$  square units per unit.

17. Rate of change of  $V = \frac{4}{3}\pi r^3$  with respect to radius  $r$  is  $\frac{dV}{dr} = 4\pi r^2$ . When  $r = 2$  m, this rate of change is  $16\pi$  m<sup>3</sup>/m.

18. Let  $A$  be the area of a square,  $s$  be its side length and  $L$  be its diagonal. Then,  $A = s^2$  and  $L = \sqrt{2}s$ . Thus, the rate of change of the area of a square with respect to its diagonal  $L$  is  $\frac{dA}{dL} = \frac{1}{2}L$ .

19. If the radius of the circle is  $r$  then  $C = 2\pi r$  and  $A = \pi r^2$ .

Rate of change of  $C$  with respect to  $A$  is  $\frac{dC}{dA} = \frac{1}{\pi r}$ .

20. Let  $s$  be the side length and  $V$  be the volume of a cube.

Then  $V = s^3$  and  $\frac{dV}{ds} = 3s^2$ . Hence,

the rate of change of the side length of a cube with respect to its volume  $V$  is  $\frac{1}{3V^{2/3}}$ .

21. Volume in tank is  $V = 350.20 - \frac{1}{2}t^2$  L at  $t$  min.

a) At  $t = 5$ , water volume is changing at rate

$$\frac{dV}{dt} = -t = -5 \text{ L/min}$$

Water is draining out at 5 L/min at that time.

At  $t = 15$ , water volume is changing at rate

$$\frac{dV}{dt} = -t = -15 \text{ L/min}$$

Water is draining out at 15 L/min at that time.

b) Average rate of change between  $t = 5$  and  $t = 15$  is

The flow rate will increase by 10% if the radius is increased by about 2.5%.

23.  $F = kr^2$  implies that  $\frac{dF}{dr} = 2kr$ . Since  $\frac{dF}{dr} = 1$  pound/mi when  $r = 4000$  mi, we have

$2k(4000) = 1$ . If  $r = 8000$ , we have

$\frac{dF}{dr} = 2k(8000) = 2$ . At  $r = 8000$  mi  $F$  decreases with respect to  $r$  at a rate of 1/8 pounds/mi.

24. If price =  $p$ , then revenue is  $R = 4000p - 10p^2$ .  
 a) Sensitivity of  $R$  to  $p$  is  $\frac{dR}{dp} = 4000 - 20p$ . If  $p = 100, 200,$  and  $300$ , this sensitivity is 2,000 \$/\$, 0 \$/\$, and -2,000 \$/\$ respectively.

b) The distributor should charge \$200. This maximizes the revenue.

25. Cost is  $C = 400x + 0.5x^2$  if  $x$  units are manufactured.

a) Marginal cost if  $x = 100$  is

$$C' = 400 + 100 = 500$$

b)  $C'(101) = 501$ ;  $C'(100) = 500$ ;  $C'(99) = 499$  which is approximately  $C'(100)$ .

26. Daily profit if production is  $x$  sheets per day is  $P = 8x - 0.005x^2$  where

$$P = 8x - 0.005x^2$$

a) Marginal profit  $P' = 8 - 0.01x$ . This is positive if  $x < 800$  and negative if  $x > 800$ .

b) To maximize daily profit, production should be 800 sheets/day.

27.  $C = 4n^2 - 100n$

$$\frac{dC}{dn} = 8n - 100$$

(a)  $n = 100$ ;  $\frac{dC}{dn} = 8(100) - 100 = 700$ . Thus, the marginal cost of  $n$  production is \$700.

(b)  $n = 300$ ;  $\frac{dC}{dn} = 8(300) - 100 = 2300$ . Thus, the marginal cost of production is approximately \$2300.

28. Daily profit  $P = 13x - C(x) = 13x - 10x - \frac{20}{1000}x^2$

$$\frac{V_{15} - V_5}{15 - 5} = \frac{350 - 225}{10} = 7,000$$

The average rate of draining is 7,000 L/min over that interval.

22. Flow rate  $F = kr^4$ , so  $F = 4kr^3 \cdot r$ . If  $F = 10$ , then

$$r = \frac{F}{40kr^3} = \frac{kr^4}{40kr^3} = 0.25r$$

$$D_x(3x^2 - 20x + 1000)$$

Graph of  $P$  is a parabola opening downward.  $P$  will be maximum where the slope is zero:

$$0 = \frac{dP}{dx} = 6x - 20 \quad \text{so } x = 1500$$

Should extract 1500 tonnes of ore per day to maximize profit.



29. One of the components comprising  $C(x)$  is usually a fixed cost,  $S$ , for setting up the manufacturing operation. On a per item basis, this fixed cost  $S/x$ , decreases as the number  $x$  of items produced increases, especially when  $x$  is small. However, for large  $x$  other components of the total cost may increase on a per unit basis, for instance labour costs when overtime is required or maintenance costs for machinery when it is over used. Let the average cost be  $A(x) = \frac{C(x)}{x}$ . The minimal average cost occurs at point where the graph of  $A(x)$  has a horizontal tangent:

$$\frac{dA}{dx} = \frac{xC'(x) - C(x)}{x^2} = 0$$

Hence,  $xC'(x) - C(x) = 0$  or  $C'(x) = \frac{C(x)}{x}$ . Thus the marginal cost  $C'(x)$  equals the average cost at the minimizing value of  $x$ .

30. If  $y = Cp^r$ , then the elasticity of  $y$  is

$$\frac{p}{y} \frac{dy}{dp} = \frac{p}{Cp^r} \cdot rCp^{r-1} = \frac{r}{p} Cp^r = r$$

**Section 2.8 The Mean-Value Theorem (page 144)**

1.  $f(x) = ax^2 + b$ ;  $f'(x) = 2ax$

$$\frac{f(b) - f(a)}{b - a} = \frac{ab^2 + b - (a^2 + b)}{b - a} = \frac{ab^2 - a^2}{b - a} = \frac{a(b^2 - a^2)}{b - a} = a(b + a)$$

2. If  $f(x) = x^2$ , and  $f'(x) = 2x$  then

$$\frac{f(2) - f(1)}{2 - 1} = \frac{4 - 1}{1} = 3 = 2 \cdot \frac{1 + 2}{2} = 2 \cdot \frac{3}{2} = 3$$

4. If  $f(x) = \cos x$ ,  $f'(x) = -\sin x$ . By the MVT, if  $x > 0$ , then

$$\frac{f(x) - f(0)}{x - 0} = \frac{\cos x - 1}{x} = -\sin c$$

for some  $c > 0$ , so  $\frac{\cos x - 1}{x} > -1$ . Thus  $\cos x > 1 - x$  for  $x > 0$ . Since both sides of the inequality are even functions, it must hold for  $x < 0$  as well.

5. Let  $f(x) = \tan x$ . If  $0 < x < \pi/2$ , then by the MVT

$$\frac{f(x) - f(0)}{x - 0} = \frac{\tan x}{x} = \sec^2 c$$

for some  $c$  in  $(0, \pi/2)$ . Thus  $\tan x > x$  since  $\sec^2 c > 1$ .

6. Let  $f(x) = 1 - Cx^r$  where  $r > 1$ .

$$\frac{f(x) - f(0)}{x - 0} = \frac{1 - Cx^r - 1}{x} = -Cx^{r-1}$$

If  $-1 < x < 0$  then  $f(x) < f(0)$ ; if  $x > 0$ , then  $f(x) > f(0)$ . Thus  $f(x) > f(0) + Cx^r$  if  $-1 < x < 0$  or  $x > 0$ . Thus  $1 - Cx^r > 1 - Cx^r$  if  $-1 < x < 0$  or  $x > 0$ .

7. Let  $f(x) = 1 - Cx^r$  where  $0 < r < 1$ . Thus,

$\frac{f(x) - f(0)}{x - 0} = \frac{1 - Cx^r - 1}{x} = -Cx^{r-1}$ . By the Mean-Value Theorem, for  $x > 0$ ,

$$\frac{f(x) - f(0)}{x - 0} = \frac{1 - Cx^r - 1}{x} = -Cx^{r-1} = -C c^{r-1}$$

for some  $c$  between  $0$  and  $x$ . Thus,  $1 - Cx^r > 1 - Cc^r$ . If  $-1 < x < 0$ , then  $c < 0$  and  $0 < 1 - Cc^r < 1$ . Hence  $1 - Cx^r > 1 - Cc^r$  since  $r < 1$ ;

$$r x \cdot 1 - Cx^{r-1} < r x \quad \text{since } x < 0$$

Hence,  $1 - Cx^r < 1 - Cc^r$ . If  $x > 0$ , then

$$\frac{f(x) - f(0)}{x - 0} = \frac{1 - Cx^r - 1}{x} = -Cx^{r-1} < -Cx^{r-1}$$

Hence,  $1 - Cx^r < 1 - Cc^r$  in this case also.

Hence,  $1 - Cx^r < 1 - Cc^r$  for either  $-1 < x < 0$  or  $x > 0$ .

8. If  $f(x) = x^3$ ,  $f'(x) = 3x^2$ . The critical points of  $f$  are  $x = 0$ .  $f$  is increasing on  $(-\infty, 0)$  and  $(0, \infty)$  where  $f'(x) > 0$ , and is decreasing on  $(-\infty, 0)$  and  $(0, \infty)$  where  $f'(x) < 0$ .

where  $c \neq 2$  lies between 1 and 2.

3.  $f(x) = x^3 - 3cx^2 + 3ax - a$ ,  $f'(x) = 3x^2 - 6cx + 3a$ ,  $a \neq 2$ ,  $b \neq 2$

$f(b) = f(a) = f(2) = f(-2)$

$$\frac{b^3 - 3cb^2 + 3ab - a}{b - a} = \frac{8 - 6c + 3a - a}{8 - 4} = \frac{4 - 2c + 2a}{4} = \frac{2 - c + a}{2}$$

$f'(c) = 3c^2 - 6c^2 + 3a = -3c^2 + 3a = 0$   
 $3c^2 - 3 = 0 \Rightarrow c^2 = 1 \Rightarrow c = \pm 1$   
 (Both points will be in  $(-2, 2)$ .)

where  $f'(x) < 0$ .

9. If  $f(x) = x^4 - 4x^3$ , then  $f'(x) = 4x^3 - 12x^2 = 4x^2(x - 3)$ . The critical point of  $f$  is  $x = 3$ .  $f$  is increasing on  $(0, 3)$  and decreasing on  $(3, \infty)$ .

10. If  $y = 1 - x^5$ , then  $y' = -5x^4 < 0$  for all  $x$ . Thus  $y$  has no critical points and is decreasing on the whole real line.

11. If  $y = x^3 - 6x^2 + 12x - 3$ , then  $y' = 3x^2 - 12x + 12 = 3(x^2 - 4) = 3(x - 2)(x + 2)$ . The critical points of  $y$  are  $x = 2$  and  $x = -2$ .  $y$  is increasing on  $(-2, 2)$  and decreasing on  $(-\infty, -2) \cup (2, \infty)$  where  $y' > 0$ , and is decreasing on  $(-\infty, -2) \cup (2, \infty)$  where  $y' < 0$ .

12. If  $f(x) = 2x^2 - 2x + 2$  then  $f'(x) = 4x - 2$ .  
Evidently,  $f'(x) > 0$  if  $x > 1/2$  and  $f'(x) < 0$  if  $x < 1/2$ .  
Therefore,  $f$  is increasing on  $(1/2, \infty)$  and decreasing on  $(-\infty, 1/2)$ .

13.  $f(x) = x^3 - 4x + 1$

$$f'(x) = 3x^2 - 4$$

$$f'(x) > 0 \text{ if } x > \frac{2}{\sqrt{3}}$$

$$f'(x) < 0 \text{ if } x < \frac{2}{\sqrt{3}}$$

$f$  is increasing on  $(\frac{2}{\sqrt{3}}, \infty)$  and  $(-\infty, \frac{2}{\sqrt{3}})$ ;  
 $f$  is decreasing on  $(\frac{2}{\sqrt{3}}, \infty)$ .

14. If  $f(x) = 3x^3 - 4x + 1$ , then  $f'(x) = 9x^2 - 4$ . Since  $f'(x) > 0$  for all real  $x$ , hence  $f(x)$  is increasing on the whole real line, i.e., on  $(-\infty, \infty)$ .

15.  $f(x) = x^2 - 4x + 2$

$$f'(x) = 2x - 4$$

$$f'(x) > 0 \text{ if } x > 2 \text{ or } 2 < x < 0$$

$$f'(x) < 0 \text{ if } x < 2 \text{ or } 0 < x < 2$$

$f$  is increasing on  $(-\infty, 2)$  and  $(0, \infty)$ ;

$f$  is decreasing on  $(2, \infty)$  and  $(0, 2)$ .

16. If  $f(x) = \frac{1}{x^2} - 1$  then  $f'(x) = -\frac{2x}{x^4} = -\frac{2}{x^3}$ . Evidently,

$$f'(x) > 0 \text{ if } x < 0 \text{ and } f'(x) < 0 \text{ if } x > 0. \text{ Therefore, } f$$

is increasing on  $(-\infty, 0)$  and decreasing on  $(0, \infty)$ .

17.  $f(x) = x^3 - 5x^2$

$$f'(x) = 3x^2 - 10x$$

$$3x^2 - 10x = x(3x - 10)$$

$$= 5x^2 - 3x$$

20. If  $f(x) = 2 \sin x$ , then  $f'(x) = 2 \cos x > 0$  if  $\cos x > 0$ , i.e.,  $x \in (-\frac{\pi}{2}, \frac{\pi}{2})$ . Thus  $f$  is increasing on the intervals  $(-\frac{\pi}{2} + 2n\pi, \frac{\pi}{2} + 2n\pi)$  where  $n$  is any integer.

21.  $f(x) = x^3$  is increasing on  $(-\infty, \infty)$  because  $f'(x) = 3x^2 \geq 0$ .

18. If  $f(x) = 2 \sin x$ , then  $f'(x) = 2 \cos x > 0$  if  $\cos x > 0$ , i.e.,  $x \in (-\frac{\pi}{2}, \frac{\pi}{2})$ . Thus  $f$  is increasing on the intervals  $(-\frac{\pi}{2} + 2n\pi, \frac{\pi}{2} + 2n\pi)$  where  $n$  is any integer.  $f$  is decreasing on  $(\frac{\pi}{2} + 2n\pi, \frac{3\pi}{2} + 2n\pi)$ .

19. If  $f(x) = x \cos x$ , then  $f'(x) = \cos x - x \sin x$ .  $f'(x) > 0$  only at isolated points  $x = 0, \pm 3, \pm 5, \dots$ . Hence  $f$  is increasing everywhere.

22. There is no guarantee that the MVT applications for  $f$  and  $g$  yield the same  $c$ .

23. CPs  $x = 0.535898$  and  $x = 7.464102$

24. CPs  $x = 1.366025$  and  $x = 0.366025$

25. CPs  $x = 0.518784$  and  $x = 0$

26. CP  $x = 0.521350$

27. If  $x_1 < x_2 < \dots < x_n$  belong to  $I$ , and  $f(x_i) = 0$ ,  $f'(x_i) = 1/n$ , then there exists  $y_i \in (x_i, x_{i+1})$  such that  $f'(y_i) = 1/n$  by MVT.

28. For  $x \neq 0$ , we have  $f(x) = 2x \sin \frac{1}{x} - \cos \frac{1}{x}$

$$\text{which has no limit as } x \rightarrow 0. \text{ However, } \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{2h \sin \frac{1}{h} - \cos \frac{1}{h} - (-1)}{h} = \lim_{h \rightarrow 0} \frac{2h \sin \frac{1}{h} - \cos \frac{1}{h} + 1}{h}$$

does exist even though  $f$  cannot be continuous at 0.

29. If  $f$  exists on  $[a, b]$  and  $f(a) > f(b)$ , let us assume,

without loss of generality, that  $f(a) > k > f(b)$ . If

$g(x) = f(x) - kx$  on  $[a, b]$ , then  $g$  is continuous on  $[a, b]$  because  $f$ , having a derivative, must be continuous there. By the Max-Min Theorem,  $g$  must have a maximum value (and a minimum value) on that interval. Suppose the maximum value occurs at  $c$ . Since  $f'(a) > 0$  we must have

$c > a$ ; since  $f'(b) < 0$  we must have  $c < b$ . By Theorem 14, we must have  $f'(c) = 0$  and so  $f'(c) = k$ . Thus  $f$

takes on the (arbitrary) intermediate value  $k$ .

$$30. f(x) = \begin{cases} 0 & \text{if } x = 0 \\ x^2 \sin \frac{1}{x} & \text{if } x \neq 0 \end{cases}$$

$$a) f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{h^2 \sin \frac{1}{h}}{h} = \lim_{h \rightarrow 0} h \sin \frac{1}{h} = 0$$

$$b) \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h^2} = \lim_{h \rightarrow 0} \frac{h^2 \sin \frac{1}{h}}{h^2} = \lim_{h \rightarrow 0} \sin \frac{1}{h} = \text{does not exist}$$

vals containing the origin.

$$\lim_{h \rightarrow 0} \frac{2h \sin 1 - h}{h^2} = \frac{h}{h} = 1$$

because  $2h \sin 1 - h = h(2 \sin 1 - 1) \neq 0$  as  $h \neq 0$ .

b) For  $x \neq 0$ , we have

$$f'(x) = \frac{2x \sin 1 - x^2}{\cos 1 - x^2}$$

There are numbers  $x$  arbitrarily close to 0 where  $f'(x) < 0$ ; namely, the numbers  $x = \frac{1}{2n\pi}$ , where  $n = 1, 2, 3, \dots$ . Since  $f'(x)$  is continuous at every  $x \neq 0$ , it is negative in a small interval about every such number. Thus  $f$  cannot be increasing on any interval containing  $x = 0$ .

31. Let  $a, b$ , and  $c$  be three points in  $I$  where  $f$  vanishes; that is,  $f(a) = f(b) = f(c) = 0$ . Suppose  $a < b < c$ . By the Mean-Value Theorem, there exist points  $r$  in  $(a, b)$  and  $s$  in  $(b, c)$  such that  $f'(r) = f'(s) = 0$ . By the Mean-Value Theorem applied to  $f'$  on  $(r, s)$ , there is some point  $t$  in  $(r, s)$  (and therefore in  $I$ ) such that  $f''(t) = 0$ .

32. If  $f^{(n)}$  exists on interval  $I$  and  $f$  vanishes at  $n + 1$  distinct points of  $I$ , then  $f^{(n)}$  vanishes at at least one point of  $I$ .

Proof: True for  $n = 2$  by Exercise 8.

Assume true for  $n = k$ . (Induction hypothesis)

Suppose  $n = k + 1$ , i.e.,  $f$  vanishes at  $k + 2$  points of  $I$  and  $f^{(k)}$  exists.

By Exercise 7,  $f^{(k)}$  vanishes at  $k + 1$  points of  $I$ .

By the induction hypothesis,  $f^{(k)}$  vanishes at a

point of  $I$  so the statement is true for  $n = k + 1$ . Therefore the statement is true for all  $n \geq 1$  by induction. (case  $n = 1$  is just MVT.)

33. Given that  $f'(0) = 1$  and  $f''(1) = 0$ :

a) By MVT,

$$f'(a) = \frac{f(2) - f(0)}{2 - 0} = \frac{1 - 0}{2} = \frac{1}{2}$$

for some  $a$  in  $(0; 2)$ .

b) By MVT, for some  $r$  in  $(0; 1)$ ,

$$f'(r) = \frac{f(1) - f(0)}{1 - 0} = \frac{0 - 0}{1} = 0$$

Also, for some  $s$  in  $(1; 2)$ ,

$$f'(s) = \frac{f(2) - f(1)}{2 - 1} = \frac{1 - 0}{1} = 1$$

Then, by MVT applied to  $f'$  on the interval  $[r; s]$ , for some  $b$  in  $(r; s)$ ,

$$f''(b) = \frac{f'(s) - f'(r)}{s - r} = \frac{1 - 0}{s - r}$$

$$f''(b) = \frac{1}{s - r} > \frac{1}{2}$$

since  $s - r < 2$ .

- c) Since  $f''(x)$  exists on  $(0; 2)$ , therefore  $f''(x)$  is continuous there. Since  $f'(0) = 1$  and  $f'(1) = 0$ , and since

3.  $x^2 - xy + y^3$   
Differentiate with respect to  $x$ :  
 $2x - y + 3y^2 = 0$

4.  $x^3 - y^2 + 3xy^5$   
 $3x^2 - 2y + 3y^5 = 0$   
 $3x^2 - y + 3y^5 = 0$

5.  $x^2 - y^3 + 2xy$   
 $2x - 3y^2 + 2y = 0$   
 $2x - 3y^2 + 2y = 0$

6.  $x^2 - 4y + 1/y$   
 $2x - 4 + 1/y^2 = 0$ , so  $y = 4.1 - y$

7.  $x - y + x^2 - y^2$   
Thus  $xy - y^2 = x^3 - x^2y - xy^2$ , or  $x^3 - x^2y - xy^2 = 0$   
Differentiate with respect to  $x$ :

$$3x^2 - 2xy - x^2 = 4y^2 = 0$$

8.  $x^p - y^q$   
 $p x^{p-1} - q y^{q-1} = 0$

$$p x^{p-1} - q y^{q-1} = 0$$

$$y = \frac{p x^{p-1}}{q}$$

9.  $2x^2 - 3y^2 + 4xy$   
 $4x - 6y + 4y = 0$   
At  $(1; 1)$ :  $4 - 6 + 4 = 2$

$0 < \frac{1}{7} < 1$ , the Intermediate-Value Theorem

assures us that  $f^{-1}(c) = \frac{1}{7}$  for some  $c$  between  $r$  and  $s$ .

**Section 2.9 Implicit Differentiation (page 149)**

1.  $xy = x^2 + y^2$   
 Differentiate with respect to  $x$ :  
 $y + xy' = 2x + 2yy'$

Thus  $y' = \frac{2x - y}{y - 2y^2}$

$2x^3 + y^3 = 1$

$\frac{x^2}{y}$

$3x^2 + 3y^2 = 0$ , so  $y' = -\frac{x^2}{y^2}$ .

Tangent line:  $y = 1 - 3x$  or  $2x + 3y = 5$

10.  $x^2 + y^3 = x^3 + y^2$   
 $2xy + 3y^2 y' = 3x^2 + 2y y'$

At  $(1, 2)$ :  $16 + 12y' = 12 + 4y'$ , so the slope is

$y' = \frac{12 - 16}{4 - 12} = \frac{28}{8} = \frac{7}{2}$

Thus, the equation of the tangent line is  $y - 2 = \frac{7}{4}(x - 1)$ , or  $7x - 4y = 1$ .

11.  $\frac{x}{y} = x^3 + y^2$

$x^4 - y^4 = 2x^3 y + 2xy^3$   
 $4x^3 - 4y^3 y' = 6x^2 y + 2x^3 y' + 2y^3 + 6xy^2 y'$

at  $(1, 1)$ :  $4 - 4y' = 6 + 2y' + 2 + 6y'$   
 $2y' = -4$ ,  $y' = -2$

Tangent line:  $y - 1 = -2(x - 1)$  or  $y = 3 - 2x$ .

12.  $x^2 y^2 C_1 D_1$

$$1 C_2 y^0 D_2 \frac{x}{x^2} \frac{1/2 y^0}{1/2} - \frac{y^2 \cdot 1/2}{x^2}$$

At  $x=2$ ;  $y=1$  we have  $1 C_2 y^0 D_2 2y^0 = 1$  so  $y^0 D_2 = \frac{1}{2}$ .  
Thus, the equation of the tangent is

$$y D_1 = \frac{1}{2} x - \frac{1}{2}, \text{ or } x C_2 y D_0:$$

13.  $2 C_1 y^0 D_1 \frac{p}{2 \cos xy} / y C_2 xy^0 D_0$   
At  $x=4$ ;  $y=1$ :  $2 C_1 y^0 D_1 = 4/y^0 / D_0 = 0$ , so

$y^0 D_1 = 4 = 4 /$ . The tangent has equation

$$y D_1 = \frac{4}{x} - 4$$

14.  $\tan xy^2 / D_2 = 1/xy$

$$\sec^2 xy^2 // y^2 C_2 xy^0 D_2 = 1/y C_2 xy^0 / y^0 D_1 = 1/2y^0, \text{ so}$$

$y^0 D_1 = 2 = 4 /$ . The tangent has equation

$$y D_1 = \frac{2}{x} - 4$$

15.  $x \sin xy^2 / D_2 x^2 = 1 / \sin xy^2 / C_2 x \cos xy^2 // y C_2 xy^0 D_2 = 2yy^0 / D_2 x$

At  $x=1$ ;  $y=1$ :  $0 C_2 1/1/1 = y^0 / D_2 = 2$ , so  $y^0 D_2 = 1$ . The tangent has equation  $y D_1 = x - 1$ , or  $y D_2 = x$ .

$$y = x^2 - 17$$

16.  $\cos x^2 y^2 D_2 y^2 = 2 : y^2 // y C_2 xy^0 D_2 = 2xy^0 / x y^2$

$$\frac{y}{x^2} \frac{xy^2}{y^2} = \frac{2xy^0}{x y^2}$$

At  $x=3$ ;  $y=1$ :  $\frac{3}{9} = \frac{2 \cdot 1}{3 \cdot 1} = \frac{2}{3} = \frac{1}{3}$  so  $y^0 D_2 = 108$

so  $y^0 D_2 = 108$ . The tangent has equation

19.  $x^3 y^2 C_1 y^3 D_1 x^2 = \frac{1}{3x^2}$

$$\frac{3x^2}{6x} \frac{2yy^0}{2y^0} C_2 3y^2 y^0 D_1 = \frac{1}{3x^2} \frac{D_2 3y^2}{D_0} \frac{2y}{2y^2}$$

$$\frac{.2}{.2} \frac{6y/y^0}{6x} \frac{6x}{6x} = \frac{.1}{.2} \frac{3x}{3y^2} \frac{1}{2y^2} \frac{1}{6x}$$

$$y^0 D_2 = \frac{3y^2}{6y \cdot 1} \frac{2y}{3x^2} = \frac{3y^2}{6x} \frac{2y}{3x^2}$$

20.  $x^3 3xy C_1 y^3 D_1 = \frac{.3y^2}{3y^2} \frac{2y^3}{2y}$

$$3x^2 3y 3xy^0 C_2 3y^2 y^0 D_0 = 0$$

Thus  $6x 3y^0 3y^0 3xy^0 C_2 6y/y^0 D_2 C_2 3y^2 y^0 D_0 = 0$

$$y = x^2$$

$$y^0 D_2 = \frac{y^2}{x^2} = \frac{0}{0} = 0$$

$$y^0 D_2 = \frac{2x}{y^2} \frac{2y}{x} = \frac{2x \cdot 2y}{y^2 \cdot x}$$

$$D_2 = \frac{2x}{y^2} \frac{2y}{x} = \frac{y^2}{y^2} \frac{x}{x^2} = \frac{y}{y^2} \frac{x^2}{x^2} = \frac{1}{y}$$

$$D_2 = \frac{1}{y} = \frac{1}{2} \frac{1}{x} = \frac{1}{2x}$$

$$\frac{1}{2} = \frac{2xy}{x^2} = \frac{4xy}{x^2}$$

21.  $x^2 C_1 y^2 D_1 a^2 = \frac{0}{0} \frac{0}{0} = \frac{x}{x}$

$2x C_2 2yy D_0$  so  $x C_2 yy D_0$  and  $y D_2 = y$

$$1 C_2 y^0 y^0 C_2 yy^0 D_0 = \frac{1}{x^2}$$

$$1 \cdot y^0 / 2 = 1 C_2 y^2$$

$$y D_2 = \frac{y}{y^2} \frac{y}{x^2} = \frac{y^2}{a^2} = y$$

$$D_2 = y^3 = D_2 y^3$$

22.  $Ax^2 C_1 By^2 D_1 C$

$$2Ax C_2 2By^0 D_0 = y^0 D_2 = \frac{Ax}{By}$$

$$108 p \bar{3}$$

$$y D 1 C 162 3 p 3 . x 3/:$$

17.  $xy D x C y$

$$y C xy^0 D 1 C y^0 y^0 D \frac{y}{1} \frac{1}{x}$$

$$y^0 C y^0 C xy^{00} D y^{00} \frac{2y^0}{2y} \frac{1}{x}$$

Therefore,  $y^{00} D \frac{1}{x} D \frac{1}{x^2}$

18.  $x^2 C 4y^2 D 4; 2x C 8yy^0 D 0; 2 C 8y^{0/2} C 8yy^{00} D 0.$

Thus,  $y^0 D 4y$  and

$$y^{00} D \frac{2 \cdot 8y^{0/2}}{8y} D \frac{16y^{1/2}}{8y} D \frac{2y^{1/2}}{y} D \frac{2y^{1/2} \cdot x^2}{16y^3} D \frac{1}{4y^3}:$$

2A C 2B.y<sup>0/2</sup> C 2Byy<sup>00</sup> D 0.  
Thus,

$$y^{00} D \frac{A \cdot B.y^{0/2}}{By} D \frac{A \cdot B \cdot \frac{1}{Ax}}{\frac{By}{2}} D \frac{A \cdot B \cdot C \cdot Ax}{B^2y^3} D \frac{AC}{B^2y^3}:$$

23. Maple gives 0 for the value.

$$\underline{206}$$

24. Maple gives the slope as  $\frac{5}{55}$ .

25. Maple gives the value  $\frac{26}{855,000}$ .

26. Maple gives the value  $\frac{371}{293}$ .



27. Ellipse:  $x^2 + 2y^2 = 2$   
 $\frac{x^2}{2} + y^2 = 1$   
 Slope of ellipse:  $y' = -\frac{x}{2y}$   
 Hyperbola:  $2x^2 - 2y^2 = 1$

$4x - 4yy' = 0$   
 Slope of hyperbola:  $y' = \frac{x}{y^2}$

At intersection points  $x^2 + 2y^2 = 2$   
 $2x^2 - 2y^2 = 1$

$3x^2 = 3$  so  $x^2 = 1$ ,  $y^2 = \frac{1}{2}$   
 Thus  $y = \pm \frac{1}{\sqrt{2}}$

$E_H = \frac{1}{2y} = \pm \sqrt{2}$

Therefore the curves intersect at right angles.

28. The slope of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is found from

$\frac{2x}{a^2} + \frac{2y}{b^2}y' = 0$ ; i.e.  $y' = -\frac{b^2x}{a^2y}$

Similarly, the slope of the hyperbola  $\frac{x^2}{A^2} - \frac{y^2}{B^2} = 1$  at  $(x, y)$  satisfies

$\frac{2x}{A^2} - \frac{2y}{B^2}y' = 0$ ; or  $y' = \frac{B^2x}{A^2y}$

If the point  $(x, y)$  is an intersection of the two curves,

then  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$   
 $\frac{x^2}{A^2} - \frac{y^2}{B^2} = 1$   
 $\frac{1}{x^2} \left( \frac{1}{A^2} - \frac{1}{a^2} \right) = \frac{1}{y^2} \left( \frac{1}{B^2} - \frac{1}{b^2} \right)$

Thus,  $\frac{x^2}{A^2} = \frac{b^2}{B^2} \frac{a^2}{a^2}$

Since  $\frac{y^2}{b^2} = \frac{1}{b^2} \left( 1 - \frac{x^2}{a^2} \right)$ , therefore  $\frac{B^2}{b^2} = \frac{a^2}{A^2}$

and  $y^2 = \frac{B^2 b^2}{a^2}$ . Thus, the product of the slope of the two curves at  $(x, y)$  is

30.  $\frac{x}{x} + \frac{y}{y} = 1$ ,  $xy = y^2$   
 $\frac{1}{x} - \frac{y}{y^2} = 0$

Differentiate with respect to  $x$ :

$2x - 4yy' = 0$  ;  $y' = \frac{2x}{4y} = \frac{x}{2y}$

However, since  $x^2 + 2y^2 = 2$  can be written

$x^2 + 4y^2 = 2$ ; or  $x^2 = 2 - 4y^2$

the only solution is  $x = 0, y = 0$ , and these values do not satisfy the original equation. There are no points on the given curve.

**Section 2.10 Antiderivatives and Initial-Value Problems (page 155)**

1.  $\int 5 dx = 5x + C$

2.  $\int x^2 dx = \frac{1}{3} x^3 + C$

3.  $\int x dx = \frac{1}{2} x^2 + C$

4.  $\int x^{12} dx = \frac{1}{13} x^{13} + C$

5.  $\int x^3 dx = \frac{1}{4} x^4 + C$

6.  $\int \cos x dx = \sin x + C$

7.  $\int \tan x \cos x dx = \int \sin x dx = -\cos x + C$

$$\frac{b^2 x}{a^2 y} = \frac{B^2 x}{A^2 y} \quad \frac{b^2 B^2}{a^2 A^2} = \frac{A^2 a^2}{B^2 b^2} \quad \mathbf{D} \quad 1:$$

Therefore, the curves intersect at right angles.

29. If  $z = \tan x = 2/t$ , then

$$\frac{1}{\sec^2 x} \cdot \frac{dx}{dz} = \frac{1}{2} \frac{dx}{dz} \quad \frac{1}{2} \frac{dx}{dz} = \frac{1}{2} \frac{dz}{dz} \quad \mathbf{D} \quad \frac{1}{2} \frac{dz}{dz}$$

Thus  $dx = dz = 2/z^2$ . Also

$$\cos x = \frac{1}{\sec x} = \frac{1}{2/z} = \frac{z}{2}$$

$$\frac{1}{2} \frac{dz}{dz} = \frac{1}{2} \frac{z^2}{z^2}$$

$$\sin x = \frac{z}{2} \quad \frac{1}{2} \frac{dz}{dz} = \frac{1}{2} \frac{z^2}{z^2} \quad \mathbf{D} \quad \frac{1}{2} \frac{dz}{dz} = \frac{1}{2} \frac{z^2}{z^2}$$

$$8. \int \frac{1}{\cos^2 x} dx = \int \sec^2 x dx = \tan x + C$$

$$9. \int \frac{1}{x^2} dx = \int x^{-2} dx = \frac{x^{-1}}{-1} + C = -\frac{1}{x} + C$$

$$10. \int \frac{1}{x^2} dx = \int x^{-2} dx = \frac{x^{-1}}{-1} + C = -\frac{1}{x} + C$$

$$11. \int \frac{1}{x^2} dx = \int x^{-2} dx = \frac{x^{-1}}{-1} + C = -\frac{1}{x} + C$$

$$12. \int \frac{1}{x^4} dx = \int x^{-4} dx = \frac{x^{-3}}{-3} + C = -\frac{1}{3x^3} + C$$

$$\int \frac{1}{x^2} dx = \int x^{-2} dx = \frac{x^{-1}}{-1} + C = -\frac{1}{x} + C$$

$$\int \frac{1}{x^2} dx = \int x^{-2} dx = \frac{x^{-1}}{-1} + C = -\frac{1}{x} + C$$

$$13. \int \frac{1}{x^3} dx = \int x^{-3} dx = \frac{x^{-2}}{-2} + C = -\frac{1}{2x^2} + C$$

14.  $\int 105 t^2 c t^4 c t^6 / dt$   
 $D 105 t^3 c t^5 c t^7 / C C$   
 $D 105 t^3 c 35 t^3 c 21 t^5 c 15 t^7 C C$

15.  $\int \cos 2x / dx D \int 2 \sin 2x / C C$

16.  $\int \sin 2x dx D \int 2 \cos 2x C C$   
 $\frac{dx}{1}$

17.  $\int .1 C x^2 D 1 C x C C$

18.  $\int \sec 1 x / \tan 1 x / dx D \sec 1 x / C C$

19.  $\int p \frac{1}{2x} C 3 dx D \int 3 .2x C 3 / 3=2 C C$

20.  $\frac{d p}{dx} = \frac{1}{x^2} C 1 D$ , therefore  
 $\frac{1}{2 x C 1}$

$\int \frac{4}{x^2} dx D 8 \frac{1}{x C 1} C C$   
 $x C 1$

21.  $\int 2x \sin x^2 / dx D \int \cos x^2 / C C$   
 $p = \frac{1}{2} x^2$

22. Since  $\frac{dx}{2x} = \frac{1}{2} \frac{dx}{x} D p = \frac{1}{2} x^2 C 1$ , therefore  
 $\int \frac{1}{p^2} dx D 2 \frac{1}{x^2} C 1 C C$   
 $C p$

23.  $\int \tan^2 x dx D \int \sec^2 x - 1 / dx D \tan x - x C C$

24.  $\int \sin x \cos x dx D \int \frac{1}{2} \sin 2x / dx D \frac{1}{4} \cos 2x / C C$

29.  $y^4 / D 3 p x } y \int 1 D 16 C \text{ so } C 15$   
 $2x^{3=2} C C$

$D 3=2 D C D$   
 Thus  $y D$

30. Given that  $2x = 15$  for  $x > 0$ .

$y^0 D x^{1=3}$   
 $y.0 / D 5$ ;

then  $y D x^{1=3} dx D \frac{3}{4} x^{4=3} C C$  and  $5 D y.0 / D C$ . Hence,  
 $y.x / D 4 x^4 C 5$  which is valid on the whole real line.

31. Since  $y^0 D A x^2 C B x C C$  we have  
 $y D \frac{A}{x} - \frac{B}{x^2}$

$3 x^3 C 2 x^2 C C x C D$ . Since  $y.1 / D T$ , therefore  
 $\frac{A}{x} - \frac{B}{x^2}$

$1 D y.1 / D 3 C 2 C C C D$ . Thus  $D D 1 - 3 - 2 C$ , and

$\frac{A}{x} - \frac{B}{x^2}$   
 $y D 3 .x - 1 / C 2 .x - 1 / C C .x - 1 / C 1$  for all  $x$

32. Given that

$y^0 D x^{9=7}$

$y.1 / D 4$ ;

then  $y D \int x^{9=7} dx D \frac{1}{2} x^{2=7} C C$ .

Also,  $4 D y.1 / D \frac{1}{2} C C$ , so  $C D \frac{1}{2}$ . Hence,

$y D \frac{1}{2} x^{2=7} + \frac{1}{2}$ , which is valid in the interval  $(0; 1/2)$ .

$y. = 6 / D 2$

33. For  $y^0 D \cos x$ , we have

$y D \int \cos x dx D \sin x C C$

$2 D \sin \frac{1}{6} C C D \frac{1}{2} C C \div C D \frac{1}{2}$

$y D \sin x C \frac{1}{2}$  (for all  $x$ );

25.  $\int \cos^2 x \, dx = \int \frac{1 + \cos 2x}{2} \, dx = \frac{x}{2} + \frac{\sin 2x}{4} + C$

26.  $\int \sin^2 x \, dx = \int \frac{1 - \cos 2x}{2} \, dx = \frac{x}{2} - \frac{\sin 2x}{4} + C$

27.  $(y'' + 2y) = x^2$   
 Thus  $y'' = x^2 - 2y$  for all  $x$ .  
 28. Given that  $y'' = x^2 - 2y$

$\int y'' \, dx = \int (x^2 - 2y) \, dx$

then  $y' = \frac{1}{3}x^3 - 2\int y \, dx + C$

and  $0 = y' - \frac{1}{3}x^3 + 2\int y \, dx - C$  so  $C = \frac{1}{3}x^3 + 2\int y \, dx$

Hence,  $y = \frac{1}{x} + \frac{1}{2x^2} + \frac{3}{2}$  which is valid on the interval  $(-\infty, 0) \cup (0, \infty)$ .

$y' = \sin 2x$

34. For  $y = \frac{1}{2} \cos 2x$ , we have  $y' = -\sin 2x$

$\frac{1}{2}$

$\int \sin 2x \, dx = -\frac{1}{2} \cos 2x + C$

$\frac{1}{2} \cos 2x = -\int \sin 2x \, dx + C = \frac{1}{2} \cos 2x + C$

$\frac{1}{2}$

$y = \frac{1}{2} \cos 2x$  (for all  $x$ ):

35. For  $y = \sec^2 x$ , we have

$y' = 2 \sec^2 x \tan x$

$\int \sec^2 x \, dx = \tan x + C$

$\frac{1}{2} \tan^2 x = \int \sec^2 x \, dx - C = \tan x + C$  (for  $-\frac{\pi}{2} < x < \frac{\pi}{2}$ )

36. For  $y = \sec^2 x$ , we

have  $y' = 2 \sec^2 x \tan x$

$$y' = 2 \sec^2 x \tan x = 2 \tan x \sec^2 x$$

$$= 2 \tan x (1 + \tan^2 x) = 2 \tan x + 2 \tan^3 x$$

37. Since  $y'' = 2$ , therefore  $y' = 2x + C_1$ .  
 Since  $y(0) = 5$ , therefore  $5 = 0 + C_1$ , and  $y' = 2x + 5$ .  
 Thus  $y = x^2 + 5x + C_2$ .  
 Since  $y(0) = 3$ , therefore  $3 = 0 + 0 + C_2$ , and  $C_2 = 3$ .  
 Finally,  $y = x^2 + 5x + 3$ , for all  $x$ .

38. Given that

$$y'' = x^4$$

$$y' = \frac{1}{5} x^5 + C_1$$

$$y = \frac{1}{30} x^6 + \frac{1}{2} C_1 x + C_2$$

then  $y' = x^4 dx = \frac{1}{5} x^5 + C_1$

Since  $y(0) = 1$ , therefore  $1 = \frac{1}{30} C_2$ , therefore  $C_2 = 30$

and  $y' = x^4$

$$y' = x^4 = \frac{1}{5} x^5 + C_1$$

$$y = \frac{1}{30} x^6 + \frac{1}{2} C_1 x + C_2$$

and  $y(0) = 1$ , therefore  $1 = \frac{1}{30} C_2$ , so that  $C_2 = 30$ . Hence,

$y = \frac{1}{30} x^6 + \frac{1}{2} C_1 x + 30$ , which is valid in the interval  $(-\infty, \infty)$ .

39. Since  $y'' = 3x^2$ , therefore  $y' = x^3 + C_1$ .

Since  $y(0) = 0$ , therefore  $0 = 0 + C_1$ , and

$$y' = x^3$$

$$y = \frac{1}{4} x^4 + C_2$$

Thus  $y = \frac{1}{4} x^4 + C_2$ .

Since  $y(0) = 8$ , we have  $8 = 0 + C_2$ .

$$y'' = \cos x$$

41. For  $y(0) = 0$ , we have

$$y' = \sin x$$

$$y = -\cos x + C_1$$

$$0 = -\cos 0 + C_1 = -1 + C_1$$

$$C_1 = 1$$

$$y = 1 - \cos x$$

$$y'' = x \cos x$$

42. For  $y'' = 2$ , we have

$$y' = 2x + C_1$$

$$y = x^2 + C_1 x + C_2$$

$$0 = 0 + 0 + C_2 = C_2$$

$$C_2 = 0$$

$$y' = 2x + C_1$$

$$2 = 0 + 0 + C_1 = C_1$$

$$C_1 = 2$$

$$y = x^2 + 2x$$

43. Let  $y = Ax + \frac{B}{x}$ . Then  $y' = A - \frac{B}{x^2}$ , and  $y'' = \frac{2B}{x^3}$ .

Thus, for all  $x \neq 0$ ,

$$y'' = \frac{2B}{x^3} = A - \frac{B}{x^2}$$

$$\frac{2B}{x^3} = A - \frac{B}{x^2}$$

$$2B = Ax^3 - Bx$$

$$2B = Ax^3 - Bx$$

We will also have  $y(0) = 2$  and  $y'(0) = 4$  provided

Hence  $y = \frac{1}{20x^5} + \frac{1}{2x^2} + C$  for all  $x$ .

40. Given that  $y'' = 8x^{1/2} - 5x^{-2} - 3x^{-1/2}$

$y(1) = 0$ ;

we have  $y' = \int (8x^{1/2} - 5x^{-2} - 3x^{-1/2}) dx = \frac{16}{3}x^{3/2} + \frac{5}{x} - 6x^{1/2} + C$ .

Also,  $y'(1) = \frac{16}{3} + 5 - 6 + C = 0$  so that  $C = -\frac{5}{3}$ . Thus,

$y' = \frac{16}{3}x^{3/2} + \frac{5}{x} - 6x^{1/2} - \frac{5}{3}$ , and

$y = \int (\frac{16}{3}x^{3/2} + \frac{5}{x} - 6x^{1/2} - \frac{5}{3}) dx = \frac{32}{15}x^{5/2} + 5 \ln|x| - 4x^{3/2} - \frac{5}{3}x + C$ .

Finally,  $y(1) = \frac{32}{15} + 5 - 4 - \frac{5}{3} + C = 0$  so that  $C = -\frac{11}{4}$ .

Hence,  $y(x) = \frac{32}{15}x^{5/2} + 5 \ln|x| - 4x^{3/2} - \frac{5}{3}x - \frac{11}{4}$ .

$AC = 2$ ; and  $A = 2, B = 4$ :

These equations have solution  $A = 2, B = 4$ , so the initial value problem has solution  $y = 2x^2 - 4x$ .

44. Let  $r_1$  and  $r_2$  be distinct rational roots of the equation

$ax^2 + bx + c = 0, x > 0$

Let  $y = Ax^{r_1} + Bx^{r_2}$

Then  $y' = Ar_1Ax^{r_1-1} + Br_2Bx^{r_2-1}$ ,  
and  $y'' = Ar_1(r_1-1)Ax^{r_1-2} + Br_2(r_2-1)Bx^{r_2-2}$ . Thus

$ax^2 + bx + c = 0$

$Aax^2 + Ar_1(r_1-1)Ax^{r_1-2} + Bbx + Br_2(r_2-1)Bx^{r_2-2} = 0$

$Cbx + Ar_1Ax^{r_1-1} + Br_2Bx^{r_2-1} = CcAx^{r_1} + CBx^{r_2} / r_1$

$A = ar_1(r_1-1) / C, B = br_2 / C$

$B = ar_2 / C$

$D = 0, C = 0, 0, x > 0$

45.  $84x^2 y^{00} C 4xy^0 y D 0 . /) a D 4; b D 4; c D 1$   
 $y.4/ D 2$

Auxiliary Equation:  $4r.r 1/ C 4r 1 D 0$   
 $4r^2 1 D 0$   
 $1$

By #31,  $y D Ax^{1=2} C Bx^{1=2}$  solves . / for  $x > 0$ .

Now  $y^0 D 2x^{1=2} C 2x^{3=2}$

Substitute the initial conditions:

$$\begin{aligned} & \frac{B}{2} + \frac{B}{16} = \frac{4}{2} \\ & \frac{2D}{4} + \frac{B}{16} = \frac{4}{2} \end{aligned}$$

Hence  $9 D \frac{2}{7}$ , so  $B D 18, AD \frac{7}{2}$ .

Thus  $y D 2x^{1=2} C 18x^{1=2}$  (for  $x > 0$ ).

46. Consider

$$\begin{aligned} & < x^2 y^{00} 6y D 0 \\ & y.1/ D 1 \\ & : y^0 .1/ D 1: \end{aligned}$$

Let  $y D x^r$ ;  $y^0 D r x^{r-1}$ ;  $y^{00} D r.r 1/x^{r-2}$ . Substituting these expressions into the differential equation we obtain

$$\begin{aligned} & x^2 C r.r 1/x^{r-2} 6x^r D 0 \\ & C r.r 1/ 6x^r D 0: \end{aligned}$$

Since this equation must hold for all  $x > 0$ , we must have

$$\begin{aligned} & r.r 1/ 6 D 0 \\ & r^2 r 6 D 0 \\ & .r 3/r C 2/ D 0: \end{aligned}$$

There are two roots:  $r_1 D 2$ , and  $r_2 D 3$ . Thus the differential equation has solutions of the form  $y D Ax^2 C Bx^3$ . Then  $y^0 D 2Ax C 3Bx^2$ . Since

$1 D y; 1/ D A B$  and  $1 D y^0 .1/ 2A C 3B$ , therefore

- d) never accelerating to the left
- e) particle is speeding up for  $t > 2$
- f) slowing down for  $t < 2$
- g) the acceleration is 2 at all times
- h) average velocity over  $0t4$  is

$$\frac{x.4/ .x.0/}{4} D \frac{16 16C3 3}{4} D 4$$

2.  $x D 4 C 5t^2, v D 5 2t, a D 2$ .

- a) The point is moving to the right if  $v > 0$ , i.e., when  $t < \frac{5}{2}$ .
- b) The point is moving to the left if  $v < 0$ , i.e., when  $t > \frac{5}{2}$ .
- c) The point is accelerating to the right if  $a > 0$ , but  $a D 2$  at all  $t$ ; hence, the point never accelerates to the right.
- d) The point is accelerating to the left if  $a < 0$ , i.e., for all  $t$ .
- e) The particle is speeding up if  $v$  and  $a$  have the same sign, i.e., for  $t > \frac{5}{2}$ .
- f) The particle is slowing down if  $v$  and  $a$  have opposite sign, i.e., for  $t < \frac{5}{2}$ .
- g) Since  $a D 2$  at all  $t$ ,  $a D 2$  at  $t D 2$  when  $v D 0$ .
- h) The average velocity over  $0t4$  is

$$\frac{x.4/ .x.0/}{4} D \frac{8 4}{4} D 2$$

3.  $x D t^3, v D dt D 3t^2, a D dt D 6t$

- a) particle moving: to the right for  $t < 2=$  3 or  $t > 2=$  3 .

A  $\frac{dx}{dt} = 5$  and B  $\frac{dx}{dt} = 5$ . Hence,  $y = 5x + C$ .

**Section 2.11 Velocity and Acceleration (page 162)**

$$1. x = \frac{1}{2}t^2 - 4t + 3, v = \frac{dx}{dt} = t - 4, a = \frac{dv}{dt} = 1$$

- a) particle is moving: to the right for  $t > 2$
- b) to the left for  $t < 2$
- c) particle is always accelerating to the right

- b) to the left for  $2 = \frac{p}{3} < t < 2 = \frac{p}{3}$
- c) particle is accelerating: to the right for  $t > 0$
- d) to the left for  $t < 0$
- e) particle is speeding up for  $t > 2 = \frac{p}{3}$  or for  $2 = \frac{p}{3} < t < 0$
- f) particle is slowing down for  $t < 2 = \frac{p}{3}$  or for  $0 < t < 2 = \frac{p}{3}$
- g) velocity is zero at  $t = \frac{p}{3}$ . Acceleration at these times is  $\frac{1}{3}$ .
- h) average velocity on  $[0, 4]$  is  $\frac{4^3 - 4}{4 \cdot 0} = \frac{1}{12}$



4.x  $D \frac{t}{t^2 C 1} ; v \frac{.t^2 C 1/.1/ .t/.2t/}{D -1 .t^2} ;$   
 $t^2 C 1 \quad D \quad .t^2 C 1/2 \quad .t^2 C 1/2$   
 a  $\frac{t^2 C 1/2 .2t/ .1 t^2/.2/.t^2 C 1/.2t/}{D \quad .t^2 C 1/4 \quad D \quad .t^2 C 1/3} \quad \frac{2t t^2 - 3/}{D \quad .t^2 C 1/3}$

a) The point is moving to the right if  $v > 0$ , i.e., when  $1 - t^2 > 0$ , or  $1 < t < 1$ .

b) The point is moving to the left if  $v < 0$ , i.e., when  $t < 1$  or  $t > 1$ .

c) The point is accelerating to the right if  $a > 0$ , i.e., when  $2t - t^3 > 0$ , that is, when

$$t > \frac{1}{3} \text{ or } \frac{1}{3} < t < 1.$$

d) The point is accelerating to the left if  $a < 0$ , i.e., for

$$t < \frac{1}{3} \text{ or } 0 < t < \frac{1}{3}.$$

e) The particle is speeding up if  $v$  and  $a$  have the same sign, i.e., for  $t < \frac{1}{3}$ , or  $1 < t < 1$  or  $\frac{1}{3} < t < 1$ .

f) The particle is slowing down if  $v$  and  $a$  have opposite sign, i.e., for  $\frac{1}{3} < t < 1$ , or  $0 < t < \frac{1}{3}$  or  $t > 1$ .

$$t > 1 \quad \frac{2t - t^3}{1 - t^2} < 0$$

g)  $v = 0$  at  $t = 1$ . At  $t = 1$ ,  $a = -2$ .

$$\text{At } t = 1, a = -2.$$

h) The average velocity over  $[0, 4]$  is

$$\frac{x(4) - x(0)}{4 - 0} = \frac{4 - 0}{4} = 1$$

$$\frac{4}{4} = 1 \quad \frac{4}{4} = 1$$

5.  $y = 9.8t - 4.9t^2$  metres (t in seconds)

$$\text{velocity } v = \frac{dy}{dt} = 9.8 - 9.8t$$

$$\text{acceleration } a = \frac{dv}{dt} = -9.8$$

The acceleration is  $9.8 \text{ m/s}^2$  downward at all times. Ball is at maximum height when  $v = 0$ , i.e., at  $t = 1$ . Thus maximum height is  $y = 9.8(1) - 4.9(1)^2 = 4.9$  metres.

$$t = 1$$

7.  $D = t^2$ , D in metres, t in seconds

$$D = t^2$$

$$\text{velocity } v = \frac{dD}{dt} = 2t$$

Aircraft becomes airborne if  $v = 200 \text{ km/h} = \frac{200,000}{3600} \text{ m/s} = 55.56 \text{ m/s}$

$$2t = 55.56 \Rightarrow t = 27.78$$

Time for aircraft to become airborne is  $t = 27.78$  s, that is, 9 about 27.8 s.

Distance travelled during takeoff run is  $t^2 = 771.6$  metres.

8. Let  $y(t)$  be the height of the projectile t seconds after it is fired upward from ground level with initial speed  $v_0$ . Then

$$y(0) = 0; \quad y'(0) = v_0; \quad y''(0) = -g$$

Two antiderivations give

$$y = \frac{1}{2}gt^2 + v_0t + y(0) = \frac{1}{2}gt^2 + v_0t$$

Since the projectile returns to the ground at  $t = 10$  s, we have  $y(10) = 0$ , so  $v_0 = 49$  m/s. On Mars, the acceleration of gravity is  $3.72 \text{ m/s}^2$  rather than  $9.8 \text{ m/s}^2$ , so the height of the projectile would be

$$y = \frac{1}{2}(3.72)t^2 + v_0t = 1.86t^2 + 49t$$

The time taken to fall back to ground level on Mars would be  $t = 49/1.86 = 26.3$  s.

9. The height of the ball after t seconds is

$y(t) = \frac{1}{2}gt^2 + v_0t$  if its initial speed was  $v_0$  m/s. Maximum height h occurs when  $dy = dt = 0$ , that is, at  $t = \frac{v_0}{g}$ .

$v_0 = g \frac{v_0}{g}$ . Hence

$$h = \frac{1}{2}g \left(\frac{v_0}{g}\right)^2 + v_0 \left(\frac{v_0}{g}\right) = \frac{v_0^2}{2g} + \frac{v_0^2}{g} = \frac{3v_0^2}{2g}$$

$$h = \frac{3}{2}g \left(\frac{v_0}{g}\right)^2 = \frac{3}{2} \frac{v_0^2}{g}$$

An initial speed of  $2v_0$  means the maximum height will be  $4 \times \frac{3}{2} \frac{v_0^2}{g} = 6 \frac{v_0^2}{g}$ . To get a maximum height of  $2h$  an

initial speed of  $\sqrt{2}v_0$  is required.

10. To get to  $3h$  metres above Mars, the ball would have to be thrown upward with speed

$$D = \frac{3}{2}g \left(\frac{v_0}{g}\right)^2 = \frac{3}{2} \frac{v_0^2}{g}$$

Ball strikes the ground when  $y = 0$ ,  $t > 0$ , i.e.,  
 $0 = 16t - 4.9t^2$  / so  $t = 2$ .  
 Velocity at  $t = 2$  is  $v = 9.8 - 9.8(2) = -9.8$  m/s.  
 Ball strikes the ground travelling at 9.8 m/s (downward).

6. Given that  $y = 100 - 2t - 4.9t^2$ , the time  $t$  at which the ball reaches the ground is the positive root of the equation

$$y = 0, \text{ i.e., } 100 - 2t - 4.9t^2 = 0, \text{ namely,}$$

$$t = \frac{2 \pm \sqrt{4 + 4 \cdot 4.9 \cdot 100}}{2 \cdot 4.9} \approx 4.318 \text{ s}$$

The average velocity of the ball is  $\frac{100}{4.318} \approx 23.16$  m/s.

Since  $v = 9.8 - 9.8t$ , then  $t = 2.159$  s.

Since  $v = 32 - 32t$  and  $y = 16t^2$ , we have  $v = 16t$  m/s.

11. If the cliff is  $h$  ft high, then the height of the rock  $t$  seconds after it falls is  $y = h - 16t^2$  ft. The rock hits the ground ( $y = 0$ ) at time  $t = \sqrt{\frac{h}{16}} = \frac{\sqrt{h}}{4}$  s. Its speed at that time is  $v = 32t = \frac{32\sqrt{h}}{4} = 8\sqrt{h}$  ft/s. Thus  $8\sqrt{h} = 20$

and the cliff is  $h = 400$  ft high.

12. If the cliff is  $h$  ft high, then the height of the rock  $t$  seconds after it is thrown down is  $y = h - 32t - 16t^2$  ft. The rock hits the ground ( $y = 0$ ) at time

$$t = \frac{32 \pm \sqrt{32^2 - 4 \cdot 16 \cdot h}}{2 \cdot 16} = \frac{32 \pm \sqrt{1024 - 64h}}{32}$$

Its speed at that time is

$$v = 32t - 16t^2 \quad \text{ft/s}$$

Solving this equation for  $h$  gives the height of the cliff as 384 ft.

13. Let  $x(t)$  be the distance travelled by the train in the  $t$  seconds after the brakes are applied. Since  $x'(t) = 60 - 6t$

m/s and since the initial speed is  $v_0 = 60$  km/h  $\approx 100$  m/s, we have

$$x(t) = 60t - 3t^2$$

The speed

of the train at time  $t$  is  $v(t) = 60 - 6t$  m/s,

so it takes the train 100 s to come to a stop. In that time it travels  $x(100) = 60(100) - 3(100)^2 = 6000 - 30000 = -24000$  metres.

14.  $x(t) = At^2 + Bt + C$ ;  $v(t) = 2At + B$ .

The average velocity over  $[t_1, t_2]$  is

$$\frac{x(t_2) - x(t_1)}{t_2 - t_1}$$

$$= \frac{A(t_2^2 - t_1^2) + B(t_2 - t_1) + C - C}{t_2 - t_1}$$

$$= \frac{A(t_2 + t_1)(t_2 - t_1) + B(t_2 - t_1)}{t_2 - t_1}$$

$$= \frac{A(t_2 + t_1) + B}{1}$$

$$= \frac{A(t_2 + t_1) + B}{1}$$

$$= \frac{A(t_2 + t_1) + B}{1}$$

The instantaneous velocity at the midpoint of  $[t_1, t_2]$  is

$$v\left(\frac{t_1 + t_2}{2}\right) = 2A\left(\frac{t_1 + t_2}{2}\right) + B = A(t_1 + t_2) + B$$

Hence, the average velocity over the interval is equal to the instantaneous velocity at the midpoint.

15.

$$v(t) = \begin{cases} 68 - 20t & 0 \leq t < 2 \\ 20 & 2 \leq t < 8 \\ 20 + 2t & 8 \leq t < 10 \end{cases}$$

Note:  $v(2) = 68 - 20(2) = 28$

$$v(t) = \begin{cases} 68 - 20t & \text{if } 0 < t < 2 \\ 28 & \text{if } 2 < t < 8 \\ 20 + 2t & \text{if } 8 < t < 10 \end{cases}$$

Since  $\lim_{t \rightarrow 2^-} v(t) = 28 = v(2)$ , therefore,  $v$  is continuous at  $t = 2$ .

Since  $\lim_{t \rightarrow 8^-} v(t) = 28 = v(8)$ , therefore  $v$  is continuous at  $t = 8$ . Hence the velocity is continuous for




Fig. 2.11-16

The rocket's acceleration while its fuel lasted is the slope of the first part of the graph, namely  $96/7 = 24$  ft/s.

17. The rocket was rising until the velocity became zero, that is, for the first 7 seconds.
18. As suggested in Example 1 on page 154 of the text, the distance travelled by the rocket while it was falling from its maximum height to the ground is the area between the velocity graph and the part of the  $t$ -axis where  $v < 0$ . The area of this triangle is  $\frac{1}{2}(14)(96) = 672$  ft. This is the maximum height the rocket achieved.
19. The distance travelled upward by the rocket while it was rising is the area between the velocity graph and the part of the  $t$ -axis where  $v > 0$ , namely  $\frac{1}{2}(7)(96) = 336$  ft. Thus the height of the tower from which the rocket was fired is  $672 + 336 = 1008$  ft.
20. Let  $s(t)$  be the distance the car travels in the  $t$  seconds after the brakes are applied. Then  $s'(t) = v(t)$  and the velocity at time  $t$  is given by

$$s(t) = \int v(t) dt = \int \begin{cases} 68 - 20t & 0 \leq t < 2 \\ 28 & 2 \leq t < 8 \\ 20 + 2t & 8 \leq t < 10 \end{cases} dt$$

where  $C_1 = 20$  m/s (that is, 72 km/h) as determined in Example 6. Thus

$$0 < t < 10$$

$$\text{acceleration } a = \frac{dv}{dt} = \begin{cases} 2 & \text{if } 0 < t < 2 \\ 0 & \text{if } 2 < t < 8 \\ 2 & \text{if } 8 < t < 10 \end{cases}$$

is discontinuous at  $t = 2$  and  $t = 8$ .  
 Maximum velocity is 4 and is attained on the interval  $2 < t < 8$ .

16. This exercise and the next three refer to the following figure depicting the velocity of a rocket fired from a tower as a function of time since firing.

$$s(t) = \begin{cases} \frac{1}{2}at^2 + v_0t + s_0 & 0 \leq t < 2 \\ v_0t + s_0 & 2 \leq t < 8 \\ \frac{1}{2}at^2 + v_0t + s_0 & 8 \leq t \leq 10 \end{cases}$$

where  $C_2 = 0$  because  $s(0) = 0$ . The time taken to come to a stop is given by  $s'(t) = 0$ , so it is  $t = 4$  s. The distance travelled is

$$s(4) = \frac{1}{2}(2)(4)^2 + 0(4) + 0 = 16 \text{ m.}$$

Review Exercises 2 (page 163)

1.  $y = 3x^2$   
 $\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{3(x+h)^2 - 3x^2}{h} = \lim_{h \rightarrow 0} \frac{9x^2 + 6xh + 3h^2 - 3x^2}{h} = \lim_{h \rightarrow 0} \frac{6xh + 3h^2}{h} = \lim_{h \rightarrow 0} (6x + 3h) = 6x$

2.  $\lim_{h \rightarrow 0} \frac{1 - (x-h)^2}{h} = \lim_{h \rightarrow 0} \frac{1 - (x^2 - 2xh + h^2)}{h} = \lim_{h \rightarrow 0} \frac{1 - x^2 + 2xh - h^2}{h} = \lim_{h \rightarrow 0} \frac{2xh - h^2}{h} = \lim_{h \rightarrow 0} (2x - h) = 2x$

3.  $f(x) = x^4$   
 $f'(x) = \lim_{h \rightarrow 0} \frac{(x+h)^4 - x^4}{h} = \lim_{h \rightarrow 0} \frac{4x^3h + 6x^2h^2 + 4xh^3 + h^4}{h} = \lim_{h \rightarrow 0} (4x^3 + 6x^2h + 4xh^2 + h^3) = 4x^3$

4.  $g(t) = t^5$   
 $g'(t) = \lim_{h \rightarrow 0} \frac{(t+h)^5 - t^5}{h} = \lim_{h \rightarrow 0} \frac{5t^4h + 10t^3h^2 + 10t^2h^3 + 5th^4 + h^5}{h} = \lim_{h \rightarrow 0} (5t^4 + 10t^3h + 10t^2h^2 + 5th^3 + h^4) = 5t^4$

5. The tangent to  $y = \cos x$  at  $x = 1$  has slope

$\frac{dy}{dx} = -\sin x$   
 at  $x = 1$ , slope =  $-\sin 1$

8.  $\frac{d}{dx} (x^4 + x^3 + x^2 + x) = 4x^3 + 3x^2 + 2x + 1$

9.  $\frac{d}{dx} (x^{2/5} - x^{5/2}) = \frac{2}{5}x^{-3/5} - \frac{5}{2}x^{3/2}$

10.  $\frac{d}{dx} (\cos^2 x) = 2 \cos x (-\sin x) = -2 \cos x \sin x = -\sin 2x$

11.  $\frac{d}{dx} (\tan^{-1} x) = \frac{1}{1+x^2}$

12.  $\frac{d}{dt} (\tan^{-1} t) = \frac{1}{1+t^2}$

13.  $\lim_{h \rightarrow 0} \frac{(x+h)^{20} - x^{20}}{h} = 20x^{19}$

14.  $\lim_{h \rightarrow 0} \frac{(4+h)^3 - 4^3}{h} = 3(4)^2 = 48$

15.  $\lim_{h \rightarrow 0} \frac{\cos(x+h) - \cos x}{h} = -\sin x$

16.  $\frac{d}{dx} (\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$

17.  $\frac{d}{dx} (\cos^{-1} \frac{2x}{3}) = \frac{-1}{\sqrt{1-(2x/3)^2}} \cdot \frac{2}{3} = \frac{-2}{3\sqrt{1-(2x/3)^2}}$

Its equation is

$$y = \frac{1}{2}x^2 - 6x + 1$$

6. At  $x = 4$  the curve  $y = \tan x$  has slope  $\sec^2 x = 4/3$ . The normal to the curve there

has equation  $y = 1 - 2x/3$ .

$$\frac{d}{dx} \frac{1}{\cos x} = \frac{1}{\cos^2 x}$$

7.  $\frac{dx}{x} = \sin x$   $\frac{d}{dx} \sin x = \cos x$

16.  $\lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} = \lim_{h \rightarrow 0} \frac{a^x(a^h - 1)}{h} = a^x \lim_{h \rightarrow 0} \frac{a^h - 1}{h} = a^x \ln a$

$\frac{d}{dx} a^{ax^3} = a^{ax^3} \ln a \cdot 3ax^2 = 3a^2 x^2 \ln a a^{ax^3}$

17.  $\frac{d}{dx} x^2 \ln x = 2x \ln x + x^2 \cdot \frac{1}{x} = 2x \ln x + x$

18.  $\frac{d}{dx} x^p = p x^{p-1}$

$\frac{d}{dx} \frac{f(x)}{g(x)} = \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}$

19.  $\frac{d}{dx} x^{2/p} = \frac{2}{p} x^{2/p-1} = \frac{2}{p} x^{(2-p)/p}$

p



20.  $\frac{d}{dx} \frac{f(x)g(x)}{f(x)g(x)^2} = \frac{f'(x)g(x) - f(x)g'(x)}{f(x)^2g(x)^2}$
21.  $\frac{d}{dx} \frac{f(x)g(x)^2}{g(x)^2} = \frac{f'(x)g(x)^2 + 2f(x)g(x)g'(x)}{g(x)^4}$
22.  $\frac{d}{dx} x^2 f(x) = 2xf(x) + x^2 f'(x)$
23.  $\frac{d}{dx} f(\sin x/g \cos x) = f'(\sin x/g \cos x) \cdot (\cos x/g \cos x - \sin x/g \sin x)$
24.  $\frac{d}{dx} \frac{\sin g(x)}{\sin g(x)} = \frac{\cos g(x)g'(x)}{\sin^2 g(x)}$
25. If  $x^3 + 2xy^3 = 12$ , then  $3x^2 + 6xy^2 = 0$ . At  $(2, 1)$ :  $12 + 6(2)(1)^2 = 24 \neq 0$ , so the slope there is  $y' = -10$ . The tangent line has equation  $y - 1 = -10(x - 2)$  or  $y = -10x + 21$ .
26.  $\frac{d}{dx} 2x \sin y = 2 \sin y + 2x \cos y y'$ . At  $(1, 3)$ ;  $1 = 4$ :  $3 \cos y + 2(1) \sin y = 0$ , so the slope there is  $y' = -\frac{3}{2}$ .
27.  $\frac{d}{dx} x^4 = 4x^3$
28.  $\frac{d}{dx} x^{-1/2} = -\frac{1}{2}x^{-3/2} = -\frac{1}{2x^{3/2}}$
29.  $\frac{d}{dx} 3 \sin x = 3 \cos x$

32. If  $g(x) = 3 \cos x$ , then  $g'(x) = -3 \sin x$ .  
If  $(2, 1)$  lies on  $y = 3 \cos x$ , then  $3 \cos x = 3$ , so  $\cos x = 1$  and  $\sin x = 0$ .  
 $\frac{d}{dx} \frac{f(x)g(x)}{g(x)^2} = \frac{f'(x)g(x) - 2f(x)g'(x)}{g(x)^3}$   
At  $(2, 1)$ :  $\frac{f'(2) \cdot 3 - 2f(2) \cdot 0}{3^3} = \frac{3f'(2)}{27} = \frac{f'(2)}{9}$
33.  $\frac{d}{dx} x \sin x = \sin x + x \cos x$
34. If  $f(x) = x^n$  and  $g(x) = x$ , then  $f'(x) = nx^{n-1}$ .  
 $\frac{d}{dx} \frac{x^n}{x} = \frac{n x^{n-1} \cdot x - x^n \cdot 1}{x^2} = \frac{n x^n - x^n}{x^2} = \frac{(n-1)x^n}{x^2} = (n-1)x^{n-2}$   
Conjecture:  $\frac{d}{dx} \frac{x^n}{x} = (n-1)x^{n-2}$  for  $n = 1, 2, 3, \dots$   
Proof: The formula is true for  $n = 1, 2$ , and  $3$  as shown above. Suppose it is true for  $n = k$ ; that is, suppose  $\frac{d}{dx} \frac{x^k}{x} = (k-1)x^{k-2}$ . Then  $\frac{d}{dx} \frac{x^{k+1}}{x} = \frac{(k+1)x^k \cdot x - x^{k+1} \cdot 1}{x^2} = \frac{(k+1)x^{k+1} - x^{k+1}}{x^2} = \frac{kx^{k+1}}{x^2} = kx^{k-1}$ . Thus the formula is also true for  $n = k + 1$ . It is therefore true for all positive integers  $n$  by induction.
35. The tangent to  $y = x^3 + 2$  at  $(a, a^3 + 2)$  has equation  $y - (a^3 + 2) = 3a^2(x - a)$ , or  $y = 3a^2x - 2a^3 + 2$ . This line passes through the origin if  $0 = 3a^2(a) - 2a^3 + 2$ , that is, if  $a = 1$ . The line then has equation  $y = 3x + 2$ .
36. The tangent to  $y = 2x^2$  at  $(a, 2a^2)$  has slope  $4a$  and equation  $y - 2a^2 = 4a(x - a)$ , or  $y = 4ax - 2a^2$ . This line passes through the origin if  $0 = 4a(a) - 2a^2$ , that is, if  $a = 1/2$ . The line then has equation  $y = 2x$ .



$y = D^p$

$$\frac{2C}{a^2} \frac{p^2 C}{a^2 x} = \frac{2C^2 p^2}{a^4 x}$$

This line passes through (0, 1/ provided

$$1 - D^p = \frac{2C^2 p^2}{a^4 x}$$

30.  $\int .2x^{-1/4} dx = \frac{2}{5} x^{3/4} + C$

or, equivalently,

$$\int .2x^{-1/4} dx = \frac{2}{5} x^{3/4} + C$$

31. If  $f(x) = \frac{1}{30} x^4 + \frac{1}{4} x^3 + C$ , then  $f'(x) = \frac{1}{3} x^3 + \frac{3}{4} x^2 + C'$ . The possibilities are  $a = 2$ , and the equations of the corresponding tangent lines are  $y = \frac{1}{3} x^3 + \frac{3}{4} x^2 + C'$ .

37.  $\frac{d}{dx} \sin^n x \sin nx / dx$

$\frac{d}{dx} \sin^{n-1} x \cos x \sin nx / \frac{d}{dx} \sin^{n-1} x \cos nx /$   
 $\frac{d}{dx} \sin^{n-1} x \cos x \sin nx / \frac{d}{dx} \sin x \cos nx /$   
 $\frac{d}{dx} \sin^{n-1} x \sin nx \cdot \frac{1}{x} /$   
 $y = \sin^n x \sin nx /$  has a horizontal tangent at  $x$

$\frac{d}{dx} \sin^n x = n \sin^{n-1} x \cos x$ , for any integer  $n$ .

38.  $\frac{d}{dx} \sin^n x \cos nx / \frac{d}{dx} \sin^n x \cos nx /$

$\frac{d}{dx} \sin^{n-1} x \cos x \cos nx / \frac{d}{dx} \sin^{n-1} x \cos nx /$   
 $\frac{d}{dx} \sin^{n-1} x \cos x \cos nx / \frac{d}{dx} \sin x \cos nx /$   
 $\frac{d}{dx} \sin^{n-1} x \cos nx \cdot \frac{1}{x} /$   
 $\frac{d}{dx} \cos^n x \sin nx /$

$\frac{d}{dx} \cos^{n-1} x \sin x \sin nx / \frac{d}{dx} \cos^{n-1} x \cos nx /$   
 $\frac{d}{dx} \cos^{n-1} x \sin x \cos nx / \frac{d}{dx} \cos x \sin nx /$   
 $\frac{d}{dx} \cos^{n-1} x \cos nx \cdot \frac{1}{x} /$   
 $\frac{d}{dx} \cos^n x \cos nx /$   
 $\frac{d}{dx} \cos^{n-1} x \sin x \cos nx / \frac{d}{dx} \cos^n x \sin nx /$   
 $\frac{d}{dx} \cos^{n-1} x \cos nx \cdot \frac{1}{x} / \frac{d}{dx} \cos x \sin nx /$

39. Q  $\frac{d}{dx} x^2 = 2x$ ;  $\frac{d}{dx} x^2 = 2x$ . If  $P(a, a^2)$  on the curve  $y = x^2$ , then the slope of  $PQ$  is  $2a$ , and the slope of  $PQ$  is  $1/a$ .

$PQ$  is normal to  $y = x^2$  if  $2a \cdot \frac{1}{a} = -1$ .  
 The points  $P$  are  $(0, 0)$  and  $(-1, 1)$ . The distances

from these points to  $Q$  are  $1$  and  $\sqrt{2}$ , respectively. The distance from  $Q$  to the curve  $y = x^2$  is the shortest of these distances, namely  $\sqrt{2}$  units.

40. The average profit per tonne if  $x$  tonnes are exported is  $P(x) = x^2$ , that is the slope of the line joining  $(0, 0)$  to  $(x, P(x))$  to the origin. This slope is maximum if the line is tangent to the graph of  $P(x)$ . In this case the slope of the line is  $P'(x)$ ,

the marginal profit.

$\frac{d}{dx} x^2 = 2x$

$2x = 2x^2$  if  $0 < x < R$

Observe that this rate is half the rate at which  $F$  decreases when  $r$  increases from  $R$ .

42.  $PV = kT$ . Differentiate with respect to  $P$  holding  $T$  constant to get

$V = k \frac{dT}{dT} = k$

Thus the isothermal compressibility of the gas is

$\frac{1}{V} \frac{dV}{dP} = \frac{1}{V} \frac{d}{dP} \left( \frac{kT}{P} \right) = -\frac{kT}{P^2}$

43. Let the building be  $h$  m high. The height of the first ball at time  $t$  during its motion is

$y_1 = h - \frac{1}{2}gt^2$

It reaches maximum height when  $\frac{dy_1}{dt} = -gt = 0$ , that is, at  $t = \sqrt{2h/g}$  s. The maximum height of the first ball is

$y_1 = h - \frac{1}{2}g \left( \frac{2h}{g} \right) = \frac{1}{2}h$

The height of the second ball at time  $t$  during its motion is

$y_2 = \frac{1}{2}gt^2$

It reaches maximum height when  $\frac{dy_2}{dt} = gt = 0$ , that is, at  $t = \sqrt{2h/g}$  s. The maximum height of the second ball is

$y_2 = \frac{1}{2}g \left( \frac{2h}{g} \right) = \frac{1}{2}h$

These two maximum heights are equal, so

$\frac{1}{2}h = \frac{1}{2}h$

- a) For continuity of  $F(r)$  at  $r = R$  we require  $mg = mkR$ , so  $k = g/R$ .
- b) As  $r$  increases from  $R$ ,  $F$  changes at rate

$$\frac{dr}{dt} = \frac{r^2}{2mgR^2} \quad \frac{dR}{dt} = \frac{R}{2mg}$$



As  $r$  decreases from  $R$ ,  $F$  changes at rate

$$\frac{dF}{dr} = \frac{mk}{r^2} \quad \frac{dF}{dt} = \frac{mk}{r^2} \frac{dr}{dt} = \frac{mk}{r^2} \frac{r^2}{2mgR^2} = \frac{mk}{2mgR^2}$$

which gives  $h = 300 = 19.6 t^2$  as the height of the building.

44. The first ball has initial height 60 m and initial velocity 0, so its height at time  $t$  is

$$y_1 = 60 - 4.9t^2 \text{ m}$$

The second ball has initial height 0 and initial velocity  $v_0$ , so its height at time  $t$  is

$$y_2 = v_0 t - 4.9t^2 \text{ m}$$

The two balls collide at a height of 30 m (at time  $T$ , say). Thus

$$30 = 60 - 4.9T^2$$

$$30 = v_0 T - 4.9T^2$$

Thus  $v_0 T = 60$  and  $T^2 = 30 = 4:9$ . The initial upward speed of the second ball is

$$v_0 = \frac{60}{T} = \frac{60}{\sqrt{30}} = 24\sqrt{5} \text{ m/s}$$

At time  $T$ , the velocity of the first ball is

$$\frac{dy}{dt} = -9.8T = -24\sqrt{5} \text{ m/s}$$

At time  $T$ , the velocity of the second ball is

$$\frac{dy}{dt} = v_0 - 9.8T = 0 \text{ m/s}$$

45. Let the car's initial speed be  $v_0$ . The car decelerates at  $20 \text{ ft/s}^2$  starting at  $t = 0$ , and travels distance  $s$  in time  $t$ ,

where  $d^2s/dt^2 = -20$ . Thus

$$\frac{ds}{dt} = v_0 - 20t$$

$$s = v_0 t - 10t^2$$

The car stops at time  $t = v_0/20$ . The stopping distance is

$s = 160$  ft, so

$$160 = v_0 \frac{v_0}{20} - 10 \left(\frac{v_0}{20}\right)^2$$

The car's initial speed cannot exceed  $p$

$v_0 = 160 \sqrt{40} = 80 \sqrt{40} \text{ ft/s}$ .

46.  $P = 2L = gL^2$

a) If  $L$  remains constant, then

$$\frac{dP}{dL} = 2g = 2L = 2g$$

$$\frac{dL}{dP} = \frac{1}{2g}$$

### Challenging Problems 2 (page 164)

1. The line through  $(a, a^2)$  with slope  $m$  has equation

$y - a^2 = m(x - a)$ . It intersects  $y = x^2$  at points  $x$  that satisfy

$$x^2 - a^2 = mx - ma$$

$$x^2 - mx + (ma - a^2) = 0$$

In order that this quadratic have only one solution  $x = a$ , the left side must be  $(x - a)^2$ , so that  $m = 2a$ . The tangent has slope  $2a$ .

This won't work for more general curves whose tangents can intersect them at more than one point.

2.  $f'(x) = x^2 - 9$

$$a) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - 9 - (x^2 - 9)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = \lim_{h \rightarrow 0} (2x + h) = 2x$$

$$b) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - 9 - (x^2 - 9)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = \lim_{h \rightarrow 0} (2x + h) = 2x$$

$$b) \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - 9 - (x^2 - 9)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = \lim_{h \rightarrow 0} (2x + h) = 2x$$

$$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - 9 - (x^2 - 9)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = \lim_{h \rightarrow 0} (2x + h) = 2x$$

$$D_{h \rightarrow 0} \frac{1}{h} = \frac{1}{h^2}$$

$$D_{f'} = \frac{2}{6} = \frac{1}{3}$$

3.  $f'(x) = 4x^3, g'(x) = 4x^3, g(x) = x^4$  if  $x \neq 4$ .

$$\lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{4x^3 - 4^3}{x - 4}$$

$$a) x \rightarrow 4, f(x) = 4x^3, g(x) = x^4$$

$$D_{f'} = \frac{4}{4} = 1, D_{g'} = \frac{4}{4} = 1$$

If  $g$  increases by 1%, then  $g = g + 0.01g$ , and  $P = P + 0.01P$ . Thus  $P$  increases by 0.5%.

b) If  $g$  remains constant, then

$$\frac{dP}{P} = \frac{dL}{L} + \frac{dL}{L} = 2 \frac{dL}{L}$$

If  $L$  increases by 2%, then  $L = L + 0.02L$ , and  $P = P + 0.04P$ . Thus  $P$  increases by 4%.

If  $L$  increases by 2%, then  $L = L + 0.02L$ , and  $P = P + 0.04P$ . Thus  $P$  increases by 4%.

$$\lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{1 - 3} = \frac{1}{2}$$

$$\lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{1 - 3} = \frac{1}{2}$$

$$\lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{1 - 3} = \frac{1}{2}$$

e)  $\lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4}$

f)  $\lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4} = \lim_{x \rightarrow 4} \frac{f(x) - f(4)}{x - 4}$

4. f(x) =  $\begin{cases} x & \text{if } x \in \mathbb{N}; 1=2, 1=3, \dots \\ nx^2 & \text{otherwise} \end{cases}$   
 a) f is continuous except at  $x=2, 1=3, 1=4, \dots$ . It is continuous at  $x \in \mathbb{N}$  and  $x \in \mathbb{Q}$  (and everywhere else).

Note that

$\lim_{x \rightarrow 1} x^2 = 1 = f(1)$   
 $\lim_{x \rightarrow 0} x^2 = 0 = f(0)$

b) If  $a \in \mathbb{N}$  and  $b \in \mathbb{N}$ , then

$\frac{f(a) - f(b)}{a - b} = \frac{a^2 - b^2}{a - b} = a + b$

If  $1=3 < x < 1=2$ , then  $f(x) = x^2 < 1=4 < 5=12$ . Thus the statement is FALSE.

c) By (a) f cannot be differentiable at  $x \in \mathbb{N}$ ,  $1=2, 1=2, \dots$ . It is not differentiable at  $x \in \mathbb{Q}$  either, since

$\lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = \lim_{h \rightarrow 0} \frac{h^2}{h} = 0$

f is differentiable elsewhere, including at  $x \in \mathbb{N}$  where its derivative is 2.

5. If  $h \neq 0$ , then

7. Given that  $g'(0) = k$  and  $g(x) = C y + D g(x) + C g(y)$ , then

a)  $g'(0) = D g'(0) + C g'(0) = C g'(0)$ . Thus  $g'(0) = D 0$ .

b)  $g'(x) = D \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h}$

c) If  $h(x) = D g(x) + kx$ , then  $h'(x) = D g'(x) + k = D 0$

for all x. Thus h(x) is constant for all x. Since  $h(0) = D g(0) + 0 = D 0$ , we have  $h(x) = D 0$  for all x, and  $g(x) = D kx$ .

8.a)  $f'(x) = D \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$D \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$

$f'(x) = D \frac{1}{2} f'(x) + C f'(x)$

$C \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h}$

$D \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h} = \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h}$

b) The change of variables used in the first part of (a) shows that

$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{f(x) - f(x-h)}{h}$  are always equal if either exists.

c) If  $f'(x) = x$ , then  $f'(0) = 0$  does not exist, but

D j j

$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{h - h}{h} = \lim_{h \rightarrow 0} 0 = 0$

9. The tangent to  $y = D x^3$  at  $x = D 3a = 2$  has equation  $y = D 8 C 4a^2 x - 2$

$$\frac{f(h) - f(0)}{h} = \frac{f(h) - f(0)}{h} > |f(h) - f(0)|$$

as  $h \rightarrow 0$ . Therefore  $f'(0)$  does not exist.

6. Given that  $f'(0) = k$ ,  $f'(0) \neq 0$ , and  $f$

is concave down, we have

$$f'(0) < f'(x) < f'(0) \quad \text{or} \quad f'(0) > f'(x) > f'(0)$$

Thus  $f'(0) < 1$ .

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h}$$

$$\lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = k < 1$$

$$\frac{27a^3}{27} = \frac{27}{3a}$$

This line passes through  $(a, 0)$  because

$$\frac{8}{27a^3} = \frac{4a^2}{27} = \frac{a}{3a} = \frac{2}{3a}$$

If  $a \neq 0$ , the x-axis is another tangent to  $y = x^3$  that passes through  $(a, 0)$ .

The number of tangents to  $y = x^3$  that pass through  $(x_0, y_0)$  is

three, if  $x_0 \neq 0$  and  $y_0$  is between 0 and  $x_0^3$ ;

two, if  $x_0 \neq 0$  and either  $y_0 = 0$  or  $y_0 = x_0^3$ ;

one, otherwise.

This is the number of distinct real solutions  $b$  of the cu-bic

equation  $2b^3 - 3b^2 - x_0 C y_0 D 0$ , which states that the tangent to  $y = Dx^3$  at  $(b, b^3)$  passes through  $(x_0, y_0)$ .

10. By symmetry, any line tangent to both curves must pass through the origin.




Fig. C-2-10

The tangent to  $y = Dx^2 + C4x + C1$  at  $x = a$  has equation

$$y = D a^2 C 4a C 1 C .2a C 4/x a/ D .2a C 4/x .a^2 1/;$$

which passes through the origin if  $a = 1$ . The two common tangents are  $y = D6x$  and  $y = D2x$ .

11. The slope of  $y = Dx^2$  at  $x = a$  is  $2a$ . The slope of the line from  $(0, b)$  to  $(a, a^2)$  is  $a^2/b = a$ .

This line is normal to  $y = Dx^2$  if either  $a = 0$  or  $2a \cdot a = b/a$ , that is, if  $a = 0$  or  $2a = D2b = 1$ .

There are three real solutions for  $a$  if  $b > 1=2$  and only one ( $a = 0$ ) if  $b = 1=2$ .

12. The point  $Q = (a, a^2)$  on  $y = Dx^2$  that is closest to

$P = (3, 0)$  is such that  $PQ$  is normal to  $y = Dx^2$  at  $Q$ . Since  $PQ$  has slope  $a^2 = a$  and  $y = Dx^2$  has slope  $2a$  at  $Q$ , we require

$$\frac{a^2}{a} = -\frac{1}{2a};$$

which simplifies to  $2a^3 - C a - 3 = 0$ . Observe that  $a = 1$  is a solution of this cubic equation. Since the slope of  $y = Dx^3 + Cx + 3$  is  $6x^2 + C$ , which is always positive, the cubic equation can have only one real solution. Thus

The curve  $y = Dx^2 + Cx + C$  has slope  $m = D2Aa + C/B$  at  $x = a$ ;  $Aa = C/B + C/C$ . Thus  $a = D \cdot m = B/2A$ , and the tangent has equation

$$y = D Aa^2 C B a C C C m x a/ D m x C \frac{m \cdot B/2}{4A} C \frac{B \cdot m \cdot B/2}{2A} C C \frac{m \cdot m \cdot B/2}{2A} D m x C C C \frac{m \cdot B/2}{4A} \frac{m \cdot B/2}{2A} D m x C f \cdot m/;$$

where  $f = m/D C$ ,  $m = B/2 = 4A/2$ .  
14. Parabola  $y = Dx^2 + Cx + C$  has tangent  $y = 2ax + a^2$  at  $(a, a^2)$ .

Parabola  $y = Dx^2 + Cx + C$  has tangent

$$y = D \cdot 2Ab C B/x + Ab^2 C C$$

at  $(b, Ab^2 C B b C C)$ . These two tangents coincide if

$$\frac{2Ab C B D 2a}{Ab^2 C D a^2} = 1/$$

The two curves have one (or more) common tangents if  $b$  has real solutions for  $a$  and  $b$ . Eliminating  $a$  between the two equations leads to

$$.2Ab C B/2 D 4Ab^2 - 4C;$$

or, on simplification,

$$4A \cdot A - 1/b^2 C 4AB b C \cdot B^2 C 4C / D 0;$$

This quadratic equation in  $b$  has discriminant

$$D D 16A B - 16A \cdot A - 1/B C 4C / D 16A \cdot B - 4 \cdot A - 1/C /;$$

There are five cases to consider:

CASE I. If  $A = 1, B \neq 0$ , then  $b = 1$  gives

$$\frac{b}{a} = \frac{B^2 C 4C}{D \cdot 4B};$$

There is a single common tangent in this case.

CASE II. If  $A = 1, B = 0$ , then  $b = 1$  forces  $C = 0$ , which is not allowed. There is no common tangent in this case.

CASE III. If  $A \neq 1$  but  $B^2 = D 4 \cdot A - 1/C$ , then

$$\frac{B}{B}$$



Q D .1; 1/ is the point on y D x<sup>2</sup> that is closest to P .

P

The distance from P to the curve is jP Qj D 5 units.

13. The curve y D x<sup>2</sup> has slope m D 2a at .a; a<sup>2</sup>/. The tangent there has equation

$$m^2$$

$$y D a^2 C m.x - a/ D mx - 4:$$

$$b D 2.A - 1/ D a:$$

There is a single common tangent, and since the points of tangency on the two curves coincide, the two curves are tangent to each other.

CASE IV. If A ≠ 1 and B<sup>2</sup> - 4.A - 1/C < 0, there are no real solutions for b, so there can be no common tangents.

CASE V. If  $A \neq 1$  and  $B^2 - 4A \cdot 1/C > 0$ , there are two distinct real solutions for  $b$ , and hence two common tangent lines.




Fig. C-2-14

15. a) The tangent to  $y = Dx^3$  at  $(a, a^3)$  has equation

$$y = D(3a^2x - 2a^3)$$

For intersections of this line with  $y = Dx^3$  we solve

$$x^3 - 3a^2x + 2a^3 = 0$$

$$(x - a)^2(x + 2a) = 0$$

The tangent also intersects  $y = Dx^3$  at  $(b, b^3)$ , where

$$b = 2a$$

- b) The slope of  $y = Dx^3$  at  $x = 2a$  is  $3(2a)^2 = 12a^2$ , which is four times the slope at  $x = a$ .

- c) If the tangent to  $y = Dx^3$  at  $x = a$  were also tangent at  $x = b$ , then the slope at  $b$  would be four times that at  $a$  and the slope at  $a$  would be four times that at  $b$ . This is clearly impossible.

- d) No line can be tangent to the graph of a cubic polynomial  $P(x)$  at two distinct points  $a$  and  $b$ , because if there was such a double tangent  $y = L(x)$ , then

- b) The tangent to  $y = Dx^4 - 2x^2$  at  $x = a$  has equation

$$y = D(4a^3x - 4a) + 2a^2 - 4a^2/x$$

Similarly, the tangent at  $x = b$  has equation

$$y = D(4b^3x - 4b) + 2b^2 - 4b^2/x$$

These tangents are the same line (and hence a double tangent) if

$$4a^3 = 4b^3 \text{ and } 2a^2 - 4a^2/x = 2b^2 - 4b^2/x$$

$$a^3 = b^3 \text{ and } a^2 - 2a^2/x = b^2 - 2b^2/x$$

The second equation says that either  $a^2 = b^2$  or  $a^2 - 2a^2/x = b^2 - 2b^2/x$

$a^2 = b^2$ ; the first equation says that

$a = 2b$  or  $a = -2b$ , or, equivalently,  $a = 2b$  or  $a = -2b$ . Then  $a = 2b$  (if  $a = -2b$  is not allowed).

Thus  $a = 2b$  and the two points are  $(a, 1)$  and  $(2a, 1)$  as discovered in part (a).

If  $a^2 - 2a^2/x = b^2 - 2b^2/x$ , then  $ab = 1$ . This is not possible

since it implies that

$$y = Dx^4 - 2x^2$$

Thus  $y = 1$  is the only double tangent to

- c) If  $y = Dx^2 + Ax + B$  is a double tangent to  $y = Dx^3 + Cx$ , then  $y = Dx^2 + Ax + B$  is a double tangent to

$$y = Dx^4 - 2x^2$$

and  $B = 1$ . Thus the only double tangent to  $y = Dx^4 - 2x^2 + Cx$  is  $y = Dx^2 + Ax + 1$ .

17. a) The tangent to

$$y = Df(x) = D(ax^4 + bx^3 + cx^2 + dx + e)$$

at  $x = p$  has equation

$$y = D(4ap^3 + 3bp^2 + 2cp + d) + f(p) - f'(p)(x - p)$$

This line meets  $y = Df(x)$  at  $x = p$  (a double root), and

$\frac{ax^2 + b}{x^2}$  would be a factor of the cubic polynomial  $P(x) = L(x)$ , and cubic polynomials do not have factors that are 4th degree polynomials.

16. a)  $y = x^4 - 2x^3$  has horizontal tangents at points  $x$  satisfying  $4x^3 - 6x^2 = 0$ , that is, at  $x = 0$  and  $x = \frac{3}{2}$ . The horizontal tangents are  $y = 0$  and  $y = \frac{27}{8}$ . Note that  $y = \frac{27}{8}$  is a double tangent; it is tangent at the two points  $(\frac{3}{2}, \frac{27}{8})$  and  $(\frac{3}{2}, \frac{27}{8})$ .

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

These two latter roots are equal (and hence correspond to a double tangent) if the expression under the square root is 0, that is, if

$$b^2 - 4ac = 0$$

This quadratic has two real solutions for  $p$  provided its discriminant is positive, that is, provided

$$16a^2 b^2 - 4.8a^2 / .4ac - b^2 / > 0:$$

This condition simplifies to

$$3b^2 > 8ac:$$

For example, for  $y = x^4 - 2x^2 + 1$ , we have  $a = 1$ ,  $b = 0$ , and  $c = 2$ , so  $3b^2 = 0 > 16 = 8ac$ ,

and the curve has a double tangent.

- b) From the discussion above, the second point of tangency is

$$q = \frac{2ap - b}{2a} = p - \frac{b}{2a}:$$

The slope of  $PQ$  is

$$\frac{f(q) - f(p)}{q - p} = \frac{b^3 - 4abc + 8a^2 d}{8a^2}:$$

Calculating  $f'(p) = 4p^3 - 4p$  leads to the same expression, so the double tangent  $PQ$  is parallel to the tangent at the point horizontally midway between  $P$

and  $Q$ .

- c) The inflection points are the real zeros of  $f''(x) = 12ax^2 - 6b$ :

$$x = \pm \sqrt{\frac{b}{2a}}:$$

This equation has distinct real roots provided  $9b^2 > 24ac$ , that is,  $3b^2 > 8ac$ . The roots are

$$r = \sqrt{\frac{3b}{12a} + \frac{\sqrt{9b^2 - 24ac}}{12a}}$$

$$s = \sqrt{\frac{3b}{12a} - \frac{\sqrt{9b^2 - 24ac}}{12a}}$$

The slope of the line joining these inflection points is

$$\frac{f(s) - f(r)}{s - r} = \frac{b^3 - 4abc + 8a^2 d}{8a^2}:$$

so this line is also parallel to the double tangent.

so the formula above is true for  $n = 1$ . Assume it is true for  $n = k$ , where  $k$  is a positive integer. Then

$$\frac{d^{k+1}}{dx^{k+1}} \cos ax = -a \frac{d^k}{dx^k} \cos ax = -a^k \sin ax$$

Thus the formula holds for  $n = 2, 3, \dots$  by induction.

- b) Claim:  $\frac{d^n}{dx^n} \sin ax = a^n \sin ax$

Proof: For  $n = 1$  we have

$$\frac{d}{dx} \sin ax = a \cos ax = a \sin ax$$

so the formula above is true for  $n = 1$ . Assume it is true for  $n = k$ , where  $k$  is a positive integer. Then

$$\frac{d^{k+1}}{dx^{k+1}} \sin ax = a \frac{d^k}{dx^k} \sin ax = a^k \sin ax$$

Thus the formula holds for  $n = 1, 2, 3, \dots$  by induction.

- c) Note that

$$\frac{d}{dx} \cos^4 x = -4 \cos^3 x \sin x$$

$$\frac{d}{dx} \sin^4 x = 4 \sin^3 x \cos x$$

$$\frac{d}{dx} \sin 4x = 4 \cos 4x$$

18. a) Claim:  $\frac{d^n}{dx^n} \cos(ax) = a^n \cos(ax)$  if  $n$  is even, and  $\frac{d^n}{dx^n} \cos(ax) = -a^n \sin(ax)$  if  $n$  is odd.

Proof: For  $n \geq 1$  we have


$$\frac{d}{dx} \cos(ax) = -a \sin(ax)$$

$$\frac{d}{dx} \sin(ax) = a \cos(ax)$$

It now follows from part (a) that

$$\frac{d^n}{dx^n} \cos(ax) = a^n \cos(ax)$$

$$\frac{d^n}{dx^n} \sin(ax) = a^n \sin(ax)$$



- a) The fuel lasted for 3 seconds.
- b) Maximum height was reached at  $t = 3$  s.
- c) The parachute was deployed at  $t = 12$  s.

d) The upward acceleration in  $0 \leq t \leq 3$  was  $30/3 = 10$  m/s<sup>2</sup>.

e) The maximum height achieved by the rocket is the distance it fell from  $t = 3$  to  $t = 15$ . This is the area under the  $t$ -axis and above the graph of  $v$  on that interval, that is,

$$\frac{1}{2} \cdot 12 \cdot 30 = 180 \text{ m}$$

f) During the time interval  $0 \leq t \leq 7$ , the rocket rose a distance equal to the area under the velocity graph and above the  $t$ -axis, that is,

$$\frac{1}{2} \cdot 7 \cdot 30 = 105 \text{ m}$$

Therefore the height of the tower was  $105 - 137 = -32$  m.

