Solution Manual for Chemistry in Focus A Molecular View of Our World 6th Edition by Nivaldo J Tro ISBN 1305084470 9781305084476

Full Link download Solution Manual

https://testbankpack.com/p/solution-manual-for-chemistry-in-focus-a-molecular-view-of-our-world-6th-edition-by-nivaldo-j-tro-isbn-1305084470-9781305084476/

Test Bank

https://testbankpack.com/p/test-bank-for-chemistry-in-focus-a-molecular-view-of-our-world-6th-edition-by-nivaldo-j-tro-isbn-1305084470-9781305084476/

CHAPTER 2

THE CHEMIST'S TOOL BOX

ANSWERS TO QUESTIONS:

- Curiosity is an important part of the scientific enterprise because scientists need a strong
 desire to investigate and learn about the behavior of nature. Science must start with the question
 why. The scientific method is then utilized to accumulate systematized knowledge about the
 physical world. A scientist's curiosity is incapable of being satisfied. Without the curious
 nature of scientists, the advancement of science would not have occurred as we presently know
 it.
- 2. Science is characterized by observations, which are used to develop experimental laws and theories for the workings of nature. Measurement lies at the heart of experiment. Nature appears to be based on a foundation that can be described by mathematics, and measurements provide the bricks to build upon this foundation.
- 3. Measured quantities are written so that the uncertainty is contained within the last digit of the number. A volume of 30.0 mL means the volume lies in the range 29.9-30.1 mL (30.0 ± 0.1 mL).
- 4. The difference between the quantities 9 inches and 9.00 inches is the amount of certainty or precision. Although numerically equivalent (the results of using 9 inches or 9.00 inches in a calculation will be indistinguishable), 9 inches tells us it is only certain in the range 8-10 inches; the quantity 9.00 inches is much more precise, having a certainty in the much narrower range of 8.99-9.01 inches.
- 5. Any measurement consists of a numerical value and the chosen unit. Units inform what is being measured and the scale used. Without a unit, a measurement is virtually useless. For example, if I say it is thirty degrees outside, the response will be thirty what? An American would presume it is cold since he is familiar with Fahrenheit; a European would turn on the air-conditioning presuming the measurement to be in Celsius. The International System of units, or SI, is generally used in the scientific arena.
- 6. Answers may vary. Three possible units for length are millimeters, centimeters and meters. Examples:

Thickness of a dime - millimeters (mm) Length of a finger - centimeters (cm) Height of an adult - meters (m)

7. Answers may vary. Three possible units for mass are grams, milligrams and kilograms.

Examples:

Mass of a penny - grams (g) Mass of a straight pin - milligrams (mg)

Mass of a bucket of water - kilograms (kg)

8. Answers may vary. Three possible units for time are milliseconds, seconds, and microseconds.

Examples:

Time between heartbeats - milliseconds (ms)

Time to run the 100 m. dash - seconds (s)

Time to blink your eye - microseconds (s)

9. Answers may vary. Three possible units for volume are milliliters, kiloliters (gallons), and liters.

Examples:

Volume of a child's juice box – milliliters (mL)

Volume of water in a swimming pool – gallons or kiloliters (kL)

Volume of a bottle of soda – liters (L)

- 10. A conversion factor is an equivalence statement that relates one unit to another. Conversion factors are commonly written in equation form or as a fraction with units in the numerator and denominator. Because it is an equivalence statement, the fraction equals one and the conversion factor can be multiplied by any number with out changing the value of the number, only its units.
- 11. Graphs are a very convenient and powerful way to illustrate relationships between different quantities. Graphs can be modified to emphasize particular features. It is important to examine the range on the *y* axis to understand the significance of the changes plotted. A change will look much bigger if the scale is smaller, whereas a small change on a large scale could go virtually unnoticed.

12.

- a) The decimal part of the number is 9.66
- b) The exponential part of the number is 10^{-5}
- c) The exponent is -5.
- 13. Density is defined as mass per unit volume. Typical units for density are g/cm³ (commonly used for solids), g/mL (used for liquids), or kg/m³ (used for gases).
- 14. Oil floating on water means that the density of the oil is less than that of the water. Denser substances will sink in less dense substances.

SOLUTIONS TO PROBLEMS:

15.

- a) $8.51 \times 10^{-4} \text{ g}$
- b) 3.8041430 x 10⁷
- c) $2.9979 \times 10^8 \text{ m/s}$
- d) 3.13914040 x 10⁸

- a) 1.9570261 x 10⁷
- b) 7.190900000 x 10⁹
- c) 7.461 x 10⁻¹¹ m
- d) $1.5 \times 10^{-5} \text{ m}$

17.

- a) 149,000,000 km
- b) 0.000000000079 m
- c) 4,540,000,000 yr
- d) 6,400,000 m

18.

- a) 602,200,000,000,000,000,000,000 carbon atoms in 12.01 grams of carbon
- b) 300,000,000 m/s
- c) 0.000000450 m
- d) 13,700,000,000 yr
- 19. Convert 40,075 km to a) meters, b) miles, c) feet.

a)
$$4.0075 \times 10^{7} \text{ m}$$
 $\frac{40075 \text{ km}}{1} = \frac{1000 \text{ m}}{1 \text{ km}} = 4.0075 \times 10^{7} \text{ m}$
b) $24,900 \text{ or } 2.490 \times 10^{4} \text{ mi}$ $\frac{40075 \text{ km}}{1} = \frac{0.6214 \text{ mi}}{1 \text{ km}} = 2.490 \times 10^{4} \text{ mi}$
c) $1.315 \times 10^{8} \text{ ft}$ $\frac{40075 \text{ km}}{1} = \frac{0.6214 \text{ mi}}{1 \text{ km}} = \frac{5280 \text{ ft}}{1.315 \times 10^{8} \text{ ft}}$
 $\frac{40075 \text{ km}}{1 \text{ km}} = \frac{1000 \text{ m}}{1 \text{ km}} = \frac{4.0075 \text{ km}}{1 \text{ mi}} = \frac{1000 \text{ m}}{1 \text{ km}} = \frac{4.0075 \text{ km}}{1 \text{ mi}} = \frac{1000 \text{ m}}{1 \text{ km}} = \frac{4.0075 \text{ m}}{1 \text{ mi}} = \frac{1000 \text{ m}}{1 \text{ km}} = \frac{10000 \text{ m}}{1 \text{ km}} = \frac{10000 \text{ m}}{1 \text{ km}} = \frac{10000$

20. Convert 2,777 miles into a) kilometers, b) meters, c) feet.

a)
$$4469 \text{ km}$$
 $\frac{2777 \text{ mi} \times 1 \text{ km}}{0.6214 \text{ mi}} = 4469 \text{ km}$
 0.6214 mi
 $\frac{2777 \text{ mi}}{1} = 1 \text{ km} = 1000 \text{ m}$
b) $4.469 \times 10^6 \text{ m}$ $\frac{10.6214 \text{ mi}}{1 \times 1000 \text{ m}} = 1 \text{ km} = 4.469 \times 10^6 \text{ m}$
 $\frac{2777 \text{ mi}}{1 \times 1000 \text{ m}} = \frac{1760 \text{ yd}}{1 \times 1000 \text{ m}} = \frac{3 \text{ ft}}{1 \times 1000 \text{ m}} = \frac{1 \text{ mi}}{1 \times 1000 \text{ m}} = \frac{1 \text{ mi}}{$

- 27 mi 1 km 43 km 23. Convert miles to kilometers: 1 km = 0.6214 mi0.6214 mi Efficiency is 43 km/gal
- 1 1km Car will travel 11 miles on one liter of fuel. 25.

a)
$$4332 \text{ mm} = 4.332 \text{ m}$$
 $\frac{4332 \text{ mm}}{1} = \frac{1 \text{ m}}{1000 \text{ mm}} 4.332 \text{ m}$

b)
$$1.76 \text{ kg} = 1760 \text{ g or } 1.76 \text{ x } 10^3 \text{ g}$$
 $1 \text{ kg} = 1,760 \text{ g}$

c)
$$4619 \text{ mg} = 4.619 \text{ x } 10^{-3} \text{ kg}$$
 $\frac{4619 \text{ mg}}{1} = \frac{1 \text{ kg}}{1000 \text{ mg}} = 4.619 \text{ x } 10^{-3} \text{ kg}$

d)
$$0.0117 L = 11.7 mL$$
 $0.0117 L = 1000 mL$ $1 L$ $1 L$ $26.$

b)
$$4912.5 \text{ g} = 4.9125 \text{ kg}$$
 $\frac{4912.5 \text{ g}}{1}$ $\frac{1 \text{ kg}}{1000 \text{ g}}$ 4.9125 kg

c) 23.1 cm = 231 mm
$$\frac{23.1 \text{ cm}}{1} \frac{10 \text{ mm}}{1 \text{ cm}} 231 \text{ mm}$$

d)
$$561 \text{ mL} = 0.561 \text{ L}$$
 $\frac{561 \text{ mL}}{1} = \frac{1 \text{ L}}{1000 \text{ mL}} = 0.561 \text{ L}$

a)
$$358 \text{ m} = 1170 \text{ ft (or } 1.17 \text{ x } 10^3 \text{ ft)}$$

$$\frac{358 \text{ m}}{1} = \frac{39.37 \text{ in}}{1 \text{ m}} = 15t = 1170 \text{ ft}$$
b) $10.0 \text{ km} = 6.21 \text{ mi}$

$$\frac{10.0 \text{ km}}{1} = \frac{0.6214 \text{ mi}}{1 \text{ km}} = 6.21 \text{ mi}$$

c)
$$1.55 \text{ m} = 61.0 \text{ in}$$
 $\frac{1.55 \text{ m}}{1 \text{ m}} = 61.0 \text{ in}$

d)
$$23 \text{ cm} = 9.1 \text{ in}$$
 $\frac{23 \text{ cm}}{1} = \frac{1 \text{ in}}{2.54 \text{ cm}} = 9.1 \text{ in}$

a)
$$4.92 \text{ in} = 125 \text{ mm}$$
 $\frac{4.92 \text{ in}}{1} = \frac{25.4 \text{ mm}}{1 \text{ in}} = 125 \text{ mm}$

b)
$$8779 \text{ yd} = 8.027 \text{ km}$$
 $\frac{8779 \text{ yd}}{1}$ $\frac{1 \text{ mi}}{1760 \text{ yd}}$ $\frac{1 \text{mi}}{0.6214 \text{ km}}$ 8.027 km

c)
$$87 \text{ ft} = 27 \text{ m}$$
 $\frac{87 \text{ ft}}{1 \text{ ft}} = \frac{12 \text{ in}}{39.37 \text{ in}} = 27 \text{ m}$

d)
$$3.8 \text{ in} = 9.7 \text{ cm}$$
 $3.8 \text{ in} = 9.7 \text{ cm}$ 1 lin 9.7 cm

29.

a)
$$1552 \text{ m}^2 = 1.671 \times 10^4 \text{ ft}^2$$

$$\frac{1552 \text{ m}^2}{1} \frac{(39.37 \text{ in})_2}{1 \text{ m}^2} \frac{1 \text{ ft}^2}{(12 \text{ in})^2} \frac{4}{1.671 \times 10} \frac{2}{1}$$

$$\frac{1552 \text{ m}^2}{1} \frac{1 \text{ km}^2}{1} \frac{1 \text{ km}^2}{1} \frac{2}{1.671 \times 10} \frac{2}{1}$$

b)
$$1552 \text{ m}^2 = 1.552 \times 10^{-3} \text{ km}^2$$
 1 $(1000 \text{ m})^2$ 1.552x10 km $\frac{1552 \text{ m}^2}{}$ $\frac{(10 \text{ dm})^2}{}$

c)
$$1552 \text{ m}^2 = 1.552 \times 10^5 \text{ dm}^2$$
 1 1 m² 1.552×10⁵ dm²

30.

$$54 \,\mathrm{cm}_3$$
 $(10 \,\mathrm{mm})_3$

a)
$$54 \text{ cm}^3 = 5.4 \times 10^4 \text{ mm}^3$$

b) $54 \text{ cm}^3 = 3.8 \text{ in}^3$

$$1 \quad 1 \text{ cm}^3 \quad 5.4 \times 10^4 \text{ mm}^3$$

$$\frac{54 \text{ cm}_3}{1} \quad \frac{1 \text{ in}^3}{(2.54 \text{ cm})^3} \quad 3.3 \text{ in}$$

c)
$$54 \text{ cm}^3 = 5.4 \times 10^{-2} \text{ dm}^3$$
 $1 \quad (10 \text{ cm})^3 \quad 5.4 \times 10 \quad \text{dm}$

31.

a) 1 square kilometer (km²) =
$$10^6$$
 square meters (m²). (1 km = 10^3 m)
b) 1 ft³ = 2.83×10^4 cm³. $\frac{1 \text{ ft }^3}{1 \text{ ft }^3} \frac{(12 \text{ in})^3}{1 \text{ ft }^3} \frac{(2.54 \text{ cm})^3}{1 \text{ in }^3} 2.83 \times 10^4 \text{ cm}^3$

c)
$$1 \text{ yd}^2 = 9 \text{ ft}^2$$
. $(1 \text{ yd} = 3 \text{ ft})$

32.

a) 1 square meter (
$$m^2$$
) = 10^4 square centimeters (cm^2). (1 m = 10^2 cm)

b) 1 cubic yard (yd₃) =
$$4.6656 \times 10^4 \text{ in}_3 \frac{1 \text{ yd}_3 (3 \text{ ft})_3 (12 \text{ in})_3}{4.6656}$$

$$x 10^4 in 3 11 yd^3 1 ft^3$$

c) 1 square foot (ft²) = 929 square centimeters (cm²)

$$\frac{1 \text{ ft}^2}{1 \text{ ft}^2} = \frac{(12 \text{ in})^2}{1 \text{ or}^2} = \frac{(2.54 \text{ cm})^2}{1 \text{ or}^2} = 9.29 \text{ x } 10^2 \text{ cm}^2$$

33. 53 min Time = distance/speed
$$\frac{10 \text{ km}}{1} = \frac{0.6214 \text{ mi}}{1 \text{ km}} = \frac{8.5 \text{ min}}{1 \text{ mi}} = 53 \text{ min}$$

$$\frac{12 \text{ mi}}{1 \text{ gal}} = \frac{1 \text{ km}}{0.6214 \text{ mi}} = \frac{1 \text{ gal}}{3.785 \text{ L}} = 5.1 \text{ km/L}$$

- a) The total decrease in the carbon monoxide is found by subtracting the final concentration in 2012 from the initial concentration in 1990.
 6.1 ppm 1.5 ppm = 4.6 ppm
- b) The average yearly decrease is found by dividing the total decrease by the total number of years (1990-2012=22 years)4.6 ppm/22 years = 0.21 ppm/yr
- c) The total percentage decrease is found by dividing the total decrease by the concentration in 1990 and multiplying by 100 % $\frac{4.6.6}{1}$ 100 % 75 %
 - d) To find the average yearly percentage decrease, the total percentage decrease (75 %) is divided by the number of years in the period (22 years):

 3.4 %/yr

 22 yr

38.

a) The total increase in the carbon dioxide is found by subtracting the initial concentration in 1950 from the final concentration in 2007.
 382 ppm – 310 ppm = 72 ppm

```
b) tal increase by the total number of years (1950 - 2007 = 57 \text{ years}) T 72 \text{ ppm/}57 \text{ years} = 1.3 \text{ ppm/}yr h
```

e

a

v e

r

a g

e e

y

e

a r

1

y

n c

r

e a

S

e i

s

f

O

u

n d

u b

y

 $\begin{array}{c} d \\ i \end{array}$

V i

d i

n

g t

h

e t

О

16

- c) The total percentage increase is found by dividing the total increase by the concentration in 1950 and multiplying by 100 % $\overline{310}$ ppm ppm x 100 %=23 %
- d) To find the average yearly percentage increase, the total percentage increase (23 %) is divided by the number of years in the period (57 years) $\frac{23 \%}{57 \text{ yr}} = 0.41 \%/\text{yr}$
- 39. The density is determined by dividing the mass by the volume: $127.8 \text{ g}/28.4 \text{ cm}^3 \text{ Density of titanium} = 4.50 \text{ g/cm}^3$
- 40. The density is determined by dividing the mass by the volume: $3.5 \text{ g/}1.5 \text{ cm}^3 \text{ Density of silicon} = 2.3 \text{ g/cm}^3$
- 41.1.26 g/mL Density is mass/volume: $\frac{6.30 \times 10^3 \text{ g}}{1.26 \text{ g/mL}} = \frac{1 \text{ L}}{1.26 \text{ g/mL}}$

1 5 L 1000 mL

42. 13.5 g/mL Density is mass/volume: $\frac{51.4 \text{ g}}{3.80 \text{ mL}} = 13.5 \text{ g/mL}$

43.

a)
$$42.7 \text{ g}$$
 38.5 mL $\frac{1 \text{ cm}^3}{2}$ $\frac{1.11 \text{ g}}{2}$ 42.7 g

b) 3.2 L 3.5 kg
$$\frac{1 \text{ mL}}{1 \text{ kg}} \times \frac{1 \text{ cm}^3}{1 \text{ 1g}} \times \frac{1 \text{ mL}}{1 \text{ cm}^3} \times \frac{1 \text{ LL}}{1 \text{ LL}} \times$$

44.

a) Mass of Gold =
$$6.8 \times 10^3 \text{ g} 350 \text{ mL} \frac{1 \text{cm}_3}{1 \text{mL 1cm}} \frac{19.32 \text{ g}}{1 \text{mL 1cm}} 6.8 \times 10^3 \text{ g}$$

Mass of Sand =
$$1.0 \times 10^3 \text{ g}$$
 350 mL $\frac{1 \text{cm}_3}{1 \text{mL 1cm}_3}$ $\frac{3.00 \text{g}}{1 \text{mL 1cm}_3}$

- b) Yes, the woman would notice the change from gold to sand since the sand weighs much less than the same volume of gold.
- 45. Density = 9.0 g/cm^3

a)
$$V \pi r^2 h \ 3.14 \ (0.55 \text{ cm})^2 \ 2.85 \text{ cm} \ 2.7 \text{ cm}^3$$

$$d \frac{m}{V} \frac{24.3 \text{ g}_3}{2.7 \text{ cm}} 9.0 \text{ g/cm}^3$$

$$V \ 2.7 \text{ cm}$$

b) The metal is copper.

m = 1.7 x
$$10^{-24}$$
g, r = 1 x 10^{-13} cm
a) V= $\frac{4}{3}$ x 3.14 x $(1x10^{-13})^3$ = 4 x 10^{-39} cm³

$$d = \frac{m}{V} = \frac{1.7 \times 10^{-24}}{4 \times 10^{-39}} = 4 \times 10^{14} \text{ g/cm}^{3}$$

$$r = 1 \times 10^{-4} \text{m} \times 100 \text{ cm/1 m} = 1 \times 10^{-2} \text{ cm}$$

$$\frac{4}{3}$$

Density from part a):
$$d = 4 \times 10^{14} \text{g/cm}^3$$

Mass = d x V = 4 x
$$10^{-6}$$
 cm³ x $\frac{4x10^{14} \text{ g}}{1\text{ cm}^3}$ = 1.6 x 10^9 g x $\frac{1 \text{ kg}}{1000 \text{ g}}$ = 1.6 x 10^6 kg

SOLUTIONS TO POINTS TO PONDER:

47. - 51. Answers will vary.


SOLUTIONS TO FEATURE PROBLEMS:

52. The most precise scale (c) measures 5.4259 g. The least precise scale (a) measures 5.42 g. The uncertainty of scale a is \pm 0.01 g

The uncertainty of scale b is ± 0.001 g

The uncertainty of scale c is ± 0.0001 g.

53. penny -1.8 cm nickel -2.0 cm dime -1.6 cm quarter -2.3 cm half-dollar -2.9cm dollar -3.8 cm



The general trend is that as the value of the coin increases, the diameter of the coin also increases. The dime is the only coin that does not fit the general trend; it is too small.

19