

Solution Manual for College Algebra 8th Edition Aufmann Nation ISBN 1285434773 9781285434773


Full link download:

<https://testbankpack.com/p/solution-manual-for-college-algebra-8th-edition-aufmann-nation-isbn-1285434773-9781285434773/>

Chapter 2 Functions and Graphs

Section 2.1 Exercises

Plot the points:



Plot the points:

a. Find the decrease: The average debt decreased between 2006 and 2007, and 2008 and 2009.

b. Find the average debt in 2011:

Increase between 2009 to 2010: 22.0 20.1 1.9

Then the increase from 2010 to 2011:

22.0 1.9 23.9, or \$23,900.


a. When the cost of a game is \$22, 60 million games

can be sold.

b. The projected numbers of sales decreases as the price of this game increases.

c. Create a table and scatter diagram:

$R = p \cdot$	
$N 8$	$8 \cdot 80 = 640$
15	$15 \cdot 70 = 1050$
22	$22 \cdot 60 = 1320$
27	$27 \cdot 50 = 1350$
31	$31 \cdot 40 = 1240$
34	$34 \cdot 30 = 1020$
36	$36 \cdot 20 = 720$
37	$37 \cdot 10 = 370$



The revenue increases to a certain point and then decreases as the price of the game increases.

Determine whether the ordered pair is a solution

$$2x + 5y = 16$$

?

$$2(-2) + 5(4) = 16$$

?

$$-4 + 20 = 16$$

$$16 = 16 \text{ True}$$

$(-2, 4)$ is a solution.

Determine whether the ordered pair is a solution

$$2x^2 - 3y = 4$$

?

$$2(1)^2 - 3(-1) = 4$$

?

$$2 + 3 = 4$$

$$5 = 4 \text{ False}$$

$(1, -1)$ is not a solution.

Determine whether the ordered pair is a solution

$$= 3x^2 - 4x + 2$$

?

3)

$$17 = 3(-)^2 - 4(-3) + 2$$

?

$$17 = 27 + 12 + 2$$

$$17 = 41 \text{ False}$$

$(-3, 17)$ is not a solution.

Determine whether the ordered pair is a solution

$$x^2 + y^2 = 169$$

?

$$(-2)^2 + (12)^2 = 169$$

?

$$4 + 144 = 169$$

$$148 = 169 \text{ False}$$

$(-2, 12)$ is not a solution.

Find the distance: $(6, 4), (-8, 11)$

Find the distance: $(-5, 8), (-10, 14)$

$$= \sqrt{(10 - (-5))^2 + (14 - 8)^2}$$

$$= \sqrt{(-5)^2 + (6)^2}$$

$$= \sqrt{25 + 36}$$

$$= \sqrt{61}$$

Find the distance: $(-4, -20), (-10, 15)$

$$= \sqrt{(-10 - (-4))^2 + (15 - (-20))^2}$$

$$= \sqrt{(-6)^2 + (35)^2}$$

$$= \sqrt{36 + 1225}$$

$$= \sqrt{1261}$$

Find the distance: $(40, 32), (36, 20)$

$$= \sqrt{(36 - 40)^2 + (20 - 32)^2}$$

$$= \sqrt{(-4)^2 + (-12)^2}$$

$$= \sqrt{16 + 144}$$

$$= \sqrt{160}$$

$$= 4\sqrt{10}$$

Find the distance: $(5, -8), (0, 0)$

$$= \sqrt{(0 - 5)^2 + (0 - (-8))^2}$$

$$= \sqrt{(-5)^2 + (8)^2}$$

$$= \sqrt{25 + 64}$$

$$= \sqrt{89}$$

Find the distance: $(0, 0), (5, 13)$

$$= \sqrt{(5 - 0)^2 + (13 - 0)^2}$$

$$= \sqrt{5^2 + 13^2}$$

$$= \sqrt{25 + 169}$$

$$= \sqrt{194}$$

15. Find the distance: $(3\sqrt{3}, \sqrt{8}), (\sqrt{2}, \sqrt{27})$

$$d = \sqrt{(3\sqrt{3} - \sqrt{2})^2 + (\sqrt{8} - \sqrt{27})^2}$$

$$= \sqrt{(3\sqrt{3} - \sqrt{2})^2 + (2\sqrt{2} - 3\sqrt{3})^2}$$

16. Find the distance: $(\sqrt{25}, \sqrt{20}), (6, 2\sqrt{5})$

$$d = \sqrt{(6 - \sqrt{25})^2 + (2\sqrt{5} - \sqrt{20})^2}$$

$$= \sqrt{(6 - 5)^2 + (2\sqrt{5} - 2\sqrt{5})^2}$$

$$= \sqrt{(6 - 5)^2 + 0^2}$$

$$= \sqrt{(6 - 5)^2} = 6 - 5 = 1$$

Note: for another form of the solution,

$$d = \sqrt{(6 - 5)^2}$$

$$= \sqrt{36 - 60 + 25} = 16\sqrt{-60 + 25}$$

Find the distance: $(a, b), (-a, -b)$

$$= \sqrt{(a - (-a))^2 + (b - (-b))^2}$$

$$= \sqrt{(2a)^2 + (2b)^2}$$

$$= \sqrt{4a^2 + 4b^2}$$

$$= \sqrt{4(a^2 + b^2)}$$

$$= 2\sqrt{a^2 + b^2}$$

Find the distance: $(a - b, b), (a, a + b)$

$$= \sqrt{(a - (a - b))^2 + (b - (a + b))^2}$$

$$= \sqrt{(a - a + b)^2 + (b - a - b)^2}$$

$$= \sqrt{b^2 + (-a)^2}$$

$$= \sqrt{a^2 + b^2}$$

Find the distance: $(x, 4x), (-2x, 3x)$

$$= \sqrt{(-2x - x)^2 + (3x - 4x)^2}$$

$$= \sqrt{(-3x)^2 + (-x)^2}$$

$$= \sqrt{9x^2 + x^2}$$

$$= \sqrt{10x^2}$$

$= -x \sqrt{10}$ (Note: since $x < 0$, $x^2 = -x$)

$$(3)^2 + (3 - 2)^2$$

Find the distance: $(x, 4x), (-2x, 3x)$

$$\sqrt{3 + (27 - 12\sqrt{4}) + 8}$$

$$\sqrt{3 + 27 - 12\sqrt{4} + 8}$$

$$\sqrt{38 - 12\sqrt{4}}$$

$$= \sqrt{(-2x - x)^2 + (3x - 4x)^2} \text{ with } x > 0$$

$$\sqrt{(-3x)^2 + (-x)^2}$$

$$\sqrt{9x^2 + x^2}$$

$$\sqrt{10x^2}$$

$$= x\sqrt{10} \text{ (since } x > 0, \sqrt{x^2} = x)$$

Find the midpoint: (1, -1), (5, 5)

$$M = \left(\frac{1+5}{2}, \frac{-1+5}{2} \right)$$

$$= \left(\frac{6}{2}, \frac{4}{2} \right)$$

$$= (3, 2)$$

Find the midpoint: (-5, -2), (6, 10)

$$M = \left(\frac{-5+6}{2}, \frac{-2+10}{2} \right)$$

$$= \left(\frac{1}{2}, \frac{8}{2} \right)$$

$$= \left(0.5, 4 \right)$$

Find the midpoint: (6, -3), (6, 11)

$$M = \left(\frac{6+6}{2}, \frac{-3+11}{2} \right)$$

$$= \left(\frac{12}{2}, \frac{8}{2} \right)$$

$$= (6, 4)$$

Find the midpoint: (4, 7), (-10, 7)

$$M = \left(\frac{4+(-10)}{2}, \frac{7+7}{2} \right)$$

$$= \left(\frac{-6}{2}, \frac{14}{2} \right)$$

$$= (-3, 7)$$

Find the midpoint: (1.75, 2.25), (-3.5, 5.57)

$$M = \left(\frac{1.75+(-3.5)}{2}, \frac{2.25+5.57}{2} \right)$$

$$= \left(\frac{-1.75}{2}, \frac{7.82}{2} \right)$$

$$= (-0.875, 3.91)$$

Find the midpoint: (-8.2, 10.1), (-2.4, -5.7)

$$M = \left(\frac{-8.2+(-2.4)}{2}, \frac{10.1+(-5.7)}{2} \right)$$

Find other endpoint: endpoint (5, 1), midpoint (9, 3)

$$M = \left(\frac{x+5}{2}, \frac{y+1}{2} \right) = (9, 3)$$

therefore $\frac{x+5}{2} = 9$ and $\frac{y+1}{2} = 3$

$$x+5 = 18 \quad y+1 = 6$$

$$x = 13 \quad y = 5$$

Thus (13, 5) is the other endpoint.

Find other endpoint: endpoint (4, -6),

midpoint (-2, 11)

$$M = \left(\frac{x+4}{2}, \frac{y+(-6)}{2} \right) = (-2, 11)$$

therefore $\frac{x+4}{2} = -2$ and $\frac{y+(-6)}{2} = 11$

$$x+4 = -4 \quad y-6 = 22$$

$$x = -8 \quad y = 28$$

Thus (8, 28) is the other endpoint.

Find other endpoint: endpoint (-3, -8),

midpoint (2, -7)

$$M = \left(\frac{x+(-3)}{2}, \frac{y+(-8)}{2} \right) = (2, -7)$$

$$\frac{x-3}{2} = 2 \quad \frac{y-8}{2} = -7$$

therefore $\frac{x-3}{2} = 2$ and $\frac{y-8}{2} = -7$

$$x-3 = 4 \quad y-8 = -14$$

$$x = 7 \quad y = -6$$

Thus (7, -6) is the other endpoint.

Find other endpoint: endpoint (5, -4), midpoint (0, 0)

$$M = \left(\frac{x+5}{2}, \frac{y+(-4)}{2} \right) = (0, 0)$$

therefore $\frac{x+5}{2} = 0$ and $\frac{y-4}{2} = 0$

$$x + 5 = 0 \qquad y - 4 = 0$$


$$x = -5 \qquad y = 4$$

$$\left(-\frac{10.6}{2}, \frac{4.4}{2} \right)$$


$$(-5.3, 2.2)$$

Thus (5, 4) is the other endpoint.


Graph the equation: $x - y = 4$




Graph the equation: $2x + y = -1$




Graph the equation: $y = x^2 - 3$




Graph the equation: $y = 0.25x^2$




Graph the equation: $y = x^2 + 1$




Graph the equation: $3x^2 + 2y = -4$




Graph the equation: $y = \frac{1}{2}(x-1)^2$




Graph the equation: $y = -2|x+3|$




Graph the equation: $y = 2(x+2)^2$




Graph the equation: $y = |x+3| - 2$




Graph the equation: $y = x^2 + 2x - 8$




Graph the equation: $y = x^2 - 2x - 8$



Graph the equation: $y = -x^2 + 2$



Graph the equation: $y = -x^2 - 1$



Find the x- and y-intercepts and graph: $2x + 5y = 12$

For the y-intercept, let $x = 0$ and solve for y .


$$2(0) + 5y = 12$$

$$y = \frac{12}{5}, \quad \text{y-intercept: } (0, \frac{12}{5})$$

For the x-intercept, let $y = 0$ and solve for x .

$$2x + 5(0) = 12$$

$$x = 6, \quad \text{x-intercept: } (6, 0)$$



Find the x- and y-intercepts and graph: $3x - 4y = 15$

For the y-intercept, let $x = 0$ and solve for y .


$$3(0) - 4y = 15$$

$$y = -\frac{15}{4}, \quad \text{y-intercept: } (0, -\frac{15}{4})$$

For the x-intercept, let $y = 0$ and solve for x .

$$3x - 4(0) = 15$$

$$x = 5, \quad \text{x-intercept: } (5, 0)$$



Find the x- and y-intercepts and graph: $x = -y^2 + 5$

For the y-intercept, let $x = 0$ and solve for y .


$$0 = -y^2 + 5$$

$$y = \sqrt{5}, \quad \text{y-intercepts: } (0, -\sqrt{5}), (0, \sqrt{5})$$

For the x-intercept, let $y = 0$ and solve for x .

$$0 = -(0)^2 + 5$$

$$x = 5, \quad \text{x-intercept: } (5, 0)$$



Find the x- and y-intercepts and graph: $x = y^2 - 6$

For the y-intercept, let $x = 0$ and solve for y .


$$0 = y^2 - 6$$

$$y = \sqrt{6}, \quad \text{y-intercepts: } (0, -\sqrt{6}), (0, \sqrt{6})$$

For the x-intercept, let $y = 0$ and solve for x .

$$0 = (0)^2 - 6$$

$$x = -6, \quad \text{x-intercept: } (-6, 0)$$



Find the x- and y-intercepts and graph: $x = |y| - 4$

For the y-intercept, let $x = 0$ and solve for y .


$$0 = |y| - 4$$

$$= 4, \quad \text{y-intercepts: } (0, -4), (0, 4)$$

For the x-intercept, let $y = 0$ and solve for x .

$$x = 0 + 4$$

$$x = -4, \text{ x-intercept: } (-4, 0)$$



Find the x- and y-intercepts and graph: $x = y^3 - 2$

For the y-intercept, let $x = 0$ and solve for y .


$$0 = y^3 - 2$$

$$y = \sqrt[3]{2}, \text{ y-intercept: } (0, \sqrt[3]{2})$$

For the x-intercept, let $y = 0$ and solve for x .

$$= (0)^3 - 2$$

$$x = -2, \text{ x-intercept: } (-2, 0)$$



Find the x- and y-intercepts and graph: $x^2 + y^2 = 4$

For the y-intercept, let $x = 0$ and solve for y .


$$(0)^2 + y^2 = 4$$

$$= 2, \text{ y-intercepts: } (0, -2), (0, 2)$$

For the x-intercept, let $y = 0$ and solve for x .

$$x^2 + (0)^2 = 4$$

$$x = \pm 2, \text{ x-intercepts: } (-2, 0), (2, 0)$$




Find the x- and y-intercepts and graph: $x^2 = y^2$

For the y-intercept, let $x = 0$ and solve for y .

For the x-intercept, let $y = 0$ and solve for x .

Intercept: $(0, 0)$



Find center and radius: $x^2 + y^2 = 36$

center $(0, 0)$, radius 6

Find center and radius: $x^2 + y^2 = 49$

center $(0, 0)$, radius 7

Find center and radius: $(x - 1)^2 + (y - 3)^2 = 49$

center $(1, 3)$, radius 7

Find center and radius: $(x - 2)^2 + (y - 4)^2 = 25$

center $(2, 4)$, radius 5

Find center and radius: $(x + 2)^2 + (y + 5)^2 = 25$

center $(-2, -5)$, radius 5

Find center and radius: $(x + 3)^2 + (y + 5)^2 = 121$

center $(-3, -5)$, radius 11

Find center and radius: $(x - 8)^2 + y^2 =$

$\frac{1}{4}$ center $(8, 0)$, radius $\frac{1}{2}$

Find center and radius: $x^2 + (y - 12)^2 = 1$

center $(0, 12)$, radius 1

Find circle equation: center $(4, 1)$, radius 2

$$(x - 4)^2 + (y - 1)^2 = 2^2$$

$$(x - 4)^2 + (y - 1)^2 = 4$$

Find circle equation: center $(5, -3)$, radius 4

$$(x - 5)^2 + (y + 3)^2 = 4^2$$

$$(x - 5)^2 + (y + 3)^2 = 16$$

63. Find circle equation: center $(\frac{1}{2}, \frac{1}{4})$, radius $\sqrt{5}$

$$(x - \frac{1}{2})^2 + (y - \frac{1}{4})^2 = (\sqrt{5})^2$$

$$(x - \frac{1}{2})^2 + (y - \frac{1}{4})^2 = 5$$

64. Find circle equation: center $(0, \frac{2}{3})$, radius $\sqrt{11}$

$$(x - 0)^2 + (y - \frac{2}{3})^2 = (\sqrt{11})^2$$

$$(x - 0)^2 + (y - \frac{2}{3})^2 = 11$$

65. Find circle equation: center $(0, 0)$, through $(-3, 4)$

$$(x - 0)^2 + (y - 0)^2 = r^2$$

$$(-3 - 0)^2 + (4 - 0)^2 = r^2$$

$$(-3)^2 + 4^2 = r^2$$

$$9 + 16 = r^2$$

$$25 = 5^2 = r^2$$

$$(x - 0)^2 + (y - 0)^2 = 25$$

Find circle equation: center $(0, 0)$, through $(5, 12)$

$$(x - 0)^2 + (y - 0)^2 = r^2$$

$$(5 - 0)^2 + (12 - 0)^2 = r^2$$

$$5^2 + 12^2 = r^2$$

$$25 + 144 = r^2$$

$$169 = 13^2 = r^2$$

$$(x - 0)^2 + (y - 0)^2 = 169$$

Find circle equation: center $(1, 3)$, through $(4, -1)$

$$(x + 2)^2 + (y - 5)^2 = r^2$$

$$(x - 1)^2 + (y - 3)^2 = r^2$$

$$(4 - 1)^2 + (-1 - 3)^2 = r^2$$

$$3^2 + (-4)^2 = r^2$$

$$9 + 16 = r^2$$

$$25 = 5^2 = r^2$$

Find circle equation: center $(-2, 5)$, through $(1, 7)$

$$(1 + 2)^2 + (7 - 5)^2 = r^2$$

$$3^2 + 2^2 = r^2$$

$$9 + 4 = r^2$$

$$13 = (\sqrt{13})^2 = r^2$$

$$(x + 2)^2 + (y - 5)^2 = 13$$

Find circle equation: center $(-2, 5)$, diameter 10

diameter 10 means the radius is 5 $r^2 = 25$.

$$(x + 2)^2 + (y - 5)^2 = 25$$

Find circle equation: center $(0, -1)$, diameter 8

diameter 8 means the radius is 4 $r^2 = 16$.

$$(x - 0)^2 + (y + 1)^2 = 16$$

Find circle equation: endpoints $(2, 3)$ and $(-4, 11)$

$$= \sqrt{(4 - 2)^2 + (11 - 3)^2}$$

$$= \sqrt{36 + 64} = \sqrt{100}$$

$$= 10$$

Since the diameter is 10, the radius is 5.

The center is the midpoint of the line segment from $(2, 3)$ to $(-4, 11)$.

$$\frac{2 + (-4)}{2}, \frac{3 + 11}{2} = (-1, 7) \text{ center}$$

$$(x + 1)^2 + (y - 7)^2 = 25$$

Find circle equation: endpoints $(7, -2)$ and $(-3, 5)$

$$d = \sqrt{(-3 - 7)^2 + (5 - (-2))^2} = \sqrt{100 + 49} = \sqrt{149}$$

Since the diameter is $\sqrt{149}$, the radius is $\frac{\sqrt{149}}{2}$.

$$\text{Center is } \frac{7 + (-3)}{2}, \frac{(-2) + 5}{2} = \frac{4}{2}, \frac{3}{2}$$

$$= (2, \frac{3}{2})$$

$$(x - 2)^2 + (y - \frac{3}{2})^2 = (\frac{\sqrt{149}}{2})^2$$

$$(x - 1)^2 + (y - 3)^2 = 25$$

$$\frac{3}{2} = \frac{149}{2}$$

$$(x - 2)^2 + (y - 2)^2 = 4$$

Find circle equation: endpoints (5,-3) and (-1,-5)

$$d = \sqrt{(5 - (-1))^2 + (-3 - (-5))^2} = \sqrt{4 + 3\sqrt{40}} = \sqrt{40}$$

Since the diameter is $\sqrt{40}$, the radius is $\frac{\sqrt{40}}{2} = \sqrt{10}$.

$$\text{Center is } \left(\frac{5 + (-1)}{2}, \frac{(-3) + (-5)}{2} \right) = (2, -4)$$

$$(x - 2)^2 + (y + 4)^2 = (\sqrt{10})^2$$

$$(x - 2)^2 + (y + 4)^2 = 10$$

Find circle equation: endpoints (4,-6) and (0,-2)

$$d = \sqrt{(4 - 0)^2 + (-6 - (-2))^2} = \sqrt{16 + 16} = \sqrt{32}$$

Since the diameter is $\sqrt{32}$, the radius is $\frac{\sqrt{32}}{2} = \sqrt{2}$.

$$\text{Center is } \left(\frac{4 + 0}{2}, \frac{(-6) + (-2)}{2} \right) = (2, -4)$$

$$(x - 2)^2 + (y + 4)^2 = (\sqrt{2})^2$$

$$(x - 2)^2 + (y + 4)^2 = 2$$

Find circle equation: center (7, 11), tangent to x-axis

Since it is tangent to the x-axis, its radius is 11.

$$(x - 7)^2 + (y - 11)^2 = 11^2$$

Find circle equation: center (-2, 3), tangent to y-axis

Since it is tangent to the y-axis, its radius is 2.

$$(x + 2)^2 + (y - 3)^2 = 2^2$$

Find center and radius: $x^2 + y^2 - 6x + 5 = 0$

$$x^2 - 6x + y^2 = -5$$

$$x^2 - 6x + 9 + y^2 = -5 + 9$$

$$(x - 3)^2 + y^2 = 2^2$$

center (3, 0), radius 2

Find center and radius: $x^2 + y^2 - 14x + 8y + 53 = 0$

$$x^2 - 14x + y^2 + 8y = -53$$

$$x^2 - 14x + 49 + y^2 + 8y + 16 = -53 + 49 + 16$$

$$(x - 7)^2 + (y + 4)^2 = 12$$

center (7, -4), radius $\sqrt{12} = 2\sqrt{3}$

Find center and radius: $x^2 + y^2 - 10x + 2y + 18 = 0$

$$x^2 - 10x + y^2 + 2y = -18$$

$$x^2 - 10x + 25 + y^2 + 2y + 1 = -18 + 25 + 1$$

$$(x - 5)^2 + (y + 1)^2 = 8$$

center (5, -1), radius $\sqrt{8} = 2\sqrt{2}$

Find center and radius: $x^2 + y^2 - x + 3y - \frac{15}{4} = 0$

$$x^2 - x + y^2 + 3y = \frac{15}{4}$$

$$x^2 - x + \frac{1}{4} + y^2 + 3y + \frac{9}{4} = \frac{15}{4} + \frac{1}{4} + \frac{9}{4}$$

$$\left(x - \frac{1}{2}\right)^2 + \left(y + \frac{3}{2}\right)^2 = \frac{15}{4} + \frac{1}{4} + \frac{9}{4}$$

$$\left(x - \frac{1}{2}\right)^2 + \left(y + \frac{3}{2}\right)^2 = \frac{25}{4}$$

Find center and radius: $x^2 + y^2 + 3x - 5y + \frac{25}{4} = 0$

$$x^2 + 3x + y^2 - 5y = -\frac{25}{4}$$

$$x^2 + 3x + \frac{9}{4} + y^2 - 5y + \frac{25}{4} = -\frac{25}{4} + \frac{9}{4} + \frac{25}{4}$$

$$\left(x + \frac{3}{2}\right)^2 + \left(y - \frac{5}{2}\right)^2 = \frac{25}{4} + \frac{9}{4} + \frac{25}{4}$$

$$\left(x + \frac{3}{2}\right)^2 + \left(y - \frac{5}{2}\right)^2 = \frac{79}{4}$$

center $\left(-\frac{3}{2}, \frac{5}{2}\right)$, radius $\frac{\sqrt{79}}{2}$

Find center and radius: $x^2 + y^2 - 6x - 4y + 12 = 0$

Find center and radius: $x^2 + y^2 + 3x - 6y + 2 = 0$

$$x^2 - 6x + 9 + y^2 - 4y + 4 = -12 + 9 + 4$$

$$x^2 - 6x + 9 + y^2 - 4y + 4 = -12 + 9 + 4$$

$$(x - 3)^2 + (y - 2)^2 = 1^2$$

center (3, 2), radius 1

$$x^2 + 3x + \frac{9}{4} + y^2 - 6y + 9 = -2 + \frac{9}{4} + 9$$

$$(x + \frac{3}{2})^2 + (y - 3)^2 = \frac{37}{4}$$

center $(-\frac{3}{2}, 3)$ radius $\frac{\sqrt{37}}{2}$

Find center and radius: $x^2 + y^2 - 5x - y - 4 = 0$

$$x^2 - 5x + y^2 - y = 4$$

$$x^2 - 5x + \frac{25}{4} + y^2 - y + \frac{1}{4} = 4 + \frac{25}{4} + \frac{1}{4}$$

$$\left(x - \frac{5}{2}\right)^2 + \left(y - \frac{1}{2}\right)^2 = \frac{49}{4}$$

center $\left(\frac{5}{2}, \frac{1}{2}\right)$, radius $\frac{7}{2}$

Find the points:

$$\sqrt{(4-x)^2 + (6-0)^2} = 10$$

$$\sqrt{(4-x)^2 + (6-0)^2} = 10^2$$

$$16 - 8x + x^2 + 36 = 100$$

$$x^2 - 8x - 48 = 0$$

$$(x-12)(x+4) = 0$$

$$x = 12 \text{ or } x = -4$$

The points are (12, 0), (-4, 0).

Find the points:

$$\sqrt{(5-0)^2 + (y-(-3))^2} = 12$$

$$\sqrt{(5)^2 + (y+3)^2} = 12^2$$

$$25 + y^2 + 6y + 9 = 144$$

$$y^2 + 6y - 110 = 0$$

$$y = \frac{-6 \pm \sqrt{6^2 - 4(1)(-110)}}{2(1)}$$

$$y = \frac{-6 \pm \sqrt{36 + 440}}{2}$$

$$y = \frac{-6 \pm \sqrt{476}}{2}$$


$$y = \frac{-6 \pm 2\sqrt{119}}{2}$$

$$y = -3 \pm \sqrt{119}$$

The points are $(0, -3 + \sqrt{119})$, $(0, -3 - \sqrt{119})$.

Find the x- and y-intercepts and graph: $|x + y| = 4$

Intercepts: (0, 4), (-4, 0)



Find the x- and y-intercepts and graph: $|x - 4y| = 8$

For the y-intercept, let $x = 0$ and solve for y .

$$|0 - 4y| = 8$$


$$4y = 8$$

$$y = 2, \text{ y-intercepts: } (0, 2), (0, -2)$$

For the x-intercept, let $y = 0$ and solve for x .

$$|x - 4(0)| = 8$$

$$x = 8, \text{ x-intercepts: } (-8, 0), (8, 0)$$



Find the formula:

$$\sqrt{(3-x)^2 + (4-y)^2} = 5$$

$$(3-x)^2 + (4-y)^2 = 5^2$$

$$9 - 6x + x^2 + 16 - 8y + y^2 = 25$$

$$x^2 - 6x + y^2 - 8y = 0$$

Find the formula:

$$\sqrt{(-5-x)^2 + (12-y)^2} = 13$$

$$(-5-x)^2 + (12-y)^2 = 13^2$$

$$25 + 10x + x^2 + 144 - 24y + y^2 = 169$$

$$x^2 + 10x + y^2 - 24y = 0$$

Prepare for Section 2.2

P1. $x^2 + 3x - 4$

$$(-3)^2 + 3(-3) - 4 = 9 - 9 - 4 = -4$$

P2. $D = \{-3, -2, -1, 0, 2\}$

R

$=$

$\{1$

,

$2,$

$4,$

$5\}$

P3. $d = \sqrt{(3 - (-4))^2 + (-2 - 1)^2} = 49\sqrt{9} = 58 \sqrt{\quad}$

P4. $2x - 6 \geq 0$
 $2x \geq 6$
 $x \geq 3$

P5. $x^2 - x - 6 = 0$
 $(x + 2)(x - 3) = 0$
 $x + 2 = 0 \quad x - 3 = 0$
 $x = -2 \quad x = 3$
 $-2, 3$

P6. $a = 3x + 4, \quad a = 6x - 5$
 $3x + 4 = 6x - 5$
 $= 3x$
 $= x$
 $= 3(3) + 4 = 13$

Section 2.2 Exercises

Write the domain and range. State whether a relation.

Domain: $\{-4, 2, 5, 7\}$; range: $\{1, 3, 11\}$

Yes. The set of ordered pairs defines y as a function of x since each x is paired with exactly one y .

Write the domain and range. State whether a relation.

Domain: $\{3, 4, 5\}$; range: $\{-2, 7, 8, 10\}$

No. The set of ordered pair does not define y as a function of x since 5 is paired with 10 and 8.

Write the domain and range. State whether a relation.

Domain: $\{4, 5, 6\}$; range: $\{-3, 1, 4, 5\}$

No. The set of ordered pair does not define y as a function of x since 4 is paired with 4 and 5.

Write the domain and range. State whether a relation.

Domain: $\{1, 2, 3\}$; range $\{0\}$

Yes. The set of ordered pairs defines y as a function of x since each x is paired with exactly one y .

Determine if the value is in the domain.

$(0) = 0 \frac{3(0)}{+4} = 0$

Yes, 0 is in the domain of the function.

Determine if the value is in the domain.

$g(-1) = 1 - (-1)^2 = 0$

Yes, -1 is in the domain of the function.

Determine if the value is in the domain.

$F(0) = \frac{-1}{-1+1} = \frac{-1}{0}$ undefined

No, -1 is not in the domain of the function.

Determine if the value is in the domain.

$y(2) = \sqrt{2(2) - 8} = \sqrt{-4}$

No, 2 is not in the domain of the function.

Determine if the value is in the domain.

$g(-1) = \frac{5(-1)-1}{-1} = \frac{-6}{-1} = 6$

Yes, -1 is in the domain of the function.

Determine if the value is in the domain.

$F(-2) = \frac{1}{(-2)^3 + 8} = \frac{1}{0}$

No, 0 is not in the domain of the function.

Is y a function of x ?

$2x + 3y = 7$
 $3y = -2x + 7$
 $y = -\frac{2}{3}x + \frac{7}{3}$, y is a function of x .

Is y a function of x ?

$5x + y = 8$
 $y = -5x + 8$, y is a function of x .

Is y a function of x ?

$-x + y^2 = 2$
 $y^2 = x + 2$
 $y = \sqrt{\quad}$ $x + 2$, y is a not function of x .

Is y a function of x ?

$x - 2y = 2$
 $-2y = -x + 2$

$y = \frac{1}{2}x - 1$, y is a function of x .

Is y a function of x ?

$$x^2 + y^2 = 9$$

$$y^2 = 9 - x^2$$

$y = \sqrt{9 - x^2}$, y is not a function of x .

Is y a function of x ?

$$y = \sqrt{x}, y \text{ is a function of } x.$$

Is y a function of x ?

$$y = x + 5, y \text{ is a function of } x.$$

Is y a function of x ?

$$y = \sqrt{x^2 + 4}, y \text{ is a function of } x.$$

Determine if the value is a zero.

$$f(-2) = 3(-2) + 6 = 0$$

Yes, -2 is a zero.

Determine if the value is a zero.

$$f(0) = 2(0)^3 - 4(0)^2 + 5(0) = 0$$

Yes, 0 is a zero.

Determine if the value is a zero.

$$f\left(-\frac{1}{3}\right) = 3\left(-\frac{1}{3}\right)^2 + 2\left(-\frac{1}{3}\right) - \frac{1}{3}$$

No, $-\frac{1}{3}$ is not a zero.

Determine if the value is a zero.

$$s(-1) = \frac{2(-1) + 6}{-1 + 1} = \frac{4}{0} \text{ undefined}$$

No, -1 is not a zero.

Determine if the value is a zero.

$$y(1) = 5(1)^2 - 2(1) - 2 = 1$$

No, 1 is not a zero.

Determine if the value is a zero.

$$g(-3) = \frac{3(-3) + 9}{(-3)^2 - 4} = \frac{0}{5} = 0$$

Yes, -3 is a zero.

25. Evaluate the function $f(x) = 3x - 1$,

$$\begin{aligned} \text{a. } f(2) &= 3(2) - 1 \\ &= 6 - 1 \\ &= 5 \end{aligned}$$

$$\begin{aligned} \text{b. } f(-1) &= 3(-1) - 1 \\ &= -3 - 1 \\ &= -4 \end{aligned}$$

$$\begin{aligned} \text{c. } f(0) &= 3(0) - 1 \\ &= 0 - 1 \\ &= -1 \end{aligned}$$

$$\text{d. } f(2) = 3(2) - 1 = 5$$

$$\begin{aligned} f(3) &= 3(3) - 1 \\ &= 9 - 1 \\ &= 8 \end{aligned}$$

$$\begin{aligned} f(k) &= 3(k) - 1 \\ &= 3k - 1 \end{aligned}$$

$$\begin{aligned} f(k + 2) &= 3(k + 2) - 1 \\ &= 3k + 6 - 1 \\ &= 3k + 5 \end{aligned}$$

Evaluate the function $g(x) = 2x^2 + 3$,

$$g(3) = 2(3)^2 + 3 = 18 + 3 = 21$$

$$g(-1) = 2(-1)^2 + 3 = 2 + 3 = 5$$

$$g(0) = 2(0)^2 + 3 = 0 + 3 = 3$$

$$\text{d. } g(2) = 2(2)^2 + 3 = 2(4) + 3 = 11$$

$$g(c) = 2(c)^2 + 3 = 2c^2 + 3$$

$$g(c + 5) = 2(c + 5)^2 + 3$$

$$\begin{aligned} &= 2c^2 + 20c + 50 + 3 \\ &= 2c^2 + 20c + 53 \end{aligned}$$

27. Evaluate the function $A(w) = \sqrt{w^2 + 5}$,

$$\text{a. } A(0) = \sqrt{(0)^2 + 5} = \sqrt{5}$$

$$\text{b. } A(2) = \sqrt{(2)^2 + 5} = \sqrt{9} = 3$$

c. $A(-2) = (-2)^2 + 5 = 9 + 5 = 14$

d. $A(4) = 4^2 + 5 = 16 + 5 = 21$

$$\sqrt{\quad} \quad \sqrt{\quad}$$

$$\sqrt{\quad} \quad \sqrt{\quad}$$

$$e. A(r+1) = \frac{\sqrt{(r+1)^2 + 5}}{\sqrt{r^2 + 2r + 1 + 5}} \cdot \sqrt{r^2 + 2r + 6}$$

$$A(-c) = (-c)\sqrt{c^2 + 5} = c^2 + 5$$

Evaluate the function $J(t) = 3t^2 - t$,

$$J(-4) = 3(-4)^2 - (-4) = 48 + 4 = 52$$

$$J(0) = 3(0)^2 - (0) = 0 - 0 = 0$$

$$c. J\left(\frac{1}{3}\right) = 3\left(\frac{1}{3}\right)^2 - \frac{1}{3} = \frac{1}{3} - \frac{1}{3} = 0$$

$$= 3 \cdot \frac{1}{9} - \frac{1}{3} = \frac{1}{3} - \frac{1}{3} = 0$$

$$d. J(-c) = 3(-c)^2 - (-c) = 3c^2 + c$$

$$e. J(x+1) = 3(x+1)^2 - (x+1) = 3x^2 + 6x + 3 - x - 1$$

$$= 3x^2 + 5x + 2$$

$$f. J(x+h) = 3(x+h)^2 - (x+h)$$

$$= 3x^2 + 6xh + 3h^2 - x - h$$

29. Evaluate the function $f(x) = \frac{1}{|x|}$,

$$a. f(2) = \frac{1}{|2|} = \frac{1}{2}$$

$$b. f(-2) = \frac{1}{|-2|} = \frac{1}{2}$$

$$c. f\left(\frac{1}{5}\right) = \frac{1}{\left|\frac{1}{5}\right|} = \frac{1}{\frac{1}{5}} = 5$$

$$f(2) + f(-2) = \frac{1}{2} + \frac{1}{2} = 1$$

30. Evaluate the function $T(x) = 5$,

$$a. T(-3) = 5$$

$$b. T(0) = 5$$

$$c. T(3) + T(1) = 5 + 5 = 10$$

$$T(x+h) = 5$$

$$T(3k+5) = 5$$

Evaluate the function $s(x) = \frac{x}{|x|}$,

$$s(4) = \frac{4}{|4|} = \frac{4}{4} = 1$$

$$s(5) = \frac{5}{|5|} = \frac{5}{5} = 1$$

$$s(-2) = \frac{-2}{|-2|} = \frac{-2}{2} = -1$$

$$s(-3) = \frac{-3}{|-3|} = \frac{-3}{3} = -1$$

e. Since $t > 0$, $t = |t|$.

$$s(t) = \frac{t}{|t|} = \frac{t}{t} = 1$$

Since $t < 0$, $t = -|t|$.

$$s(t) = \frac{t}{|t|} = \frac{-|t|}{|t|} = -1$$

Evaluate the function $r(x) = x + 4$,

$$a. r(0) = 0 + 4 = 4$$

$$r(-1) = -1 + 4 = 3$$

$$r(-3) = -3 + 4 = 1$$

$$\frac{1}{2} \left(\frac{1}{2} \right)$$

e. $f(c^2 + 4) = \frac{1}{c^2 + 4} = \frac{1}{c^2 + 4}$

1

f. $f(2 + h) = \frac{1}{2 + h}$

d. $\frac{f(2) - f(2+h)}{2 - (2+h)}$

$= \frac{\frac{1}{2} - \frac{1}{2+h}}{2 - (2+h)} = \frac{\frac{2+h}{2(2+h)} - \frac{2}{2(2+h)}}{-h} = \frac{\frac{2+h-2}{2(2+h)}}{-h} = \frac{\frac{h}{2(2+h)}}{-h} = -\frac{1}{2(2+h)}$

e. $r(0.1) = \frac{0.1}{0.1+4} = \frac{0.1}{4.1} = \frac{1}{41}$

f. $r(10,000) = \frac{10,000}{10,000+4} = \frac{10,000}{10,004} = \frac{2500}{2501}$

a. Since $x = -4 < 2$, use $P(x) = 3x + 1$.

$P(-4) = 3(-4) + 1 = -12 + 1 = -11$

Since $x = 5^3 \sqrt[3]{\text{use } P(x) = -x^2 + 11}$.

$P(\sqrt[3]{5}) = -(5)^{\frac{2}{3}} + 11 = -5 + 11 = 6$

Since $x = c < 2$, use $P(x) = 3x + 1$. $P(c$

$) = 3c + 1$

Since $k^3 \geq 1$, then $x = k + 1^3 \geq 2$,

so use $P(x) = -x^2 + 11$.

$$\begin{aligned} P(k+1) &= -(k+1)^2 + 11 = -(k^2 + 2k + 1) + 11 \\ &= -k^2 - 2k - 1 + 11 \\ &= -k^2 - 2k + 10 \end{aligned}$$

a. Since $t = 0$ and $0 \leq t \leq 5$, use $Q(t) = 4$.

$Q(0) = 4$

Since $t = e$ and $6 < e < 7$, then $5 < t \leq 8$, so

use $Q(t) = -t + 9$.

$Q(e) = -e + 9$

Since $t = n$ and $1 < n < 2$, then $0 \leq t \leq 5$, so

use $Q(t) = 4$

$Q(0) = 4$

Since $t = m^2 + 7$ and $1 < m \leq 2$,

then $1^2 < m^2 \leq 2^2$

$1^2 + 7 < m^2 + 7 \leq 2^2 + 7$

$1 + 7 < m^2 + 7 \leq 4 + 7$

$8 < m^2 + 7 \leq 11$ thus

$8 < t \leq 11$,

so use $Q(t) = \sqrt{t-7}$

$Q(m^2 + 7) = \sqrt{(m^2 + 7) - 7}$

For $f(x) = 3x - 4$, the domain is the set of all real numbers.

36. For $f(x) = -2x + 1$, the domain is the set of all real numbers.

37. For $f(x) = x^2 + 2$, the domain is the set of all real numbers.

38. For $f(x) = 3x^2 + 1$, the domain is the set of all real numbers.

$\frac{4}{|x|}$

For $f(x) = x + 2$, the domain is $\{x \in \mathbb{R}\}$.

For $f(x) = x - 5$, the domain is $\{x \in \mathbb{R}\}$.

\sqrt{x}

41. For $f(x) = \sqrt{x+7}$, the domain is $\{x \in \mathbb{R} \mid x \geq -7\}$.

For $f(x) = 4 - x$, the domain is $\{x \in \mathbb{R}\}$.

43. For $f(x) = \sqrt{4-x^2}$, the domain is $\{x \in \mathbb{R} \mid -2 \leq x \leq 2\}$.

For $f(x) = \sqrt{2-x^2}$, the domain is

$\{x \in \mathbb{R} \mid -\sqrt{2} \leq x \leq \sqrt{2}\}$.

$(-\infty, \frac{1}{x+4}) \cup (1, \infty)$


45. For $f(x) = \frac{1}{x+4}$, the domain is $\{x \in \mathbb{R} \mid x \neq -4\}$.

$(-\infty, \frac{1}{5-x}) \cup (5, \infty)$

46. For $f(x) = \sqrt{5-x}$, the domain is $\{x \in \mathbb{R} \mid x \leq 5\}$.

To graph $f(x) = 3x - 4$, plot points and draw a smooth graph.


x	-1	0	1	2
$y = f(x) = 3x - 4$	-7	-4	-1	2



$$= \sqrt{m^2} = |m| = m \text{ since } m > 0$$


48. To graph $f(x) = 2 - \frac{1}{2}x$, plot points and draw a smooth graph.

x	-4	-2	0	4
$y = f(x) = 2 - \frac{1}{2}x$	4	3	2	0




To graph $g(x) = x^2 - 1$, plot points and draw a smooth graph.

x	-2	-1	0	1	2
$y = g(x) = x^2 - 1$	3	0	-1	0	3




To graph $g(x) = 3 - x^2$, plot points and draw a smooth graph.

x	-3	-1	0	1	3
$y = g(x) = 3 - x^2$	-6	2	3	2	-6




51. To graph $f(x) = \sqrt{x+4}$, plot points and draw a smooth graph.

x	-4	-2	0	2	5
$y = f(x) = \sqrt{x+4}$	0	$\sqrt{2}$	2	$\sqrt{6}$	3




To graph $h(x) = \sqrt{5-x}$, plot points and draw a smooth graph.

x	-4	0	1	3	5
$y = h(x) = \sqrt{5-x}$	3	$\sqrt{5}$	2	$\sqrt{2}$	0




To graph $f(x) = |x - 2|$, plot points and draw a smooth graph.

x	-3	0	2	4	6
$y = f(x) = x - 2 $	5	2	0	2	4




54. To graph $h(x) = 3 - |x|$, plot points and draw a smooth graph.

x	-3	-1	0	1	3
$y = h(x) = 3 - x $	0	2	3	2	0




To graph $L(x) = \frac{1}{3}x$ for $-6 \leq x \leq 6$, plot points

x	-6	-4	-3	-1	0	4	6
$y = L(x) = \frac{1}{3}x$	-2	-2	-1	-1	0	1	2




To graph $L(x) = x + 2$ for $0 \leq x \leq 4$, plot points and draw a smooth graph.


x	0	1	2	3	4
$y = L(x) = x + 2$	2	3	4	5	6



To graph $N(x) = \text{int}(-x)$ for $-3 \leq x \leq 3$, plot points and draw a smooth graph.




To graph $N(x) = \text{int}(x) + x$ for $0 \leq x \leq 4$, plot points and draw a smooth graph.




59. Graph $f(x) = \begin{cases} 1-x & x < 2 \\ 2x & x \geq 2 \end{cases}$

Graph $y = 1 - x$ for $x < 2$ and graph $y = 2x$ for $x \geq 2$.




60. Graph $f(x) = \begin{cases} 2x & x \leq -1 \\ x & x > -1 \end{cases}$

Graph $y = 2x$ for $x \leq -1$ and graph $y = x$ for $x > -1$.




61. Graph $r(x) = \begin{cases} -x^2 + 4 & x < -1 \\ -x + 2 & -1 \leq x \leq 1 \\ 3x - 2 & x > 1 \end{cases}$

Graph $y = -x^2 + 4$ for $x < -1$, graph $y = -x + 2$ for $-1 \leq x \leq 1$, and graph $y = 3x - 2$ for $x > 1$.



62. Graph $A(x) = \begin{cases} |x| & x < 1 \\ x^2, & 1 \leq x < 3 \\ -x + 2, & x \geq 3 \end{cases}$

Graph $y = |x|$ for $x < 1$, graph $y = x^2$ for $1 \leq x < 3$, and graph $y = -x + 2$ for $x \geq 3$.



Find the value of a in the domain of $f(x) = 3x - 2$ for which $f(a) = 10$.

$$\begin{aligned} 3a - 2 &= 10 && \text{Replace } f(a) \text{ with } 3a - 2 \\ 3a &= 12 \\ a &= 4 \end{aligned}$$

Find the value of a in the domain of $f(x) = 2 - 5x$ for which $f(a) = 7$.

$$\begin{aligned} 2 - 5a &= 7 && \text{Replace } f(a) \text{ with } 2 - 5a \\ -5a &= 5 \\ a &= -1 \end{aligned}$$

Find the values of a in the domain of

$$\begin{aligned} f(x) &= x^2 + 2x - 2 \text{ for which } f(a) = 1. \\ a^2 + 2a - 2 &= 1 && \text{Replace } f(a) \text{ with } a^2 + 2a - 2 \\ a^2 + 2a - 3 &= 0 \end{aligned}$$

$$(a + 3)(a - 1) = 0$$

$$\begin{aligned} a + 3 &= 0 && a - 1 = 0 \\ a &= -3 && a = 1 \end{aligned}$$

Find the values of a in the domain of

$$\begin{aligned} f(x) &= x^2 - 5x - 16 \text{ for which } f(a) = -2. \\ a^2 - 5a - 16 &= -2 && \text{Replace } f(a) \text{ with } a^2 - 5a - 16 \\ a^2 - 5a - 14 &= 0 \\ (a + 2)(a - 7) &= 0 \end{aligned}$$

$$\begin{aligned} a + 2 &= 0 && a - 7 = 0 \\ a &= -2 && a = 7 \end{aligned}$$

Find the values of a in the domain of $f(x) = |x|$ for

which $f(a) = 4$.

$$\begin{aligned} |a| &= 4 && \text{Replace } f(a) \text{ with } |a| \\ a &= -4 && a = 4 \end{aligned}$$

68. Find the values of a in the domain of $f(x) = |x + 2|$

for which $f(a) = 6$.

$$\begin{aligned} |a + 2| &= 6 && \text{Replace } f(a) \text{ with } |a + 2| \\ a + 2 &= -6 && a + 2 = 6 \\ a &= -8 && a = 4 \end{aligned}$$

69. Find the values of a in the domain of $f(x) = x^2 + 2$ for which $f(a) = 1$.

$$\begin{aligned} a^2 + 2 &= 1 && \text{Replace } f(a) \text{ with } a^2 + 2 \\ a^2 &= -1 \end{aligned}$$

There are no real values of a .

Find the values of a in the domain of $f(x) = |x| - 2$

for which $f(a) = -3$.

$$\begin{aligned} |a| - 2 &= -3 && \text{Replace } f(a) \text{ with } |a| - 2 \\ |a| &= -1 \end{aligned}$$

There are no real values of a .

71. Find the zeros of f for $f(x) = 3x - 6$.

$$\begin{aligned} f(x) &= 0 \\ 3x - 6 &= 0 \\ 3x &= 6 && x = \\ & && 2 \end{aligned}$$

72. Find the zeros of f for $f(x) = 6 + 2x$.

$$\begin{aligned} f(x) &= 0 \\ 6 + 2x &= 0 \\ 2x &= -6 \\ x &= -3 \end{aligned}$$

73. Find the zeros of f for $f(x) = 5x + 2$.

$$\begin{aligned} f(x) &= 0 \\ 5x + 2 &= 0 \\ 5x &= -2 \\ x &= -\frac{2}{5} \end{aligned}$$

74. Find the zeros of f for $f(x) = 8 - 6x$.

$$\begin{aligned} f(x) &= 0 \\ 8 - 6x &= 0 \\ -6x &= -8 \\ x &= \frac{4}{3} \end{aligned}$$

Find the zeros of f for $f(x) = x^2 - 4$.

$$\begin{aligned} f(x) &= 0 \\ x^2 - 4 &= 0 \quad (x + 2)(x - 2) = 0 \\ x + 2 &= 0 & x - 2 &= 0 \\ x &= -2 & x &= 2 \end{aligned}$$

76. Find the zeros of f for $f(x) = x^2 + 4x - 21$.

$$\begin{aligned} f(x) &= 0 \\ x^2 + 4x - 21 &= 0 \\ (x + 7)(x - 3) &= 0 \\ x + 7 &= 0 & x - 3 &= 0 \\ x &= -7 & x &= 3 \end{aligned}$$

Find the zeros of f for $f(x) = x^2 - 5x - 24$.

$$\begin{aligned} f(x) &= 0 \\ x^2 - 5x - 24 &= 0 \\ (x + 3)(x - 8) &= 0 \\ x + 3 &= 0 & x - 8 &= 0 \\ x &= -3 & x &= 8 \end{aligned}$$

78. Find the zeros of f for $f(x) = 2x^2 + 3x - 5$.

$$\begin{aligned} f(x) &= 0 \\ 2x^2 + 3x - 5 &= 0 \\ (2x + 5)(x - 1) &= 0 \\ 2x + 5 &= 0 & x - 1 &= 0 \\ x &= -\frac{5}{2} & x &= 1 \end{aligned}$$

Determine which graphs are functions.

Yes; every vertical line intersects the graph in one point.

Yes; every vertical line intersects the graph in one point.

No; some vertical lines intersect the graph at more than one point.

Yes; every vertical line intersects the graph in one point.

a. Yes; every vertical line intersects the graph in one point.

No; some vertical lines intersect the graph at more than one point.

No; a vertical line intersects the graph at more than one point.

Yes; every vertical line intersects the graph in one point.

Determine where the graph is increasing, constant, or decreasing. Decreasing on $(-, 0)$; increasing on $[0,)$

Determine where the graph is increasing, constant, or decreasing. Decreasing on $(-\infty, \infty)$

Determine where the graph is increasing, constant, or decreasing. Increasing on $(-\infty, \infty)$

Determine where the graph is increasing, constant, or decreasing. Increasing on $(-\infty, 2)$; decreasing on $[2, \infty)$

Determine where the graph is increasing, constant, or decreasing. Decreasing on $(-\infty, -3)$; increasing on $[-3, 0]$; decreasing on $[0, 3]$; increasing on $[3, \infty)$

Determine where the graph is increasing, constant, or decreasing. Increasing on $(-\infty, \infty)$

Determine where the graph is increasing, constant, or decreasing. Constant on $(-\infty, 0]$; increasing on $[0, \infty)$

Determine where the graph is increasing, constant, or decreasing. Constant on $(-\infty, \infty)$

Determine where the graph is increasing, constant, or decreasing. Decreasing on $(-\infty, 0]$;

constant on $[0, 1]$; increasing on $[1, \infty)$

Determine where the graph is increasing, constant, or decreasing. Constant on $(-\infty, 0]$;

decreasing on $[0, 3]$; constant on $[3, \infty)$

Determine which functions from 77-81 are one-to-one.

g and F are one-to-one since every horizontal line intersects the graph at one point.

f , V , and p are not one-to-one since some horizontal lines intersect the graph at more than one point.

Determine which functions from 82-86 are one-to-one.

s is one-to-one since every horizontal line intersects the graph at one point.

t , m , r and k are not one-to-one since some horizontal lines intersect the graph at more than one point.

a. $C(2.8) = 0.90 - 0.20\text{int}(1 - 2.8)$


$$0.90 - 0.20\text{int}(-1.8)$$

$$0.90 - 0.20(-2)$$

$$0.90 + 0.4$$

$$\text{\$}1.30$$

b. Graph $C(w)$.



a. Domain: $[0, \infty)$

b. $T(50,020) = 0.25(50,020 - 35,350) + 4867.50$

$$0.25(14,670) + 4867.50$$

$$3667.50 + 4867.50$$

$$\text{\$}8535$$

$$T(123,500) = 0.28(123,500 - 85,650) + 17,442.50$$

$$0.28(37,850) + 17,442.50$$

$$10,598 + 17,442.50$$

$$\text{\$}28,040.50$$

a. Write the width.

$$2l + 2w = 50$$

$$2w = 50 - 2l$$

$$= 25 - l$$

b. Write the area.

$$A = lw$$

$$= l(25 - l)$$

$$A = 25l - l^2$$

a. Write the length.

$$= \frac{12l}{d + l}$$

$$4(d + l) = 12l$$

$$4d + 4l = 12l$$

$$4d = 8l$$

$$\frac{1}{2}d = l$$

$$l(d) = \frac{1}{2}d$$

Find the domain. Domain: $[0, \infty)$

$$\frac{1}{2}$$

Find the length. $l(8) = \frac{1}{2}(8) = 4$ ft

Write the function.

$$v(t) = 80,000 - 6500t, \quad 0 \leq t \leq 10$$

Write the function.

$$v(t) = 44,000 - 4200t, \quad 0 \leq t \leq 8$$

a. Write the total cost function.

$$C(x) = 5(400) + 22.80x$$

$$2000 + 22.80x$$

Write the revenue function. $R(x) = 37.00x$

Write the profit function.

$$P(x) = 37.00x - C(x)$$

$$37.00x - [2000 + 22.80x]$$

$$37.00x - 2000 - 22.80x$$

$$14.20x - 2000$$

Note x is a natural number.

a. Write the volume function.

$$V = lwh$$

$$V = (30 - 2x)(30 - 2x)(x)$$

$$= (900 - 120x + 4x^2)(x)$$

$$= 900x - 120x^2 + 4x^3$$

State the domain.

$= lwh$ the domain of V is dependent on the domains of l , w , and h . Length, width and height must

be positive values $30 - 2x > 0$ and $x > 0$.

$$-2x > -30$$

$$x < 15$$

Thus, the domain of V is $\{x \mid 0 < x < 15\}$.

101. Write the function.

$$\frac{15}{3} = \frac{15-h}{r}$$

$$= \frac{15-h}{r}$$

$$5r = 15 - h$$

$$= 15 - 5r$$

$$h(r) = 15 - 5r$$

a. Write the function.

$$- =$$

$$h = 4$$

$$r = \frac{4}{2}$$

$$r = \frac{1}{2}h$$

Write the function. $V =$

$$\frac{1}{3} \pi r^2 h$$

$$= \frac{1}{3} \pi \left(\frac{1}{2} h \right)^2 h = \frac{1}{3} \pi$$

$$\left(\frac{1}{4} h^2 \right) h V = 12 \frac{1}{3} \pi h^3$$

103. Write the function.

$$= \sqrt{(3t)^2 + (50)^2}$$

$$d = \sqrt{9t^2 + 2500} \text{ meters, } 0 \leq t \leq 60$$

104. Write the function.

$$t = \frac{d}{r}$$

105. Write the function.

$$= \sqrt{(45 - 8t)^2 + (6t)^2} \text{ miles}$$

where t is the number of hours after 12:00 noon

106. Write the function.

$$= \sqrt{(60 - 7t)^2 + (10t)^2} \text{ miles}$$

where t is the number of hours after 12:00 noon

a. Write the function.

Left side triangle

$$c^2 = 20^2 + (40 - x)^2$$

$$c = \sqrt{400 + (40 - x)^2}$$

Right side triangle

$$c^2 = 30^2 + x^2$$

$$c = \sqrt{900 + x^2}$$

$$\text{Total length} = \sqrt{900 + x^2} + \sqrt{400 + (40 - x)^2}$$

Complete the table.

x	0	10	20	30	40
Total					
Length	74.72	67.68	64.34	64.79	70

Find the domain. Domain: $[0, 40]$.

108. Complete the table.

p	40	50	60	75	r
$f(p)$	4900	4300	3800	3200	$\frac{2}{2800}$

Answers accurate to nearest 100 feet.

109. Complete the table.

x	5	10	12.5	15	20
$Y(x)$	275	375	385	390	394

Answers accurate to the nearest apple.

110. Complete the table.

x	100	200	500	750	1000
$C(x)$	57,121	59,927	65,692	69,348	72,507

Answers accurate to the nearest dollar.

111. Find c .

$$(c) = c^2 - c - 5 = 1$$

$$c^2 - c - 6 = 0$$

$$(c - 3)(c + 2) = 0$$

$$c - 3 = 0 \quad \text{or} \quad c + 2 = 0$$

$$c = 3 \qquad c = -2$$
$$t = \frac{\sqrt{1+x^2}}{2} + \frac{3-x}{8} \text{ hours}$$

Chapter 2 Functions and Graphs

112. Find c .

$$\begin{aligned}
 g(c) &= -2c^2 + 4c - 1 = -4 \\
 &= -2c^2 + 4c + 3 = 0 \\
 c &= \frac{-4 \pm \sqrt{(-2)(3)}}{2(-2)} \\
 c &= \frac{-4 \pm \sqrt{24}}{-4} = \frac{-4 \pm 2\sqrt{6}}{-4} \\
 c &= \frac{-4 \pm 2\sqrt{6}}{-4} \\
 c &= \frac{2 \pm \sqrt{6}}{2}
 \end{aligned}$$

113. Determine if 1 is in the range.

1 is not in the range of $f(x)$, since


$$1 = \frac{x-1}{x+1} \text{ only if } x+1 = x-1 \text{ or } 1 = -1.$$

114. Determine if 0 is in the range.

0 is not in the range of $g(x)$, since

$$0 = \frac{1}{-3} \text{ only if } (x-3)(0) = 1 \text{ or } 0 = 1.$$

115. Graph functions. Explain how the graphs are related.




The graph of $g(x) = x^2 - 3$ is the graph of $f(x) = x^2$

shifted down 3 units. The graph of $h(x) = x^2 + 2$ is

2

the graph of $f(x) = x^2$ shifted up 2 units.

116. Graph functions. Explain how the graphs are related.



The graph of $g(x) = (x-3)^2$ is the graph of

2

$f(x) = x^2$ shifted 3 units to the right. The graph of

$h(x) = (x+2)^2$ is the graph of $f(x) = x^2$ shifted 2

117. Find all fixed points.

$$\begin{aligned}
 a^2 + 3a - 3 &= a \\
 a^2 + 2a - 3 &= 0 \\
 (a-1)(a+3) &= 0 \\
 a &= 1 \text{ or } a = -3
 \end{aligned}$$

118. Find all fixed points.

$$\begin{aligned}
 \frac{a}{a+5} &= a \\
 &= a(a+5) \\
 a &= a^2 + 5a \\
 0 &= a^2 + 4a \\
 0 &= a(a+4)
 \end{aligned}$$

$$a = 0 \text{ or } a = -4$$

a. Write the function.

$$A = xy$$

$\frac{1}{2}$

$$A(x) = x \left(-\frac{1}{2}x + 4 \right)$$

$$A(x) = -\frac{1}{2}x^2 + 4x$$

Complete the table.

	12467			
Area	3.5	6	8	6 3.5

c. Find the domain. Domain: $[0, 8]$.

a. Write the function.

$$m = \frac{0-2}{-2-0} = -2$$

$$\begin{aligned}
 PB &= \frac{x-2}{x-2} \\
 m_{AB} &= \frac{0-y}{-2-0} = \frac{-y}{-2} = \frac{y}{2}
 \end{aligned}$$

$$m_{PB} = m_{AB}$$

$$\frac{-2}{x-2} = \frac{-y}{2}$$

$$\frac{2x}{x-2} = y$$

$$\text{Area} = \frac{1}{2}bh = \frac{1}{2}xy$$

$$\begin{aligned}
 \frac{1}{2}x \frac{2x}{x-2} \\
 \frac{x^2}{x-2}
 \end{aligned}$$

b. Find the domain. Domain: $(2, \infty)$

units to the left.

121. a. Write the function.

Circle
 $C = 2r$
 $x = 2r$
 $r = \frac{x}{2}$

Square
 $C = 4s$
 $20 - x = 4s$
 $s = 5 - \frac{x}{4}$

Area = $r^2 = \left(\frac{x}{2}\right)^2$ Area = $s^2 = \left(5 - \frac{x}{4}\right)^2$

$= \frac{x^2}{4}$ $= 25 - \frac{5x}{4} + \frac{x^2}{16}$

Total Area = $4 \cdot \frac{x^2}{4} + 25 - 2 \cdot \frac{5x}{4} + 16 \cdot \frac{x^2}{16}$

$\left(\frac{1}{4} + 16\right)x^2 -$

$2\frac{5}{4}x + 25$ b. Complete the table.

x	0	4	8	12	16	20
Total Area	25	17.27	14.09	15.46	21.37	31.83

c. Find the domain. Domain: [0, 20].

a. Let $m = 10$, $d = 7$, $c = 19$, and $y = 41$. Then

$$z = \frac{13m-1}{5} + \frac{y}{4} + \frac{c}{4} + d+y-2c$$

$$\frac{13 \cdot 10 - 1}{5} + \frac{41}{4} + \frac{19}{4} + 7 + 41 - 2 \cdot 19$$

$$25 + 10 + 4 + 7 + 41 - 38$$

$$49$$

The remainder of 49 divided by 7 is 0.

Thus December 7, 1941, was a Sunday.

This one is tricky. Because we are finding a date in the month of January, we must use 11 for the month and we must use the previous year, which is 2019. Thus we let $m = 11$, $d = 1$, $c = 20$, and $y = 19$.

Then

$$z = \frac{13m-1}{5} + \frac{y}{4} + \frac{c}{4} + d+y-2c$$

$$\frac{13 \cdot 11 - 1}{5} + \frac{19}{4} + \frac{20}{4} + 1 + 19 - 2 \cdot 20$$

$$28 + 4 + 5 + 1 + 19 - 40$$

$$17$$

The remainder of 17 divided by 7 is 3.

Thus January 1, 2020, will be a Wednesday.

Let $m = 5$, $d = 4$, $c = 17$, and $y = 76$. Then

$$z = \frac{13m-1}{5} + \frac{y}{4} + \frac{c}{4} + d+y-2c$$

$$\frac{13 \cdot 5 - 1}{5} + \frac{76}{4} + \frac{17}{4} + 4 + 76 - 2 \cdot 17$$

$$12 + 19 + 4 + 4 + 76 - 34$$

$$81$$

The remainder of 81 divided by 7 is 4.

Thus July 4, 1776 was a Thursday.

Answers will vary.

Prepare for Section

2.3 P1. $d = 5 - (-2) = 7$

P2. The product of any number and its negative reciprocal is -1 . For example,

$$\frac{1}{-7} \cdot 7 = -1$$

P3. $\frac{-4-4}{2-(-3)} = \frac{-8}{5}$

P4. $y - 3 = -2(x - 3)$
 $y - 3 = -2x + 6$
 $= -2x + 9$

P5. $3x - 5y = 15$
 $-5y = -3x + 15$
 $= \frac{3}{5}x - 3$

P6. $y = 3x - 2(5 - x)$
 $= 3x - 2(5 - x)$
 $= 3x - 10 + 2x$
 $= 5x$
 $= x$

Section 2.3 Exercises

If a line has a negative slope, then as the value of y increases, the value of x decreases.

If a line has a positive slope, then as the value of y decreases, the value of x decreases.

The graph of a line with zero slope is horizontal.

The graph of a line whose slope is undefined is vertical.

Determine the slope and y-intercept.

$$y = 4x - 5 : m = 4, \text{ y-intercept: } (0, -5)$$

Chapter 2 Functions and Graphs 6.

Determine the slope and y-intercept.

$$y = 3 - 2x : m = -2, \text{ y-intercept: } (0, 3) \quad 7.$$

Determine the slope and y-intercept.

$$(x) = \frac{2}{3}x - \frac{2}{3} : m = \frac{2}{3}, \text{ y-intercept: } (0, 0)$$

Determine the slope and y-intercept.

$$(x) = -1 : m = 0, \text{ y-intercept: } (0, -1)$$

Determine whether the graphs are parallel, perpendicular, or neither.

$$= 3x - 4 : m = 3, \\ = -3x + 2 : m = -3$$

The graphs are neither parallel nor perpendicular

Determine whether the graphs are parallel, perpendicular, or neither.

$$y = -\frac{2}{3}x + 1 : m = -\frac{2}{3},$$

$$y = 2 - \frac{2}{3}x : m = -\frac{2}{3}$$

The graphs are parallel.

Determine whether the graphs are parallel, perpendicular, or neither.

$$f(x) = 3x - 1 : m = 3, \\ \frac{x}{3} - \frac{1}{3}$$

$$y = -3x - 1 : m = -3$$

$$3\left(-\frac{1}{3}\right) = -1$$

The graphs are perpendicular.

Determine whether the graphs are parallel, perpendicular, or neither.

$$(x) = \frac{4}{3}x + 2 : m = \frac{4}{3}, \\ \frac{3}{4}$$

$$(x) = 2 - \frac{3}{4}x : m = -\frac{3}{4}$$

$$\frac{3}{4} \cdot \frac{4}{3} = 1$$

Find the slope.

$$m = \frac{1-4}{5-(-2)} = \frac{-3}{7} = -\frac{3}{7}$$

Find the slope. m

$$\frac{2-0}{0-4} = -\frac{1}{2}$$

Find the slope.

$$m = \frac{4-4}{2-(-3)} = \frac{0}{5} = 0$$

Find the slope.

$$m = \frac{-7-2}{3-3} = \frac{-9}{0} \text{ undefined}$$

Find the slope.

$$m = \frac{0-0}{3-0} = \frac{0}{3} = 0$$

Find the slope.

$$m = \frac{-2-4}{-4-(-3)} = \frac{-6}{-1} = 6$$

Find the slope.

$$m = \frac{4-(-1)}{-3-(-5)} = \frac{5}{2}$$

Find the slope.


$$m = \frac{-7-1}{\frac{2}{3}-\frac{2}{3}} = \frac{-8}{0} = \text{undefined}$$

22. Find the slope.

$$m = \frac{2-4}{\frac{7-1}{4}-\frac{5}{2}} = \frac{-2}{\frac{5}{4}-\frac{10}{4}} = \frac{-2}{-\frac{5}{4}} = \frac{8}{5}$$

Find the slope, y-intercept, and graph. $y = 2x - 4$

$$m = 2, \text{ y-intercept } (0, -4)$$




The graphs are perpendicular.

Find the slope.

$$m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{7 - 4}{1 - 3} = \frac{3}{-2} = -\frac{3}{2}$$


Find the slope, y-intercept, and graph. $y = -x + 1$

$m = -1$, y-intercept $(0, 1)$




Find the slope, y-intercept, and graph. $y = 4\frac{3}{4}x + 1$

$m = 4\frac{3}{4}$, y-intercept $(0, 1)$




Find the slope, y-intercept, and graph. $y = -2\frac{3}{4}x + 4$

$m = -2\frac{3}{4}$, y-intercept $(0, 4)$




Find the slope, y-intercept, and graph. $y = -2x + 3$

$m = -2$, y-intercept $(0, 3)$




Find the slope, y-intercept, and graph. $y = 3x - 1$

$m = 3$, y-intercept $(0, -1)$




Find the slope, y-intercept, and graph. $y = 3$

$m = 0$, y-intercept $(0, 3)$




Find the slope, y-intercept, and graph. $y = -2$

$m = 0$, y-intercept $(0, -2)$




Find the slope, y-intercept, and graph. $y = 2x$

$m = 2$, y-intercept $(0, 0)$




Find the slope, y-intercept, and graph. $y = -3x$

$m = -3$, y-intercept $(0, 0)$




Find the slope, y-intercept, and graph. $y = x$

$m = 1$, y-intercept $(0, 0)$



Find the slope, y-intercept, and graph. $y = -x$

$m = -1$, y-intercept $(0, 0)$




Write slope-intercept form, find intercepts, and graph.

$$2x + y = 5$$

$$= -2x + 5$$

x-intercept $(\frac{5}{2}, 0)$, y-intercept $(0, 5)$




Write slope-intercept form, find intercepts, and graph.

$$x - y = 4$$

$$= x - 4$$

x-intercept $(4, 0)$, y-intercept $(0, -4)$



Write slope-intercept form, find intercepts, and graph.


$$4x + 3y - 12 = 0$$

$$3y = -4x + 12$$

$$\underline{3}$$

$$= -\frac{4}{3}x + 4$$

x-intercept $(3, 0)$, y-intercept $(0, 4)$



Write slope-intercept form, find intercepts, and graph.


$$2x + 3y + 6 = 0$$

$$3y = -2x - 6$$

$$\underline{3}$$

$$= -\frac{2}{3}x - 2$$

x-intercept $(-3, 0)$, y-intercept $(0, -2)$



Write slope-intercept form, find intercepts, and graph.


$$2x - 5y = -15$$

$$-5y = -2x - 15$$

$$\underline{5}$$

$$y = \frac{2}{5}x + 3$$

x-intercept $(-\frac{15}{2}, 0)$, y-intercept $(0, 3)$



Write slope-intercept form, find intercepts, and graph.

$$3x - 4y = 8$$


$$-4y = -3x + 8$$

$$y = \frac{3}{4}x - 2$$

$$\underline{4}$$

$$\frac{8}{4}$$

x-intercept $(\frac{8}{3}, 0)$, y-intercept $(0, -2)$




Write slope-intercept form, find intercepts, and graph.

$$+2y = 6$$

$$= -\frac{1}{2}x + 3$$

x-intercept $(6, 0)$, y-intercept $(0, 3)$




Write slope-intercept form, find intercepts, and graph.

$$-3y = 9$$

$$= \frac{1}{3}x - 3$$

x-intercept $(9, 0)$, y-intercept $(0, -3)$



Find the equation.

Use $y = mx + b$ with $m = 1, b = 3$.
 $= x + 3$

Find the equation.

Use $y = mx + b$ with $m = -2, b = 5$.
 $= -2x + 5$

Find the equation.

Use $y = mx + b$ with $m = 4\frac{3}{4}, b = \frac{1}{2}$.
 $= 4\frac{3}{4}x + \frac{1}{2}$

Find the equation.

Use $y = mx + b$ with $m = -\frac{2}{3}, b = 4\frac{3}{4}$.
 $= -\frac{2}{3}x + 4\frac{3}{4}$

Find the equation.

Use $y = mx + b$ with $m = 0, b = 4$.
 $y = 4$

Find the equation.

Use $y = mx + b$ with $m = \frac{1}{2}, b = -1$.
 $= \frac{1}{2}x - 1$

Find the equation.

$-2 = -4(x - (-3))$
 $-2 = -4x - 12$
 $= -4x - 10$

Find the equation.

$+1 = -3(x + 5)$
 $= -3x - 15 - 1$
 $= -3x - 16$

Find the equation.

$m = \frac{4-1}{-1-3} = \frac{3}{-4} = -\frac{3}{4}$
 $y - 1 = -\frac{3}{4}(x - 3)$
 $y = -\frac{3}{4}x + \frac{9}{4} + \frac{4}{4}$

Find the equation.

$m = \frac{-8 - (-6)}{2-5} = \frac{-2}{-3} = \frac{2}{3}$
 $y - (-6) = \frac{2}{3}(x - 5)$
 $y + 6 = \frac{2}{3}x - \frac{10}{3}$
 $y = \frac{2}{3}x - \frac{10}{3} - 6$
 $y = \frac{2}{3}x - \frac{28}{3}$

Find the equation.

$= \frac{-1-11}{2-7-55} = \frac{-12}{-55} = \frac{12}{55}$
 $y - 11 = \frac{12}{55}(x - 7)$
 $y = \frac{12}{55}x - \frac{84}{55} + \frac{55}{55}$
 $y = \frac{12}{55}x - \frac{29}{55}$

Find the equation.

$m = \frac{-4-6}{-3-(-5)} = \frac{-10}{2} = -5$
 $y - 6 = -5(x + 5)$
 $-6 = -5x - 25$
 $= -5x - 25 + 6$
 $y = -5x - 19$

Find the equation.

$y = 2x + 3$ has slope $m = 2$.
 $y - y_1 = 2(x - x_1)$
 $+4 = 2(x - 2)$
 $y + 4 = 2x - 4$
 $y = 2x - 8$

Find the equation.

$y = -x + 1$ has slope $m = -1$.
 $y - y_1 = -1(x - x_1)$
 $y - 4 = -1(x + 2)$
 $y - 4 = -x - 2$
 $= -x + 2$

$$= -4\frac{3}{4}x + \frac{13}{4}$$

Chapter 2 Functions and Graphs

Find the equation.

$$y = -\frac{3}{4}x + 3 \text{ has slope } m = -\frac{3}{4}.$$

$$y - y_1 = -\frac{3}{4}(x - x_1)$$

$$1 \quad 4 \quad 1$$

$$y - 2 = -\frac{3}{4}(x - 3)$$

$$= -\frac{3}{4}x + 1$$

Find the equation.

$$y = \frac{2}{3}x - 1 \text{ has slope } m = \frac{2}{3}.$$

$$y - y_1 = \frac{2}{3}(x - x_1)$$

$$y + 5 = \frac{2}{3}(x + 3)$$

$$= \frac{2}{3}x - 3$$

Find the equation.

$$2x - 5y = 2$$

$$-5y = -2x + 2$$

$$y = \frac{2}{5}x - \frac{2}{5} \text{ has slope } m = \frac{2}{5}.$$

$$\frac{2}{5}$$

$$y - y_1 = \frac{2}{5}(x - x_1)$$

$$y - 2 = \frac{2}{5}(x - 5)$$

$$\frac{2}{5}$$

$$y = \frac{2}{5}x$$

Find the equation.

$$x + 3y = 4$$

$$3y = -x + 4$$

$$\frac{1}{3} \quad 4$$

$$\frac{1}{3}$$

$$y = -\frac{1}{3}x + \frac{4}{3} \text{ has slope } m = -\frac{1}{3}.$$

$$\frac{1}{3}$$

$$y - y_1 = -\frac{1}{3}(x - x_1)$$

Find the equation.

$$y = 2x - 5 \text{ has perpendicular slope } m = -\frac{1}{2}.$$

$$y - y_1 = -\frac{1}{2}(x - x_1)$$

$$y + 4 = -\frac{1}{2}(x - 3)$$

$$y + 4 = -\frac{1}{2}x + \frac{3}{2}$$

$$= -\frac{1}{2}x - \frac{5}{2}$$

Find the equation.

$$y = -x + 3 \text{ has perpendicular slope } m = 1.$$

$$y - y_1 = 1(x - x_1)$$

$$y - 2 = 1(x + 5)$$

$$y - 2 = x + 5$$

$$= x + 7$$

Find the equation.

$$y = -\frac{3}{4}x + 1 \text{ has perpendicular slope } m = \frac{4}{3}.$$

$$y - y_1 = \frac{4}{3}(x - x_1)$$

$$y = \frac{4}{3}x + 8$$

Find the equation.

$$3x - 2y = 5$$

$$-2y = -3x + 5$$

$$y = \frac{3}{2}x - \frac{5}{2} \text{ has perpendicular slope } m = -\frac{2}{3}.$$

$$y - y_1 = -\frac{2}{3}(x - x_1)$$

$$y - 4 = -\frac{2}{3}(x + 3)$$

$$y - 4 = -\frac{2}{3}x - 2$$

$$= -\frac{2}{3}x + 2$$

$$+ 1 = - 3(x - x_1)$$

$$+ 1 = - \frac{1}{3}(x + 3)$$

$$= - \frac{1}{3}x - 1$$

$$\frac{1}{3}x - 2$$

Find the equation.

$$\begin{aligned}
 -x - 4y &= 6 \\
 -4y &= x + 6 \\
 y &= -\frac{1}{4}x - \frac{3}{2} \text{ has perpendicular slope } m = 4.
 \end{aligned}$$

$$\begin{aligned}
 y - y_1 &= 4(x - x_1) \\
 y - 2 &= 4(x - 5) \\
 y - 2 &= 4x - 20 \\
 &= 4x - 18
 \end{aligned}$$

Find the equation.


$$\begin{aligned}
 5x - y &= 2 \\
 y &= -5x + 2 \\
 y &= 5x - 2 \text{ has perpendicular slope } m = -\frac{1}{5}. \\
 &\frac{1}{5}
 \end{aligned}$$

$$\begin{aligned}
 y - y_1 &= -\frac{1}{5}(x - x_1) \\
 + 2 &= -\frac{1}{5}(x - 10) \\
 y + 2 &= -\frac{1}{5}x + 2 \\
 &= -\frac{1}{5}x
 \end{aligned}$$

Find the zero of f .

$$\begin{aligned}
 f(x) &= 3x - 12 \\
 3x - 12 &= 0 \\
 3x &= 12 \\
 x &= 4
 \end{aligned}$$

The x -intercept of the graph of $f(x)$ is (4, 0).




$$\begin{aligned}
 X_{\min} &= 4, X_{\max} = 6, X_{\text{scl}}=2, \\
 Y_{\min} &= -4, Y_{\max} = 4, Y_{\text{scl}} = 2
 \end{aligned}$$

Find the zero of f .

$$\begin{aligned}
 f(x) &= -2x - 4 \\
 -2x - 4 &= 0 \\
 -2x &= 4 \\
 x &= -2
 \end{aligned}$$

The x -intercept of the graph of $f(x)$ is (-2, 0).




$$\begin{aligned}
 X_{\min} &= 4, X_{\max} = 6, X_{\text{scl}}=2, \\
 Y_{\min} &= -2, Y_{\max} = 2, Y_{\text{scl}} = 2
 \end{aligned}$$

Find the zero of f .

$$\begin{aligned}
 &\frac{1}{4} \\
 f(x) &= 4x + 5
 \end{aligned}$$

$$\begin{aligned}
 \frac{1}{4}x + 5 &= 0 \\
 \frac{1}{4}x &= -5 \\
 &= -20
 \end{aligned}$$

The x -intercept of the graph of $f(x)$ is (-20, 0).



$$X_{\min} = -30, X_{\max} = 30, X_{\text{scl}} = 10,$$


$$Y_{\min} = -10, Y_{\max} = 10, Y_{\text{scl}} = 1$$

Find the zero of f .

$$f(x) = -\frac{1}{3}x + 2$$

$$\begin{aligned}
 \frac{1}{3}x + 2 &= 0 \\
 \frac{1}{3}x &= -2 \\
 x &= 6
 \end{aligned}$$

The x -intercept of the graph of $f(x)$ is (6, 0).



$$X_{\min} = 2, X_{\max} = 8, X_{\text{scl}} = 2,$$

$$Y_{\min} = 0, Y_{\max} = 4, Y_{\text{scl}} = 2$$

Find the slope and explain the meaning.

$$= \frac{1505 - 1482}{28 - 20} = 2.875$$

The value of the slope indicates that the speed of sound

in water increases 2.875 m/s for a one-degree Celsius increase in temperature.

Find the slope and explain the meaning.

$$m = \frac{40 - 10}{100 - 25} = 0.4$$

The value of the slope indicates that the file is being downloaded at 0.4 megabytes per second.

$$\mathbf{a.} \quad m = \frac{31 - 20}{23 - 12} = 1$$

$$(c) - 20 = 1(c - 12)$$

$$H(c) = c + 8$$

$$H(19) = (19) + 8 = 27 \text{ mpg}$$

$$\underline{864.9 - 1008.1}$$

$$\mathbf{a.} \quad m = \frac{864.9 - 1008.1}{2011 - 2007} = -35.8$$

$$C(t) - 864.9 = -35.8(t - 2011)$$

$$C(t) = -35.8t + 72,858.7$$

$$750 = -35.8t + 72,858.7 - 72,108.7$$

$$= -35.8t$$

$$2014.2 \gg t$$

The debt will fall below \$750 billion in 2014.

$$\mathbf{a.} \quad m = \frac{316,500 - 279,200}{2020 - 2010} = 3730$$

$$N(t) - 279,200 = 3730(t - 2010)$$

$$N(t) = 3730t - 7,218,100$$

$$300,000 = 3730t - 7,218,100$$

$$7,518,100 = 3730t$$

$$2015.6 \gg t$$

The number of jobs will exceed 300,000 in 2015.

$$\mathbf{a.} \quad m = \frac{2200 - 2150}{15 - 20} = -10$$

$$(t) - 2200 = -10(t - 15) T$$

$$(t) = -10t + 2350$$

The value of the slope means that the temperature is

decreasing at a rate of 10 F per minute.

$$T(180) = -10(180) + 2350 = 550 \text{ F}$$

After 3 hours, the temperature will be 550 F.

$$\mathbf{a.} \quad m = \frac{240 - 180}{18 - 16} = 30$$

$$B(d) - 180 = 30(d - 16)$$

$$B(d) = 30d - 300$$

The value of the slope means that a 1-inch increase in the diameter of a log 32 ft long results in an increase of 30 board-feet of lumber that can be obtained from the log.

$$B(19) = 30(19) - 300 = 270 \text{ board feet}$$

$$\mathbf{a.} \quad m = \frac{1640 - 800}{60 - 40} = 42$$

$$E(T) - 800 = 42(T - 40)$$

$$E(T) = 42T - 880$$

The value of the slope means that an additional 42 acre-feet of water evaporate for a one F increase in temperature.

$$E(75) = 42(75) - 880 = 2270 \text{ acre-feet}$$

Line *A* represents Michelle.

Line *B* represents Amanda.

Line *C* represents the distance between Michelle and Amanda.

$$\mathbf{a.} \quad m_{AB} = \frac{1}{8} - \frac{9}{6} = -4 \text{ F}$$

$$m_{AB} = \frac{1}{8} - \frac{9}{6} = -4 \text{ F}$$

$$m = \frac{-4 - 5}{5 - 6} = 9 \text{ F}$$

The temperature changed most rapidly between points *D* and *E*.

c. The temperature remained constant (zero slope)

between points *C* and *D*.

$$\underline{80.5 - 19.9}$$

$$\mathbf{81. a.} \quad m = \frac{80.5 - 19.9}{0 - 65} \gg -0.9323$$

$$-80.5 = -0.9323(x - 0) y$$

$$= -0.9323x + 80.5$$

$$\mathbf{b.} \quad y = -0.9323(25) + 80.5 = 57.19 \gg 57 \text{ years}$$

a. $m = \frac{75.5 - 17.2}{65} \approx -0.8969$

$-75.5 = -0.8969(x - 0) y$
 $= -0.8969x + 75.5$

b. $y = -0.8969(25) + 75.5 = 53.08 \approx 53$ years

Determine the profit function and break-even point.

$P(x) = 92.50x - (52x + 1782)$

$P(x) = 92.50x - 52x - 1782$

$P(x) = 40.50x - 1782$

$40.50x - 1782 = 0$

$40.50x = 1782$

$x = 40.50 \frac{1782}{40.50}$

$x = 44$, the break-even point

Determine the profit function and break-even point.

$P(x) = 124x - (78.5x + 5005)$

$P(x) = 124x - 78.5x - 5005$

$P(x) = 45.5x - 5005$

$45.5x - 5005 = 0$

$45.5x = 5005$

$x = \frac{5005}{45.5}$

$x = 110$, the break-even point

Determine the profit function and break-even point.

$P(x) = 259x - (180x + 10,270)$

$P(x) = 259x - 180x - 10,270$

$P(x) = 79x - 10,270$

$79x - 10,270 = 0$

$79x = 10,270$

$x = \frac{10,270}{79}$

$x = 130$, the break-even point

Determine the profit function and break-even point.

$P(x) = 14,220x - (8010x + 1,602,180)$

$P(x) = 14,220x - 8010x - 1,602,180$

$P(x) = 6210x - 1,602,180$

$6210x - 1,602,180 = 0$

$6210x = 1,602,180$

$x = \frac{1,602,180}{6210}$

$x = 258$, the break-even point

a. $C(0) = 8(0) + 275 = 0 + 275 = \275

$C(1) = 8(1) + 275 = 8 + 275 = \283

$C(10) = 8(10) + 275 = 80 + 275 = \355

The marginal cost is the slope of $C(x) = 8x + 275$,

which is \$8 per unit.

a. $R(0) = 210(0) = \$0$

$R(1) = 210(1) = \$210$

$R(10) = 210(10) = \$2100$

The marginal revenue is the slope of $R(x) = 210x$,

which is \$210 per unit.

a. $C(t) = 19,500.00 + 6.75t$

$R(t) = 55.00t$

$P(t) = R(t) - C(t)$

$P(t) = 55.00t - (19,500.00 +$

$6.75t) P(t) = 55.00t - 19,500.00 -$

$6.75t P(t) = 48.25t - 19,500.00$

$48.25t = 19,500.00$

$= \frac{19,500.00}{48.25}$

$t = 404.1451$ days ≈ 405 days

$m = \frac{117,500 - 98,000}{35,000 - 32,000} = \frac{19,500}{3000} = 6.5$

$P(s) - 98,000 = 6.5(s - 32,000) P(s)$

$= 6.5s - 208,000 + 98,000 P(s)$

$= 6.5s - 110,000$

$P(50,000) = 6.5(50,000) - 110,000$

$325,000 - 110,000$

$\$215,000$

Let $6.5s - 110,000 = 0$. Then

$6.5s = \frac{110,000}{6.5}$
 $s = \frac{110,000}{6.5} \approx 16,924$ subscribers

The equation of the line through $(0,0)$ and $P(3,4)$

has slope $\frac{4}{3}$.

The path of the rock is on the line through $P(3,4)$ with

slope $-4\frac{3}{4}$, so $y - 4 = -4\frac{3}{4}(x - 3)$.


$$\begin{aligned} y - 4 &= -4\frac{3}{4}x + \frac{9}{4} \\ &= -4\frac{3}{4}x + \frac{9}{4} + \frac{3}{4} \\ 4y &= -4\frac{3}{4}x + \frac{25}{4} \end{aligned}$$

The point where the rock hits the wall at $y = 10$ is the point of intersection of $y = -4\frac{3}{4}x + \frac{25}{4}$ and $y = 10$.

$$\begin{aligned} 4\frac{3}{4}x + \frac{25}{4} &= 10 \\ -3x + 25 &= 40 \\ -3x &= 15 \\ x &= -5 \text{ feet} \end{aligned}$$

Therefore the rock hits the wall at $(-5, 10)$.

The x -coordinate is -5 .



The equation of the line through $(0,0)$ and

$P(\sqrt{15}, 1)$ has slope $\frac{1}{\sqrt{15}}$.

The path of the rock is on the line through

$P(\sqrt{15}, 1)$ with slope $-15\sqrt{15}$


$$\begin{aligned} y - 1 &= -15\sqrt{15}(x - \sqrt{15}) \\ y - 1 &= -15\sqrt{15}x + 15 \\ &= -15\sqrt{15}x + 15 + 1 \\ &= -15\sqrt{15}x + 16 \end{aligned}$$

The point of impact with the wall at $y = 14$ is the point of intersection of $y = -15\sqrt{15}x + 16$ and $y = 14$ intersect.

$$\begin{aligned} \sqrt{15}x + 16 &= 14 \\ \sqrt{15}x &= -2 \\ x &= \frac{-2}{\sqrt{15}} \approx -0.52 \text{ feet} \end{aligned}$$

$$\begin{aligned} \frac{2}{\sqrt{15}} &= \frac{2\sqrt{15}}{15} \\ &= \frac{2\sqrt{15}}{15} \end{aligned}$$

The x -coordinate is $\frac{2}{\sqrt{15}}$ or approximately 0.52.



Find a .

$$\begin{aligned} (a) &= 2a + 3 = -1 \\ 2a &= -4 \\ a &= -2 \end{aligned}$$

Find a .

$$\begin{aligned} (a) &= 4 - 3a = 7 \\ -3a &= 3 - a \\ &= -1 \end{aligned}$$

Find a .

$$\begin{aligned} (a) &= 1 - 4a = 3 \\ -4a &= 2 \\ &= -\frac{1}{2} \end{aligned}$$

Find a .

$$\begin{aligned} (a) &= \frac{2}{3}a + 2 = 4 \\ \frac{2}{3}a &= 2 \\ a &= 3 \end{aligned}$$

$a, h = 1$ so

$$Q(1+h, [1+h]^2 + 1) = Q(2, 2^2 + 1) = Q(2, 5)$$

$$m = \frac{5 - 2}{2 - 1} = \frac{3}{1} = 3$$

$h = 0.1$ so

$$Q(1+h, [1+h]^2 + 1) = Q(1.1, 1.1^2 + 1) = Q(1.1, 2.21)$$

$$m = \frac{2.21 - 2}{1.1 - 1} = \frac{0.21}{0.1} = 2.1$$

$h = 0.01$ so

$$Q(1+h, [1+h]^2 + 1) = Q(1.01, 1.01^2 + 1) \\ Q(1.01, 2.0201)$$

$$m = \frac{2.0201 - 2}{1.01 - 1} = \frac{0.0201}{0.01} = 2.01$$

As h approaches 0, the slope of PQ seems to be approaching 2.

$$x_1 = 1, y_1 = 2, x_2 = 1+h, y_2 = [1+h]^2 + 1 \\ = \frac{y_2 - y_1}{x_2 - x_1} = \frac{[1+h]^2 + 1 - 2}{(1+h) - 1} = \frac{(1+2h+h^2) - 1}{h}$$

$$\frac{x_2 - x_1(1+h) - 1h}{h} = \frac{2h + h^2}{h} = 2 + h$$

a. $h = 1$, so

$$Q(-2+h, 9-[-2+h]^2) = Q(-2+1, 9-[-2+1]^2) \\ = Q(-1, 8)$$

$$m = \frac{8-5}{-1-(-2)} = \frac{3}{1} = 3$$

$h = 0.1$ so

$$Q(-2+h, 9-[-2+h]^2) = Q(-2+0.1, 9-[-2+0.1]^2) \\ = Q(-1.9, 5.39)$$

$$m = \frac{5.39-5}{-1.9-(-2)} = \frac{0.39}{0.1} = 3.9$$

$h = 0.01$ so

$$Q(-2+h, 9-[-2+h]^2) = Q(-2+0.01, 9-[-2+0.01]^2) \\ = Q(-1.99, 5.0399)$$

$$m = \frac{5.0399-5}{-1.99-(-2)} = \frac{0.0399}{0.01} = 3.99$$

As h approaches 0, the slope of PQ seems to be approaching 4.

$$x_1 = -2, y_1 = 5, x_2 = -2+h, y_2 = 9-[-2+h]^2 \\ = \frac{y_2 - y_1}{x_2 - x_1} = \frac{9-[-2+h]^2 - 5}{(-2+h) - (-2)} \\ = \frac{9-(4-4h+h^2) - 5}{h}$$

The slope of the line through $(3, 9)$ and (x, y)

$$\text{is } \frac{15}{2}, \text{ so } \frac{y-9}{x-3} = \frac{15}{2}.$$

Therefore

$$2(y-9) = 15(x-3) \\ 2y-18 = 15x-45 \\ 2y-15x+27=0 \\ 2x^2-15x+27=0 \quad \text{Substituting } y=x^2 \\ (2x-9)(x-3)=0 \\ x = \frac{9}{2} \text{ or } x=3$$

$$\text{If } x = \frac{9}{2}, y = x^2 = \left(\frac{9}{2}\right)^2 = \frac{81}{4} \quad \left(\frac{9}{2}, \frac{81}{4}\right).$$

If $x = 3, y = x^2 = (3)^2 = 9$ $(3, 9)$, but this is the

point itself. The point $\left(\frac{9}{2}, \frac{81}{4}\right)$ is on the graph of

$y = x^2$, and the slope of the line containing $(3, 9)$ and $\left(\frac{9}{2}, \frac{81}{4}\right)$ is $\frac{15}{2}$.

100. The slope of the line through $(3, 2)$ and (x, y) is $\frac{3}{8}$, so $\frac{y-2}{x-3} = \frac{3}{8}$.

$$8(y-2) = 3(x-3)$$

Therefore

$$8(y-2) = 3(x-3) \\ 8y-16 = 3x-9 \\ 8y = 3x+7 \\ 8\sqrt{x+1} = 3x+7 \quad \text{Substituting } y = x+1 \\ + \sqrt{x+1} = (3x+7)^2$$

$$64(x+1) = 9x^2 + 42x + 49$$

$$64x + 64 = 9x^2 + 42x + 49 \\ = 9x^2 - 22x - 15 \\ = (9x+5)(x-3)$$

$$x = -\frac{5}{9} \text{ or } x=3$$

$$\text{If } x = -\frac{5}{9},$$

h

$$= \frac{4h - h^2}{h} = 4 - h$$

h

Section 2.3 2 160

$$y = x + 1 \quad \sqrt{-9 + 9} = \sqrt{0} = 0 \quad \left(-9, 3 \right)$$

If $x = 3$, $y = \sqrt{x + 1} = \sqrt{3 + 1} = \sqrt{4} = 2 \quad (3, 2)$, but

Chapter 2 Functions and Graphs this

is the point itself.

The point $(-9, \frac{2}{3})$ is on the graph of $y = \frac{2}{3}x + 1$, and

the slope of the line containing $(3, 2)$ and

$(-9, \frac{2}{3})$ is $\frac{2}{3}$.

Find the equation.

$$2x + 3y = 5$$

$$3y = -2x + 5$$

$$y = -\frac{2}{3}x + \frac{5}{3} \text{ has slope } m = -\frac{2}{3}$$

$$y - y_1 = -\frac{2}{3}(x - x_1)$$

$$-(-1) = -\frac{2}{3}(x - 3)$$


$$\frac{2}{3}$$

$$+1 = -\frac{2}{3}x + 2$$

$$\frac{2}{3}$$

$$y = -\frac{2}{3}x + 1$$

$f(x) = -\frac{2}{3}x + 1$ has y-intercept $(0, 1)$.



Mid-Chapter 2 Quiz

Find the midpoint and length.

$$\left(\frac{-3+1}{2}, \frac{4+2}{2} \right) = (-1, 1)$$

$$= \sqrt{1+16} = \sqrt{17}$$

Find the center and radius.

$$x^2 + y^2 - 6x + 4y - 2 = 0$$

$$x^2 - 6x + y^2 + 4y = 2$$

$$x^2 - 6x + 9 + y^2 + 4y + 4 = 2 + 9 + 4$$

$$(x - 3)^2 + (y + 2)^2 = 15$$

center $(3, -2)$, radius $\sqrt{15}$

Evaluate.

$$f(x) = x^2 - 6x + 1$$

$$(-3) = (-3)^2 - 6(-3) + 1 = 9 + 18 + 1 = 28$$

Find the domain.

For $f(x) = \sqrt{x}$, the domain is $[0, \infty)$.

Find the zeros of f for $f(x) = x^2 - x - 12$.

$$f(x) = 0$$

$$x^2 - x - 12 = 0$$

$$(x + 3)(x - 4) = 0$$

$$x + 3 = 0 \quad x - 4 = 0$$

$$x = -3 \quad x = 4$$

Find the slope.

Prepare for Section 2.4

P1. $3x^2 + 10x - 8 = (3x - 2)(x + 4)$

P2. $x^2 - 8x = x^2 - 8x + 16 = (x - 4)^2$

P3. $f(-3) = 2(-3)^2 - 5(-3) - 7$

$$18 + 15 - 7$$

$$26$$

P4. $2x^2 - x = 1$

$$2x^2 - x - 1 = 0$$

$$(2x + 1)(x - 1) = 0$$

$$2x + 1 = 0 \quad x - 1 = 0$$

$$x = -\frac{1}{2} \quad x = 1$$

P5. $x^2 + 3x - 2 = 0$

$$m = \frac{3 - (-2)}{-2 - 8} = \frac{5}{-10} = -\frac{1}{2}$$

$$x = \frac{-3 \pm \sqrt{3^2 - 4(1)(-2)}}{2(1)}$$

$$\frac{\sqrt{\quad}}{\sqrt{\quad}}$$

P6. $53 = -16t^2 + 64t + 5$
 $16t^2 - 64t + 48 = 0$
 $t^2 - 4t + 3 = 0$
 $(t - 1)(t - 3) = 0$
 $t = 1, 3$

Section 2.4 Exercises

- d
- f
- b
- h
- g
- e
- c
- a

Write in standard form, find the vertex, the axis


of symmetry and graph.

$$(x) = (x^2 + 4x) + 1$$

$$(x^2 + 4x + 4) + 1 - 4$$

$$= (x + 2)^2 - 3 \quad \text{standard form,}$$

vertex (-2, -3), axis of symmetry $x = -2$




Write in standard form, find the vertex, the axis

of symmetry and graph.

$$(x) = (x^2 + 6x) - 1$$

$$= (x + 3)^2 - 10 \quad \text{standard form,}$$

vertex (-3, -10), axis of symmetry $x = -3$




Write in standard form, find the vertex, the axis of symmetry and graph.

$$(x) = (x^2 - 8x) + 5$$

$$(x^2 - 8x + 16) + 5 - 16$$

$$= (x - 4)^2 - 11 \quad \text{standard form,}$$

vertex (4, -11), axis of symmetry $x = 4$




Write in standard form, find the vertex, the axis of symmetry and graph.

$$(x) = (x^2 - 10x) + 3$$

$$(x^2 - 10x + 25) + 3 - 25$$

$$= (x - 5)^2 - 22 \quad \text{standard form,}$$

vertex (5, -22), axis of symmetry $x = 5$



Write in standard form, find the vertex, the axis of symmetry and graph.


$$(x) = (x^2 + 3x) + 1$$

$$\left(x^2 + 3x + \frac{9}{4} \right) + 1 - \frac{9}{4}$$

$$\left(x + \frac{3}{2} \right)^2 + \frac{4}{4} - \frac{9}{4}$$

$$= \left(x + \frac{3}{2} \right)^2 - \frac{5}{4} \quad \text{standard form,}$$

vertex $\left(-\frac{3}{2}, -\frac{5}{4} \right)$, axis of symmetry $x = -\frac{3}{2}$



Write in standard form, find the vertex, the axis of symmetry and graph.


$$(x) = (x^2 + 7x) + 2$$

$$\left(x^2 + 7x + \frac{49}{4}\right) + 2 - \frac{49}{4}$$

$$\left(x + \frac{7}{2}\right)^2 + \frac{8}{4} - \frac{49}{4}$$

$$= \left(x + \frac{7}{2}\right)^2 - \frac{41}{4} \text{ standard form, vertex}$$

$\left(-\frac{7}{2}, -\frac{41}{4}\right)$, axis of symmetry $x = -\frac{7}{2}$



Write in standard form, find the vertex, the axis of symmetry and graph.


$$(x) = -x^2 + 4x + 2$$

$$-(x^2 - 4x) + 2$$

$$-(x^2 - 4x + 4) + 2 + 4$$

$$= -(x - 2)^2 + 6 \text{ standard form,}$$

vertex (2, 6), axis of symmetry $x = 2$



Write in standard form, find the vertex, the axis of symmetry and graph.


$$(x) = -x^2 - 2x + 5$$

$$-(x^2 + 2x) + 5$$

$$-(x^2 + 2x + 1) + 5 + 1$$

$$= -(x + 1)^2 + 6 \text{ standard form,}$$

vertex (-1, 6), axis of symmetry $x = -1$



Write in standard form, find the vertex, the axis of symmetry and graph.

$$(x) = -3x^2 + 3x + 7$$


$$-3(x^2 - 1x) + 7$$

$$-3\left(x^2 - 1x + \frac{1}{4}\right) + 7 + 4$$

$$-3\left(x - \frac{1}{2}\right)^2 + \frac{28}{4} + 4$$

$$= -3\left(x - \frac{1}{2}\right)^2 + \frac{31}{4} \text{ standard form,}$$

vertex $\left(\frac{1}{2}, \frac{31}{4}\right)$, axis of symmetry $x = \frac{1}{2}$



Write in standard form, find the vertex, the axis of symmetry and graph.


$$(x) = -2x^2 - 4x + 5$$

$$-2(x^2 + 2x) + 5$$

$$-2(x^2 + 2x + 1) + 5 + 2$$

$$= -2(x + 1)^2 + 7 \text{ standard form,}$$

vertex (-1, 7), axis of symmetry $x = -1$



Find the vertex, write the function in standard form.

$$= -2a \frac{b}{2a} = -2(1) \frac{10}{2} = -5$$

$$= f(5) = (5)^2 - 10(5)$$

$$= 25 - 50 = -25$$

vertex (5, -25)

$$(x) = (x - 5)^2 - 25$$

Find the vertex, write the function in standard form.

$$= -\frac{b}{2a} = \frac{6}{2} = 3$$

$$= f(3) = (3)^2 - 6(3) = 9 - 18 = -9$$

vertex (3, -9)

$$f(x) = (x - 3)^2 - 9$$

Find the vertex, write the function in standard form.

$$= -\frac{b}{2a} = \frac{0}{2} = 0$$

$$= f(0) = (0)^2 - 10 = -10$$

vertex (0, -10)

$$f(x) = x^2 - 10$$

Find the vertex, write the function in standard form.

$$= -\frac{b}{2a} = \frac{0}{2} = 0$$

$$y = f(0) = (0)^2 - 4 = -4$$

vertex (0, -4)

$$f(x) = x^2 - 4$$

23. Find the vertex, write the function in standard form.

$$x = \frac{-b}{2a} = \frac{-6}{2(-1)-2} = \frac{-6}{-4} = 3$$

$$= f(3) = -(3)^2 + 6(3) + 1 = -9 + 18 + 1 = 10$$

10

vertex (3, 10)

$$f(x) = -(x - 3)^2 + 10$$

24. Find the vertex, write the function in standard form.

$$x = \frac{-b}{2a} = \frac{-4}{2(-1)-2} = \frac{-4}{-4} = 2$$

$$= f(2) = -(2)^2 + 4(2) + 1 = -4 + 8 + 1 = 5$$

Find the vertex, write the function in standard form.

$$= -\frac{b}{2a} = \frac{7}{2} = 3.5$$

$$= f\left(3.5\right) = 2\left(3.5\right)^2 - 3\left(3.5\right) + 7$$

$$= 2\left(12.25\right) - \frac{10.5}{2} + 7$$

$$= 24.5 - 5.25 + 7 = 16.25$$

$$\frac{16.25}{2}$$

vertex $\left(3.5, \frac{16.25}{2}\right)$

$$f(x) = 2\left(x - 3.5\right)^2 + \frac{16.25}{2}$$

Find the vertex, write the function in standard form.

$$= -\frac{b}{2a} = \frac{10}{-2(3)} = \frac{10}{-6} = -\frac{5}{3}$$

$$= f\left(-\frac{5}{3}\right) = 3\left(-\frac{5}{3}\right)^2 - 10\left(-\frac{5}{3}\right) + 2$$

$$= 3\left(\frac{25}{9}\right) - \frac{50}{-3} + 2$$

$$= \frac{25}{3} - \frac{50}{-3} + 2 = \frac{25}{3} - \frac{50}{-3} + \frac{6}{3}$$

$$= \frac{19}{3}$$

vertex $\left(-\frac{5}{3}, \frac{19}{3}\right)$

$$f(x) = 3\left(x - \left(-\frac{5}{3}\right)\right)^2 - \frac{19}{3}$$

Find the vertex, write the function in standard form.

$$x = \frac{-b}{2a} = \frac{-1}{2(-4)} = \frac{1}{8}$$

$$= f\left(\frac{1}{8}\right) = -4\left(\frac{1}{8}\right)^2 + \left(\frac{1}{8}\right) + 1$$

$$= -4\left(\frac{1}{64}\right) + \frac{1}{8} + 1$$

$$= -\frac{4}{64} + \frac{8}{64} + \frac{64}{64} = \frac{-4 + 8 + 64}{64} = \frac{68}{64} = \frac{17}{16}$$

$$8 + \frac{1}{8} + 1 = -16$$

5
vertex (2, 5)

$$f(x) = -(x - 2)^2 + 5$$

$$\frac{1}{2} + \frac{1}{6} = \frac{1}{6}$$

16
vertex $\left(8, \frac{17}{16}\right)$

$$f(x) = -4\left(x - \frac{1}{8}\right)^2 + \frac{17}{16}$$

Find the vertex, write the function in standard form.

$$x = \frac{-b}{2a} = \frac{-6}{2(-5)} = \frac{6}{-10} = -\frac{3}{5}$$

$$= f\left(-\frac{3}{5}\right) = -5\left(-\frac{3}{5}\right)^2 - 6\left(-\frac{3}{5}\right) + 3$$

$$= -5\left(\frac{9}{25}\right) + \frac{18}{5} + 3$$

$$= \frac{-9}{5} + \frac{18}{5} + \frac{15}{5} = \frac{9}{5}$$

vertex $\left(-\frac{3}{5}, \frac{24}{5}\right)$

$$f(x) = -5\left(x + \frac{3}{5}\right)^2 + \frac{24}{5}$$

Find the range, find x .

$$(x) = x^2 - 2x - 1$$

$$(x^2 - 2x) - 1$$

$$(x^2 - 2x + 1) - 1 - 1$$

$$(x - 1)^2 - 2$$

vertex (1, -2)

The y -value of the vertex is -2.

The parabola opens up since $a = 1 > 0$.

Thus the range is $\{y \mid y \geq -2\}$.

Find the range, find x .

$$(x) = -x^2 - 6x - 2$$

$$-(x^2 + 6x) - 2$$

$$-(x^2 + 6x + 9) - 2 + 9$$

$$-(x + 3)^2 + 7$$

vertex (3, 7)

The y -value of the vertex is 7.

The parabola opens down since $a = -1 < 0$.

Thus the range is $\{y \mid y \leq 7\}$.

Find the range, find x .

$$(x) = -2x^2 + 5x - 1$$

$$-2\left(x^2 - \frac{5}{2}x\right) - 1$$

$$-2\left(x^2 - \frac{5}{2}x + \frac{25}{16}\right) - 1 + 2\left(\frac{25}{16}\right)$$

$$-2\left(x - \frac{5}{4}\right)^2 + \frac{25}{8} - 1$$

$$-2\left(x - \frac{5}{4}\right)^2 + \frac{17}{8}$$

vertex $\left(\frac{5}{4}, \frac{17}{8}\right)$

$\frac{17}{8}$

The y -value of the vertex is $\frac{17}{8}$.

The parabola opens down since $a = -2 < 0$.

Thus the range is $\left\{y \mid y \leq \frac{17}{8}\right\}$.

Find the range, find x .

$$(x) = 2x^2 + 6x - 5$$

$$2(x^2 + 3x) - 5$$

$$2\left(x^2 + 3x + \frac{9}{4}\right) - 5 - 2\left(\frac{9}{4}\right)$$

$$2\left(x + \frac{3}{2}\right)^2 - \frac{10}{2} - \frac{9}{2}$$

$$2\left(x + \frac{3}{2}\right)^2 - \frac{19}{2}$$

vertex $\left(-\frac{3}{2}, -\frac{19}{2}\right)$

The y -value of the vertex is $-\frac{19}{2}$.

The parabola opens up since $a = 2 > 0$.

Find the real zeros and x -intercepts.

$$(x) = x^2 + 2x - 24$$

$$(x + 6)(x - 4)$$

$$x + 6 = 0 \quad x - 4 = 0$$

$$x = -6 \quad x = 4$$

$$(-6, 0) \quad (4, 0)$$

Find the real zeros and x -intercepts.

$$\begin{aligned} f(x) &= -x^2 + 6x + 7 \\ &= -(x^2 - 6x - 7) \\ &= -(x + 1)(x - 7) \\ x + 1 = 0 & \quad x - 7 = 0 \\ x = -1 & \quad x = 7 \\ (-1, 0) & \quad (7, 0) \end{aligned}$$

Find the real zeros and x -intercepts.

$$\begin{aligned} f(x) &= 2x^2 + 11x + 12 \\ &= (x + 4)(2x + 3) \\ x + 4 = 0 & \quad 2x + 3 = 0 \\ x = -4 & \quad x = -\frac{3}{2} \\ (-4, 0) & \quad \left(-\frac{3}{2}, 0\right) \end{aligned}$$

Find the real zeros and x -intercepts.

$$\begin{aligned} f(x) &= 2x^2 - 9x + 10 \\ &= (x - 2)(2x - 5) \\ x - 2 = 0 & \quad 2x - 5 = 0 \\ x = 2 & \quad x = \frac{5}{2} \\ (2, 0) & \quad \left(2, 0\right) \end{aligned}$$

Find the minimum or maximum.

$$\begin{aligned} f(x) &= x^2 + 8x \\ &= (x^2 + 8x + 16) - 16 \\ &= (x + 4) - 16 \end{aligned}$$

minimum value of -16 when $x = -4$
Find the minimum or maximum.

$$\begin{aligned} f(x) &= -x^2 - 6x \\ &= -(x^2 + 6x) \\ &= -(x^2 + 6x + 9) + 9 \end{aligned}$$

Find the minimum or maximum.

$$\begin{aligned} f(x) &= -x^2 + 6x + 2 \\ &= -(x^2 - 6x) + 2 \\ &= -(x^2 - 6x + 9) + 2 + 9 \\ &= -(x - 3)^2 + 11 \end{aligned}$$

maximum value of 11 when $x = 3$
Find the minimum or maximum.

$$\begin{aligned} f(x) &= -x^2 + 10x - 3 \\ &= -(x^2 - 10x) - 3 \\ &= -(x^2 - 10x + 25) - 3 + 25 \\ &= -(x - 5)^2 + 22 \end{aligned}$$

maximum value of 22 when $x = 5$
Find the minimum or maximum.

$$\begin{aligned} f(x) &= 2x^2 + 3x + 1 \\ &= 2\left(x^2 + 2\frac{3}{4}x\right) + 1 \\ &= 2\left(x^2 + 2\frac{3}{4}x + 16^9\right) + 1 - 2\left(16^9\right) \\ &= 2\left(x + 4\frac{3}{8}\right)^2 + \frac{8}{8} - \frac{9}{8} \\ &= \left(x + 4\frac{3}{8}\right)^2 - \frac{1}{8} \end{aligned}$$

minimum value of $-\frac{1}{8}$ when $x = -4\frac{3}{8}$

Find the minimum or maximum.

$$\begin{aligned} f(x) &= 3x^2 + x - 1 \\ &= 3\left(x^2 + \frac{1}{3}x\right) - 1 \\ &= 3\left(x^2 + \frac{1}{3}x + 36\right) - 1 - 3\left(36\right) \\ &= 3\left(x + 6\right)^2 - 12 - 12 \\ &= 3\left(x + 6\right)^2 - 24 \end{aligned}$$

minimum value of -24 when $x = -6$

$$3\left(x + \frac{1}{6}\right)^2 - 12 \frac{13}{6}$$

maximum value of 9 when $x = 3$

minimum value of $-12 \frac{13}{6}$ when $x = -\frac{1}{6}$

Chapter 2 Functions and Graphs

Find the minimum or maximum.

$$(x) = 5x^2 - 11$$

$$5(x^2) - 11$$

$$5(x - 0)^2 - 11$$

minimum value of -11 when $x = 0$

Find the minimum or maximum.

$$(x) = 3x^2 - 41$$

$$3(x^2) - 41$$

$$3(x - 0)^2 - 41$$

minimum value of -41 when $x = 0$

Find the minimum or maximum.

$$(x) = -\frac{1}{2}x^2 + 6x + 17$$

$$-\frac{1}{2}(x^2 - 12x) + 17$$

$$-\frac{1}{2}(x^2 - 12x + 36) + 17 + 18$$

$$-\frac{1}{2}(x - 6)^2 + 35$$

maximum value of 35 when $x = 6$

Find the minimum or maximum.

$$(x) = -4\frac{3}{4}x^2 - \frac{2}{5}x + 7$$

$$-\frac{3}{4}\left(x^2 + \frac{8}{15}x + \frac{16}{225}\right) + 7 + \frac{529}{75}$$

$$-\frac{3}{4}\left(x + \frac{4}{15}\right)^2 + \frac{529}{75}$$

maximum value of $\frac{529}{75} = 7\frac{4}{75}$ when $x = -\frac{4}{15}$

$$A(t) = -4.9t^2 + 90t + 9000$$

Microgravity begins and ends at a height of 9000 m.

$$9000 = -4.9t^2 + 90t + 9000$$

$$4.9t^2 - 90t = 0$$

$$t(4.9t - 90) = 0$$

$$t = 0 \quad 4.9t - 90 = 0$$

$$h(t) = -4.9t^2 + 12.8t$$

$$0 = -4.9t^2 + 12.8t$$

$$0 = t(-4.9t + 12.8)$$

$$t = 0 \quad -4.9t + 12.8 = 0$$

$$t = 0 \quad t = \frac{12.8}{4.9} \approx 2.6$$

The ball is in the air 2.6 seconds.

$$h(x) = -64x^2 + 27 = -64(x - 0)^2 + 27$$

The maximum height of the arch is 27 feet.

$$h(10) = -64(10)^2 + 27$$

$$-64(100) + 27$$

$$-1675 + 27$$

$$-1675 + \frac{432}{5} = -1675 + 86.4$$

$$= -1588.6$$

$$1616$$

$$3$$

$$h(x) = 8 = -64x^2 + 27$$

$$\frac{3}{64}$$

$$8 - 27 = -64x^2$$

$$-19 = -64x^2$$

$$\frac{64(-19)}{-64} = x^2$$

$$\sqrt{\frac{64(-19)}{-64}} = x$$

$$\sqrt{19} = x$$

$$t = 0 \quad t = \frac{90}{4.9} \approx 18.4$$

$$4$$

$$9$$

The time of microgravity is 18.4 seconds.

$$\frac{19.3}{3} = x$$

$$\frac{57}{3} = x$$

$$20.1 = x$$

$h(x) = 8$ when $x = 20.1$ feet

$$l + w = 240$$

$$w = 240 - l$$

$$A = l(240 - l) \quad A =$$

$$240l - l^2$$

$$\sqrt{\quad} \sqrt{\quad}$$

$$\sqrt{\quad}$$

$$A = -l^2 + 240l$$

$$A = -(l^2 - 240l + 120^2)$$

$$+120^2 A = -(l - 120)^2 + 120^2$$

Thus $l = 120$ and $w = 120$ produce the greatest area.

a. $3w + 2l = 600$

$$3w = 600 - 2l$$

$$= \frac{600 - 2l}{3}$$

$$A = w \cdot l$$

$$A = \left(\frac{600 - 2l}{3} \right) l$$

$$A = -\frac{2}{3}(l^2 - 300l)$$

$$A = -\frac{2}{3}(l^2 - 300l + 150^2)$$

+15,000 In standard form,

$$A = -\frac{2}{3}(l - 150)^2 + 15,000$$

The maximum area of 15,000 ft² is produced when $l = 150$ ft and the width $w = 100$ ft.

a. Find the temperature for maximum surviving larvae.

$$N(t) = -0.6t^2 + 32.1t - 350$$

$$= -0.6\left(t^2 - \frac{32.1}{0.6}t\right) - 350$$

$$= -0.6\left[t^2 - 53.5t + (26.75)^2\right] - 350 + 0.6(26.75)^2$$

$$= -0.6(t - 26.75)^2 + 79.3375$$

$$t = \frac{-32.1 \pm \sqrt{(-32.1)^2 - 4(-0.6)(-350)}}{2(-0.6)}$$

$$t = \frac{-32.1 \pm \sqrt{1030.41 - 840}}{-1.2}$$

$$t = \frac{-32.1 \pm 13.8}{-1.2}$$

$$t = \frac{-32.1 + 13.8}{-1.2} \quad \text{or} \quad t = \frac{-32.1 - 13.8}{-1.2}$$

$$= 15.25 \approx 15 \quad \text{or} \quad = 38.25 \approx 38$$

Thus the x -intercepts to the nearest whole number for $N(t)$ are (15, 0) and (38, 0).

When the temperature is less than 15 C or greater

than 38 C, none of the larvae survive.

a. $T(t) = -0.7t^2 + 9.4t + 59.3$

$$-0.7\left(t^2 - \frac{9.4}{0.7}t\right) + 59.3$$

$$-0.7\left(t^2 - \frac{94}{7}t\right) + 59.3$$

$$= -0.7\left(t^2 - \frac{94}{7}t + \left(\frac{47}{7}\right)^2\right) + 59.3 + 0.7\left(\frac{47}{7}\right)^2$$

$$-0.7\left(t - \frac{47}{7}\right)^2 + 90.857$$

$$-0.7\left(t - 6\frac{5}{7}\right)^2 + 91$$

The temperature is a maximum when $t = \frac{47}{7} = 6\frac{5}{7}$

The maximum number of larvae will survive at 27 C. A maximum of 79 larvae will survive.

$$N(t) = 0 = -0.6t^2 + 32.1t - 350$$

hours after 6:00 A.M. Note $7\frac{5}{6}$ (60 min) » 43

min. Thus the temperature is a maximum at

12:43 P.M.

The maximum temperature is approximately 91 F.

$$h(t) = -9.8t^2 + 100t$$

$$(t) = -9.8(t^2 - 10.2t)$$

$$h(t) = -9.8(t - 5.1)^2 + 254.9$$

The maximum height is 255 m.

$$55. t = -\frac{b}{2a} = -\frac{82.86}{2(-279.67)} = 0.14814$$

$$E(0.14814) = -279.67(0.14814)^2 + 82.86(0.14814)$$

6.
1

The maximum energy is 6.1 joules.

$$h(x) = -0.0009x^2 + 6$$

$$h(60.5) = -0.0009(60.5)^2 + 6 \approx 2.7$$

Since 2.7 is less than 5.4 and greater than 2.5, yes,

the pitch is a strike.

a. $E(v) = -0.018v^2 + 1.476v + 3.4$

$$-0.018 \left(v^2 - \frac{1.476}{0.018}v \right) + 3.4$$

$$-0.018(v^2 - 82v) + 3.4$$

$$-0.018(v^2 - 82v + 41^2) + 3.4 + 0.018(41)^2$$

$$-0.018(v - 41)^2 + 33.658$$

The maximum fuel efficiency is obtained at a speed of 41 mph.

The maximum fuel efficiency for this car, to the nearest mile per gallon, is 34 mpg.

$$h(x) = -0.0002348x^2 + 0.0375x$$

$$= -0.0002348(x^2 - 0.0002348^{-1} \cdot 0.0375x)$$

$$= -0.0002348 \left(x^2 - \frac{0.0375}{0.0002348}x \right)$$

$$= -0.0002348 \left(x^2 - 160x + 160^2 \right) + 0.0002348(160)^2$$

The maximum height of the field, to the nearest tenth of a foot, is 1.5 feet.

59. $-\frac{b}{2a} = -\frac{96}{2(-0.2)} = 740$

$$R(740) = 296(740) - 0.2(740)^2 = 109,520$$

Thus, 740 units yield a maximum revenue of \$109,520.

60. $-\frac{b}{2a} = -\frac{10}{2(-0.6)} = 675$

$$R(675) = 810(675) - 0.6(675)^2 = 273,375$$

Thus, 675 units yield a maximum revenue of \$273,375.

61. $-\frac{b}{2a} = -\frac{1.7}{2} = 85$

62. $-\frac{b}{2a} = -\frac{1.68}{2(-0.00014)} = 11,760$

$$P(11,760) = -4000(11,760) + 1.68(11,760)^2 = 5878.40$$

Thus, 11,760 units yield a maximum profit of \$5878.40.

$$P(x) = R(x) - C(x)$$

$$x(102.50 - 0.1x) - (52.50x + 1840)$$

$$-0.1x^2 + 50x - 1840$$

The break-even points occur when $R(x) = C(x)$

or $P(x) = 0$.

$$\text{Thus, } 0 = -0.1x^2 + 50x - 1840$$

$$x = \frac{-50 \pm \sqrt{50^2 - 4(-0.1)(-1840)}}{2(-0.1)}$$

$$= \frac{-50 \pm \sqrt{1764}}{-0.2}$$

$$-0.2$$

$$= \frac{-50 \pm 42}{-0.2}$$

$$x = 40 \text{ or } x = 460$$

The break-even points occur when $x = 40$ or $x = 460$.

$$P(x) = R(x) - C(x)$$

$$x(210 - 0.25x) - (78x + 6399)$$

$$-0.25x^2 + 132x - 6399$$

$$-\frac{b}{2a} = -\frac{32}{2(-0.25)} = 264$$

$$P(264) = -0.25(264)^2 + 132(264) - 6399 = 11,025$$

The break-even points occur when $P(x) = 0$.

$$\text{Thus, } 0 = -0.25x^2 + 132x - 6399$$

$$x = \frac{-132 \pm \sqrt{132^2 - 4(-0.25)(-6399)}}{2(-0.25)}$$

$$2a = 2(-0.01)$$

$$P(85) = -0.01(85)^2 + 1.7(85) - 48 = 24.25 \text{ Thus,}$$

85 units yield a maximum profit of \$24.25.

$$\frac{-132 \pm \sqrt{11025}}{-0.5}$$
$$= \frac{-132 \pm 105}{-0.5} x = 54 \text{ or } x = 474$$

The break-even points occur when $x = 54$ or $x = 474$.

Let x = the number of people that take the tour.

$$R(x) = x(15.00 + 0.25(60 - x))$$

$$x(15.00 + 15 - 0.25x)$$

$$-0.25x^2 + 30.00x$$

$$P(x) = R(x) - C(x)$$

$$(-0.25x^2 + 30.00x) - (180 + 2.50x)$$

$$-0.25x^2 + 27.50x - 180$$

$$-2 \frac{b}{a} = -2 \left(\frac{27.50}{-0.25} \right) = 55$$

$$P(55) = -0.25(55)^2 + 27.50(55) - 180 = \$576.25$$

The maximum profit occurs when $x = 55$ tickets.

Let x = the number of parcels.

$$R(x) = xp = x(22 - 0.01x) = -0.01x^2 + 22x$$

$$P(x) = R(x) - C(x)$$

$$(-0.01x^2 + 22x) - (2025 + 7x)$$

$$-0.01x^2 + 15x - 2025$$

c. $-\frac{b}{2a} = -\frac{15}{2(-0.01)} = 750$

$$P(750) = -0.01(750)^2 + 15(750) - 2025 = \$3600$$

$$p(750) = 22 - 0.01(750) = \$14.50$$

The break-even points occur when $R(x) = C(x)$.

$$-0.01x^2 + 22x = 2025 +$$

$$7x - 0.01x^2 + 15x - 2025 = 0$$

$$x = \frac{-(15) \pm \sqrt{15^2 - 4(-0.01)(-2025)}}{2(-0.01)}$$

$x = 150$ or $x = 1350$ are the break-even points. Thus

the minimum number of parcels the air freight

company must ship to break even is 150 parcels.

Let x = the number of \$0.05 price

reductions. Then the price per gallon is

$$p(x) = 3.95 - 0.05x$$

The number of gallons sold each day

$$\text{is } q(x) = 10,000 + 500x$$

Then the revenue is

$$R(x) = (3.95 - 0.05x)(10,000 + 500x)$$

$$-25x^2 + 1475x + 39,500$$

The cost is

$$C(x) = 2.75(10,000 + 500x) = 27,500 + 1375x$$

The profit equals revenue minus cost.

$$P(x) = R(x) - C(x)$$

$$= -25x^2 + 1475x + 39,500 - (27,500 + 1375x)$$

$$= -25x^2 + 100x + 12,000$$

The maximum profit occurs when

$$x = -\frac{b}{2a} = -\frac{100}{2(-25)} = 2$$

The price per gallon that maximizes profit is

$$p(2) = 3.95 - 0.05(2) = \$3.85$$

Let x = the number of \$10 price

reductions. Then the price per ticket is

$$p(x) = 390 - 10x$$

The number of tickets sold each day

$$\text{is } q(x) = 350 + 25x$$

Then the revenue is

$$R(x) = (390 - 10x)(350 + 25x)$$

$$-250x^2 + 6250x + 136,500$$

The cost is

$$C(x) = 150(350 + 25x) = 52,500 +$$

3750x The profit equals revenue minus cost.

$$P(x) = R(x) - C(x)$$

$$= -250x^2 + 6250x + 136,500 - (52,500 + 3750x)$$

$$= -250x^2 + 2500x + 84,000$$

The maximum profit occurs when

$$-\frac{b}{2a} = -\frac{2500}{2(-250)} = 5$$

$$x = -\frac{b}{2a} = -\frac{2500}{2(-250)} = 5$$

The price per ticket that maximizes profit

$$\text{is } p(5) = 390 - 10(5) = \$340$$

Chapter 2 Functions and Graphs

$$h(t) = -16t^2 + 128t$$

a. $-\frac{b}{2a} = -\frac{128}{2(-16)} = 4$ seconds

$$h(4) = -16(4)^2 + 128(4) = 256 \text{ feet}$$

$$0 = -16t^2 + 128t$$

$$0 = -16t(t - 8)$$

$$-16t = 0 \quad \text{or} \quad t - 8 = 0$$

$$t = 0 \quad \quad \quad t = 8$$

The projectile hits the ground at $t = 8$ seconds.

$$h(t) = -16t^2 + 64t + 80$$

a. $-\frac{b}{2a} = -\frac{64}{2(-16)} = 2$

$$h(2) = -16(2)^2 + 64(2) + 80$$

b. $\frac{b}{2a} = \frac{144}{2(-16)}$

2 seconds

$$0 = -16t^2 + 64t + 80$$

$$16(t^2 - 4t - 5) = 0$$

$$16(t - 5)(t + 1)$$

$$t - 5 = 0 \quad \text{or} \quad t + 1 = 0$$

$$t = 5 \quad \quad \quad t = -1 \text{ No}$$

The projectile has height 0 feet at $t = 5$ seconds.

$$y(x) = -0.014x^2 + 1.19x + 5$$

$$-\frac{b}{2a} = -\frac{1.19}{2(-0.014)}$$

42.5

$$y(42.5) = -0.014(42.5)^2 + 1.19(42.5) + 5$$

$$= 30.2875$$

» 30 feet

72. $h(t) = -6.6t^2 + 430t + 28,000$

$$-\frac{b}{2a} = -\frac{430}{2(-6.6)}$$

32.6

$$h(32.6) = -6.6(32.6)^2 + 430(32.6) + 28,000$$

$$35,000.784$$

» 35,000 feet

$$h(x) = -0.002x^2 - 0.03x + 8$$

$$h(39) = -0.002(39)^2 - 0.03(39) + 8 = 3.788 > 3$$

Solve for x using quadratic formula.

$$-0.002x^2 - 0.03x + 8 = 0$$

$$x^2 + 15x - 4000 = 0$$

$$x = \frac{-15 \pm \sqrt{(15)^2 - 4(1)(-4000)}}{2(1)}$$

$$\frac{-15 \pm 16\sqrt{225}}{2}$$

, use positive value of x

» 56.2

Yes, the conditions are satisfied.

$$4w + 2l = 1200$$

$$2l = 1200 - 4w$$

$$= \frac{1200 - 4w}{2}$$

$$= 600 - 2w$$

$$= w(600 - 2w) \quad A =$$

$$\frac{600w - 2w^2}{2} \quad A =$$

$$2w^2 + 600w \quad A =$$

$$2(w^2 - 300w)$$


$$A = -2(w^2 - 300w + 150^2) +$$

$$2 \cdot 150^2 \quad A = -2(w - 150)^2 + 45,000$$

Thus when $w = 150$, the length $l = \frac{1200 - 4(150)}{2} = 300$.

Thus the dimensions that yield the greatest enclosed area are $w = 150$ ft and $l = 300$ ft.

Find height and radius.



The perimeter is $48 = \pi r + h + 2r + h$.

Solve for h .

$$48 - \pi r - 2r = 2h$$

$$\frac{1}{2} (48 - \pi r - 2r) = h$$

Area = semicircle + rectangle

$$A = \frac{1}{2} \pi r^2 + 2rh$$

$$\frac{1}{2} \pi r^2 + 2r \left(\frac{1}{2} \right) (48 - \pi r - 2r)$$


$$\frac{1}{2} \pi r^2 + r(48 - \pi r - 2r)$$

$$\frac{1}{2} \pi r^2 + 48r - \pi r^2 - 2r^2$$

$$\left(\frac{1}{2} \pi - \pi - 2 \right) r^2 + 48r$$

$$\left(-\frac{1}{2} \pi - 2 \right) r^2 + 48r$$

Graph the function A to find that its maximum occurs when r 6.72 feet.



Xmin = 0, Xmax = 14, Xscl = 1

Ymin = 50, Ymax = 200, Yscl = 50

$$= \frac{1}{2} (48 - \pi r - 2r)$$

$$\frac{1}{2} (48 - \pi(6.72) - 2(6.72))$$

6.72 feet

Hence the optimal window has its semicircular radius equal to its height.

Note: Using calculus it can be shown that the exact

$$\frac{48}{\pi}$$

value of $r = h = \frac{48}{\pi} + 4$.

$$y = a(x - h)^2 + k$$

$$y = a(x - 0)^2 + 6$$

$$y = ax^2 + 6$$

$$500 = a(2100)^2 + 6$$

$$494 = a(2100)^2$$

$$\frac{494}{2100^2} = a$$

Prepare for Section 2.5

P1. $f(x) = x^2 + 4x - 6$

$$\frac{b}{2a} = \frac{-4}{2(1)} = -2$$

$$-2x = -2$$

4

P2. $f(3) = \frac{3(3)}{4} = \frac{243}{10} = 24.3$

$$(3)^2 + 1 = 10$$

$$f(-3) = \frac{3(-3)}{4} = \frac{243}{10} = 24.3$$

$$(-3)^2 + 1 = 10$$

$$f(3) = f(-3)$$

P3. $f(-2) = 2(-2)^3 - 5(-2) = -16 + 10 = -6$

$$f(2) = -[2(2)^3 - 5(2)] = -[16 - 10] = -6 = -f(-2)$$

$$(-2) = -f(2)$$

P4. $f(-2) - g(-2) = (-2)^2 - [-2 + 3] = 4 - 1 = 3$

$$(-1) - g(-1) = (-1)^2 - [-1 + 3] = 1 - 2 = -1 = -f(-2)$$

$$(0) - g(0) = (0)^2 - [0 + 3] = 0 - 3 = -3$$

$$f(1) - g(1) = (1)^2 - [1 + 3] = 1 - 4 = -3 = -f(1)$$

$$(2) - g(2) = (2)^2 - [2 + 3] = 4 - 5 = -1 = -f(2)$$

P5. $\frac{-a+a}{2} = 0, \frac{b+b}{2} = b$


midpoint is $(0, b)$

P6. $\frac{-a+a}{2} = 0, \frac{-b+b}{2} = 0$


midpoint is $(0, 0)$

Section 2.5 Exercises

Plot the points.



Plot the points.




$$0.000112018 \gg a$$


$$= 0.000112018x^2 + 6$$

Chapter 2 Functions and Graphs 3.


Plot the points.




Plot the points.



Plot the points.



Plot the points.



Determine how the graph is symmetric.

The graph is symmetric with respect to the origin.

Determine how the graph is symmetric.

The graph is symmetric with respect to the y -axis.


Determine how the graph is symmetric.

The graph is symmetric with respect to the x -axis, the y -axis, and the origin.


Determine how the graph is symmetric.

The graph is symmetric with respect to the x -axis.


Sketch the graph symmetric to the x -axis.




Sketch the graph symmetric to the x -axis.




Sketch the graph symmetric to the y -axis.




Sketch the graph symmetric to the y -axis.



Sketch the graph symmetric to the origin.



Sketch the graph symmetric to the origin.



Determine if the graph is symmetric.

No

Yes

Determine if the graph is symmetric.

Yes

No

Determine if the graph is symmetric.

No

No

Determine if the graph is symmetric.

No

No

Determine if the graph is symmetric.

Yes

Yes

Determine if the graph is symmetric.

Yes

Yes

Determine if the graph is symmetric.

Yes

Yes

Determine if the graph is symmetric.

No

No

Determine if the graph is symmetric.

Yes

Yes

Determine if the graph is symmetric to the origin.

Not symmetric with respect to the origin since $(-y) = (-x) + 1$ does not simplify to the original equation $y = x + 1$.

Determine if the graph is symmetric to the origin.

No, since $(-y) = 3(-x) - 2$ simplifies to

$(y) = 3x - 2$, which is not equivalent to the original equation $y = 3x - 2$.

Determine if the graph is symmetric to the origin.

Yes, since $(-y) = (-x)^3 - (-x)$ simplifies to

$y = -x^3 + x$, which is equivalent to the original equation $y = x^3 - x$.

Determine if the graph is symmetric to the origin.

3

Yes, since $(-y) = -(-x)$ implies

$y = x^3$ or $y = -x^3$, which is the original equation.

Determine if the graph is symmetric to the origin.

Yes, since $(-y) = \frac{9}{(-x)}$ is equivalent to the original

Determine if the graph is symmetric to the origin.

Yes, since $(-x)^2 + (-y)^2 = 10$ simplifies to the original equation.

Determine if the graph is symmetric to the origin.

))

Yes, since $(-x)^2 - (-y)^2 = 4$ simplifies to the original equation.

Determine if the graph is symmetric to the

origin. Yes, since $\sqrt{-y} = -x$ simplifies to the original equation.

Determine if the graph is symmetric to the

origin. Yes, since $|y| = -x$ simplifies to the original equation.

Determine if the function is odd, even or neither.

Even since $g(-x) = (-x)^2 - 7 = x^2 - 7 = g(x)$.

Determine if the function is odd, even or neither.

Even, since $h(-x) = (-x)^2 + 1 = x^2 + 1 = h(x)$.

Determine if the function is odd, even or

neither. Odd, since $F(-x) = (-x)^5 + (-x)^3 - x^5 - x^3 = -F(x)$.

Determine if the function is odd, even or neither.

Neither, since $G(-x) \neq G(x)$ and $G(-x) \neq -G(x)$.

Determine if the function is odd, even or neither. Even

Determine if the function is odd, even or neither. Even

Determine if the function is odd, even or neither. Even

equation $y = \frac{9}{x}$.

Determine if the function is odd, even
or neither. Neither

Determine if the function is odd, even
or neither. Even

Determine if the function is odd, even
or neither. Even

Determine if the function is odd, even or neither.

Even

Determine if the function is odd, even or neither.

Neither


Determine if the function is odd, even or neither.

Neither


Determine if the function is odd, even or neither.

Odd

Sketch the graphs.




Sketch the graphs.




Sketch the graphs.

$f(x + 2)$




$f(x) + 2$




Sketch the graphs.

$g(x - 1)$




$g(x) - 1$




Sketch the graphs.

$y = f(x - 2) + 1$




$y = f(x + 3) - 2$




Sketch the graphs.

$y = f(x + 3) + 2$



$y = f(x - 2) - 1$



a. Give three points on the graph.

$(x + 3)$

$(-2-3, 5) = (-5, 5)$

$(0-3, -2) = (-3, -2)$

$(1-3, 0) = (-2, 0)$

Give three points on the graph.

$(x) + 1$
 $(-2, 5+1) = (-2, 6)$
 $(0, -2+1) = (0, -1)$
 $(1, 0+1) = (1, 1)$

a. Give three points on the graph.

$g(x - 2)$
 $(-3+2, -1) = (-1, -1)$
 $(1+2, -3) = (3, -3)$
 $(4+2, 2) = (6, 2)$

Give three points on the graph.

$g(x) - 2$
 $(-3, -1-2) = (-3, -3)$
 $(1, -3-2) = (1, -5)$
 $(4, 2-2) = (4, 0)$

a. Give two points on the graph.

$(-x)$
 $(-1, 3) = (1, 3)$
 $(-2, -4)$

Give two points on the graph.

$f(x)$
 $(-1, -3)$
 $(2, -4) = (2, 4)$

a. Give two points on the graph.


$-g(x)$
 $(4, -5) = (4, 5)$
 $(-3, -2)$

Give two points on the graph.


$(-x)$
 $(-4, -5)$
 $(-3, 2) = (3, 2)$

Sketch the graphs.

a. $f(x)$




$f(x)$




Sketch the graphs.


$g(x)$




$g(x)$



Sketch the graphs.




Sketch the graphs.




Sketch the graphs.

$2g(x)$




Chapter 2 Functions and Graphs **b.**

$\frac{1}{2}g(x)$




Sketch the graphs.

$2f(x)$




$\frac{1}{2}f(x)$




Sketch the graphs.

$3h(x)$




$\frac{1}{2}h(x)$




Sketch the graphs.

$4f(x)$




$\frac{1}{4}f(x)$




Sketch the graphs.

$f(2x)$




$f\left(\frac{1}{3}x\right)$




Sketch the graphs.

$g(2x)$




$$g\left(\frac{1}{2}x\right)$$




Sketch the graphs.

$$h(2x)$$




$$h\left(\frac{1}{2}x\right)$$




Sketch the graphs.


$$j(2x)$$




$$j\left(\frac{1}{3}x\right)$$



Sketch the graph.




Sketch the graph.




Sketch the graphs.

$$\frac{1}{2}$$

a. $y = -\frac{1}{2}j(x) + 1$




$y = 2j(x) - 1$




Sketch the graphs.

a. $y = \frac{1}{2}h(x) - 1$




Chapter 2 Functions and Graphs **b. y**


$= -2h(x) + 1$




Graph using a graphing utility.




Graph using a graphing utility.




Graph using a graphing utility.




Graph using a graphing utility.




Reflect the graph about the y-axis and then about the origin.




Reflect the graph about the origin and then about the y-axis.



Reflect the graph about the y-axis and then about the x-axis.



Reflect the graph about the x-axis and then about the origin.



Prepare for Section 2.6

P1. $(2x^2 + 3x - 4) - (x^2 + 3x - 5) = x^2 + 1$

P2. $(3x^2 - x + 2)(2x - 3) = 6x^3 - 2x^2 + 4x - 9x^2 + 3x - 6$
 $6x^3 - 11x^2 + 7x - 6$

P3. $f(3a) = 2(3a)^2 - 5(3a) + 2$

$18a^2 - 15a + 2$

P4. $f(2 + h) = 2(2 + h)^2 - 5(2 + h) + 2$

$2h^2 + 8h + 8 - 5h - 10 + 2$

$2h^2 + 3h$

P5. Domain: all real numbers except $x = 1$

P6. $2x - 8 = 0$

$x = 4$

Domain: $x \geq 4$ or $[4, \infty)$

Section 2.6 Exercises

Evaluate.

$$(f + g)(-2) = f(-2) + g(-2)$$

$$\begin{array}{r} 3 + (-6) \\ -3 \end{array}$$

Evaluate.

$$(f - g)(-2) = f(-2) - g(-2)$$

$$\begin{array}{r} 3 - (-6) \\ 9 \end{array}$$

Evaluate.

$$(f \cdot g)(-2) = f(-2) \cdot g(-2)$$

$$\begin{array}{r} 3(-6) \\ -18 \end{array}$$

Evaluate.

$$\frac{f(-2)}{g(-2)} = \frac{f(-2)}{g(-2)}$$

$$\begin{array}{r} 3 \\ -6 \\ -\frac{1}{2} \end{array}$$

Evaluate.

$$g[f(-5)] = g[7] = -2$$

Evaluate.

$$(4) = f[g(0)] = 3$$

Simplify.

$$(2 + h) = 3(2 + h) - 4$$

$$\begin{array}{r} 6 + 3h - 4 \\ 2 + 3h \end{array}$$

Simplify.

$$(2 + h) = (2 + h)^2 + 1$$

$$\begin{array}{r} 4 + 4h + h^2 + 1 \\ 5 + 4h + h^2 \end{array}$$

Perform the operations and find the domain.

$$(x) + g(x) = (x^2 - 2x - 15) + (x + 3)$$

$$x^2 - x - 12 \text{ Domain all real numbers}$$

$$(x) - g(x) = (x^2 - 2x - 15) - (x + 3)$$

$$x^2 - 3x - 18 \text{ Domain all real numbers}$$

$$f(x)g(x) = (x^2 - 2x - 15)(x + 3)$$

$$= x^3 + x^2 - 21x - 45$$

$$\text{Domain all real numbers}$$

$$(x) / g(x) = (x^2 - 2x - 15) / (x + 3)$$

$$x - 5 \text{ Domain } \{x \mid x \neq -3\}$$

Perform the operations and find the domain.

$$(x) + g(x) = (x^2 - 25) + (x - 5)$$

$$x^2 + x - 30 \text{ Domain all real numbers}$$

$$(x) - g(x) = (x^2 - 25) - (x - 5)$$

$$x^2 - x - 20 \text{ Domain all real numbers}$$

$$(x)g(x) = (x^2 - 25)(x - 5)$$

$$x^3 - 5x^2 - 25x + 125$$

Domain all real numbers

$$(x) / g(x) = (x^2 - 25) / (x - 5)$$

$$x + 5 \text{ Domain } \{x \mid x \neq 5\}$$

Perform the operations and find the domain.

$$(x) + g(x) = (2x + 8) + (x + 4)$$

$$3x + 12 \text{ Domain all real numbers}$$

$$(x) - g(x) = (2x + 8) - (x + 4)$$

$$x + 4 \text{ Domain all real numbers}$$

$$(x)g(x) = (2x + 8)(x + 4)$$

$$2x^2 + 16x + 32 \text{ Domain all real numbers}$$

$$(x) / g(x) = (2x + 8) / (x + 4)$$

$$[2(x + 4)] / (x + 4)$$

$$2 \text{ Domain } \{x \mid x \neq -4\}$$

Perform the operations and find the domain.

$$(x) + g(x) = (5x - 15) + (x - 3)$$

$$6x - 18 \text{ Domain all real numbers}$$

$$(x) - g(x) = (5x - 15) - (x - 3)$$

$$4x - 12 \text{ Domain all real numbers}$$

$$(x)g(x) = (5x - 15)(x - 3)$$

$$2$$

$$5x^2 - 30x + 45 \text{ Domain all real numbers}$$

$$(x) / g(x) = (5x - 15) / (x - 3)$$

$$[5(x - 3)] / (x - 3)$$

$$5 \text{ Domain } \{x \mid x \neq 3\}$$

Perform the operations and find the domain. f

$$(x) + g(x) = (x^3 - 2x^2 + 7x) + x$$

$$x^3 - 2x^2 + 8x \text{ Domain all real numbers}$$

$$(x) - g(x) = (x^3 - 2x^2 + 7x) - x$$

$$x^3 - 2x^2 + 6x \text{ Domain all real}$$

$$\text{numbers } f(x)g(x) = (x^3 - 2x^2 + 7x)x$$

$$x^4 - 2x^3 + 7x^2 \text{ Domain all real numbers}$$

$$(x) / g(x) = (x^3 - 2x^2 + 7x) / x$$

$$x^2 - 2x + 7 \text{ Domain } \{x \mid x \neq 0\}$$

Perform the operations and find the domain.

$$(x) + g(x) = (x^2 - 5x - 8) + (-x)$$

$$x^2 - 6x - 8 \text{ Domain all real numbers}$$

$$(x) - g(x) = (x^2 - 5x - 8) - (-x)$$

$$x^2 - 4x - 8 \text{ Domain all real numbers}$$

$$(x)g(x) = (x^2 - 5x - 8)(-x)$$

$$-x^3 + 5x^2 + 8x \text{ Domain all real numbers}$$

$$(x) / g(x) = (x^2 - 5x - 8x) / (-x)$$

$$= -x + 5 + \frac{8}{x} \text{ Domain } \{x \mid x \neq 0\}$$

Perform the operations and find the domain. f

$$(x) + g(x) = (4x - 7) + (2x^2 + 3x - 5)$$

$$2x^2 + 7x - 12 \text{ Domain all real numbers}$$

$$(x) - g(x) = (4x - 7) - (2x^2 + 3x - 5)$$

$$-2x^2 + x - 2 \text{ Domain all real numbers}$$

$$(x)g(x) = (4x - 7)(2x^2 + 3x - 5)$$

$$8x^3 - 14x^2 + 12x^2 - 20x - 21x + 35$$

$$8x^3 - 2x^2 - 41x + 35$$

$$\text{Domain all real numbers}$$

$$(x) / g(x) = (4x - 7) / (2x^2 + 3x - 5)$$

$$= \frac{4x - 7}{2x^2 + 3x - 5}$$

$$\text{Domain } \left\{x \mid x \neq 1, x \neq \frac{5}{2}\right\}$$

Perform the operations and find the domain. f

$$(x) + g(x) = (6x + 10) + (3x^2 + x - 10)$$

$$3x^2 + 7x \text{ Domain all real numbers}$$

$$(x) - g(x) = (6x + 10) - (3x^2 + x - 10)$$

$$-3x^2 + 5x + 20 \text{ Domain}$$

$$\text{all real numbers}$$

$$(x)g(x) = (6x + 10)(3x^2 + x - 10)$$

$$18x^3 + 6x^2 - 60x + 30x^2 + 10x - 100$$

$$18x^3 + 36x^2 - 50x - 100$$

Domain all real numbers

$$(x) / g(x) = (6x + 10) / (3x^2 + x - 10)$$

$$\frac{6x + 10}{3x^2 + x - 10}$$

$$\text{Domain } \left\{x \mid x \neq -2, x \neq \frac{5}{3}\right\}$$

Perform the operations and find the domain.

$$f(x) + g(x) = \sqrt{x - 3} + x \quad \text{Domain } \{x \mid x \geq 3\}$$

$$f(x) - g(x) = \sqrt{x - 3} - x \quad \text{Domain } \{x \mid x \geq 3\}$$

$$f(x)g(x) = x\sqrt{x - 3} \quad \text{Domain } \{x \mid x \geq 3\}$$

$$f(x) / g(x) = \frac{\sqrt{x - 3}}{x} \quad \text{Domain } \{x \mid x \geq 3\}$$

Perform the operations and find the domain.

$$f(x) + g(x) = \sqrt{x - 4} - x \quad \text{Domain } \{x \mid x \geq 4\}$$

$$f(x) - g(x) = \sqrt{x - 4} + x \quad \text{Domain } \{x \mid x \geq 4\}$$

$$f(x)g(x) = -x\sqrt{x - 4} \quad \text{Domain } \{x \mid x \geq 4\}$$

$$f(x) / g(x) = -\frac{\sqrt{x - 4}}{x} \quad \text{Domain } \{x \mid x \geq 4\}$$

Perform the operations and find the domain.

$$f(x) + g(x) = \sqrt{4 - x^2} + 2 + x$$

$$\text{Domain } \{x \mid -2 \leq x \leq 2\}$$

$$(x) - g(x) = \sqrt{4 - x^2} - 2 - x$$

$$\text{Domain } \{x \mid -2 \leq x \leq 2\}$$

$$f(x)g(x) = (\sqrt{4 - x^2})(2 + x)$$

Domain $\{x \mid -2 \leq x \leq 2\}$

$$f(x) / g(x) = \frac{\sqrt{4-x^2}}{2+x} \text{ Domain } \{x \mid -2 < x \leq 2\}$$

Perform the operations and find the domain.

$$f(x) + g(x) = \sqrt{x^2 - 9} + x - 3$$

Domain $\{x \mid x \leq -3 \text{ or } x \geq 3\}$

$$f(x) - g(x) = \sqrt{x^2 - 9} - x + 3$$

Domain $\{x \mid x \leq -3 \text{ or } x \geq 3\}$

$$(x)g(x) = (\sqrt{x^2 - 9})(x - 3)$$

Domain $\{x \mid x \leq -3 \text{ or } x \geq 3\}$

$$f(x) / g(x) = \frac{\sqrt{x^2 - 9}}{x - 3} \text{ Domain } \{x \mid x \leq -3 \text{ or } x > 3\}$$

21. Evaluate the function.

$$(f + g)(x) = x^2 - x - 2$$

$$(f + g)(5) = (5)^2 - (5) - 2$$

$$25 - 5 - 2$$

$$18$$

Evaluate the function.

$$(f + g)(x) = x^2 - x - 2$$

$$(f + g)(-7) = (-7)^2 - (-7) - 2$$

$$49 + 7 - 2$$

$$54$$

Evaluate the function.

$$(f + g)(x) = x^2 - x - 2$$

$$(f + g)\left(\frac{1}{2}\right) = \left(\frac{1}{2}\right)^2 - \left(\frac{1}{2}\right) - 2$$

$$\frac{1}{4} - \frac{1}{2} - 2$$

$$\underline{9}$$

$$- 4$$

Evaluate the function.

$$(f + g)(x) = x^2 - x - 2$$

$$(f + g)\left(\frac{2}{3}\right) = \left(\frac{2}{3}\right)^2 - \left(\frac{2}{3}\right) - 2$$

$$9\frac{4}{9} - \frac{2}{3} - 2$$

$$-\frac{20}{9}$$

Evaluate the function.

$$(f - g)(x) = x^2 - 5x + 6$$

$$(f - g)(-3) = (-3)^2 - 5(-3) + 6$$

$$9 + 15 + 6$$

$$30$$

Evaluate the function.

$$(f - g)(x) = x^2 - 5x + 6$$

$$(f - g)(24) = (24)^2 - 5(24) + 6$$

$$576 - 120 + 6$$

$$462$$

Evaluate the function.

$$2$$

$$(f - g)(x) = x^2 - 5x + 6$$

$$(f - g)(-1) = (-1)^2 - 5(-1) + 6$$

$$1 + 5 + 6$$

$$12$$

Evaluate the function.

$$2$$

$$(f - g)(x) = x^2 - 5x + 6$$

$$(f - g)(0) = (0)^2 - 5(0) + 6$$

$$6$$

Evaluate the function.

$$(fg)(x) = (x^2 - 3x + 2)(2x - 4)$$

$$2x^3 - 6x^2 + 4x - 4x^2 + 12x - 8$$

$$2x^3 - 10x^2 + 16x - 8$$

$$(fg)(7) = 2(7)^3 - 10(7)^2 + 16(7) - 8$$

$$686 - 490 + 112 - 8$$

$$300$$

Chapter 2 Functions and Graphs

Evaluate the function.

$$(fg)(x) = 2x^3 - 10x^2 + 16x - 8$$

$$(fg)(-3) = 2(-3)^3 - 10(-3)^2 + 16(-3) - 8$$

$$= -54 - 90 - 48 - 8$$

$$= -200$$

Evaluate the function.

$$(fg)(x) = 2x^3 - 10x^2 + 16x - 8$$

$$(fg)\left(\frac{2}{5}\right) = 2\left(\frac{2}{5}\right)^3 - 10\left(\frac{2}{5}\right)^2 + 16\left(\frac{2}{5}\right) - 8$$

$$= \frac{16}{125} - \frac{40}{25} + \frac{32}{5} - 8$$

$$= -125 \frac{384}{125} = -3.072$$

Evaluate the function.

$$(fg)(x) = 2x^3 - 10x^2 + 16x - 8$$

$$(fg)(-100) = 2(-100)^3 - 10(-100)^2 + 16(-100) - 8$$

$$= -2,000,000 - 100,000 - 1600 - 8$$

$$= -2,101,608$$

Evaluate the function.

$$\frac{f(x) - f(x-h)}{h} = \frac{x^2 - 3x + 4 - (x-h)^2 - 3(x-h) + 4}{h}$$

$$= \frac{x^2 - 3x + 4 - (x^2 - 2xh + h^2) - 3x + 3h + 4}{h}$$

$$= \frac{-2xh + 3h - h^2}{h}$$

$$= -2 - 2 \text{ or } -2 \frac{5}{2}$$

Evaluate the function.

$$\frac{f(x) - f(x-h)}{h} = \frac{1}{2}x - \frac{1}{2}$$

$$= \frac{1}{2} - \frac{1}{2}$$

$$= (1) -$$

Evaluate the function.

$$\frac{f(x) - f(x-h)}{h} = \frac{1}{2}x - \frac{1}{2}$$

$$= \frac{1}{2} - \frac{1}{2}$$

$$= \frac{1}{4}$$

Evaluate the function.

$$\frac{f(x) - f(x-h)}{h} = \frac{1}{2}x - \frac{1}{2}$$

$$= \frac{1}{2} - \frac{1}{2}$$

$$= \frac{1}{8} - \frac{1}{2}$$

$$= -\frac{3}{8}$$

Find the difference quotient.

$$\frac{(x+h) - f(x)}{hh} = \frac{[2(x+h) + 4] - (2x + 4)}{hh}$$

$$= \frac{2x + 2(h) + 4 - 2x - 4}{h}$$

$$= \frac{2h}{h}$$

$$= 2$$

Find the difference quotient.

$$\frac{(x+h) - f(x)}{hh} = \frac{[4(x+h) - 5] - (4x - 5)}{hh}$$

$$= \frac{4x + 4(h) - 5 - 4x + 5}{h}$$

$$= \frac{4(h)}{h}$$

$$= 4$$

Find the difference quotient.

$$\frac{(x+h) - f(x)}{h} = \frac{(x+h)^2 - 6 - (x^2 - 6)}{h}$$

$$= \frac{2xh + h^2}{h}$$

$$\begin{aligned}
 \text{è } g \text{ ø} & \quad 2 \quad 2 \\
 & = \underline{11} - \underline{1} \\
 & \quad 2 \quad 2 \\
 & = \underline{10} = 5 \\
 & \quad 2
 \end{aligned}$$

$$\begin{aligned}
 hh & \\
 & \quad 2 \quad 2 \quad 2 \\
 & = \frac{x^2 + 2x(h) + (h)^2 - 6x + 6}{6h} \\
 & = \frac{2x(h) + h^2}{h} \\
 & \quad 2x + h
 \end{aligned}$$

Find the difference quotient.

$$\begin{aligned} \frac{(x+h)-f(x)}{hh} &= \frac{x^2 + 2xh + (h)^2 + 11 - (x^2 + 11)}{h} \\ &= \frac{2xh + h^2}{h} \\ &= 2x + h \end{aligned}$$

Find the difference quotient.

$$\begin{aligned} \frac{f(x+h)-f(x)}{h} &= \frac{2(x+h)^2 + 4(x+h) - 3 - (2x^2 + 4x - 3)}{h} \\ &= \frac{2x^2 + 4xh + 2h^2 + 4x + 4h - 3 - 2x^2 - 4x + 3}{h} \\ &= \frac{4xh + 2h^2 + 4h}{h} \\ &= 4x + 2h + 4 \end{aligned}$$

Find the difference quotient.

$$\begin{aligned} \frac{f(x+h)-f(x)}{h} &= \frac{2(x+h)^2 - 5(x+h) + 7 - (2x^2 - 5x + 7)}{h} \\ &= \frac{2x^2 + 4xh + 2h^2 - 5x - 5h + 7 - 2x^2 + 5x - 7}{h} \\ &= \frac{4xh + 2h^2 - 5h}{h} \\ &= 4x + 2h - 5 \end{aligned}$$

Find the difference quotient.

$$\begin{aligned} \frac{(x+h)-f(x)}{hh} &= \frac{-4(x+h)^2 + 6 - (-4x^2 + 6)}{h} \\ &= \frac{-4x^2 - 8xh - 4h^2 + 6 + 4x^2 - 6}{h} \\ &= \frac{-8xh - 4h^2}{h} \\ &= -8x - 4h \end{aligned}$$

Find the difference quotient.

$$\begin{aligned} f(x) &= -5x^2 - 4x \\ \frac{f(x+h)-f(x)}{h} &= \frac{-5(x+h)^2 - 4(x+h) - (-5x^2 - 4x)}{h} \\ &= \frac{-5x^2 - 10x(h) - 5h^2 - 4x - 4h + 5x^2 + 4x}{h} \\ &= \frac{-10x(h) - 5h^2 - 4h}{h} \\ &= -10x - 5h - 4 \end{aligned}$$

a. On $[0, 1]$, $a = 0$

$$\begin{aligned} t &= 1 - 0 = 1 \\ C(a+t) &= C(1) = 99.8 \text{ (mg/L)/h} \\ C(a) &= C(0) = 0 \\ \text{Average rate of change} &= \frac{C(1)-C(0)}{1-0} = 99.8 - 0 = 99.8 \end{aligned}$$

This is identical to the slope of the line through $(0, C(0))$ and $(1, C(1))$ since

$$\frac{C(1)-C(0)}{1-0} = C(1)-C(0)$$

On $[0, 0.5]$, $a = 0$, $t = 0.5$

$$\begin{aligned} \text{Average rate of change} &= \frac{C(0.5)-C(0)}{0.5-0} = \frac{78.1-0}{0.5} = 156.2 \text{ (mg/L)/h} \end{aligned}$$

On $[1, 2]$, $a = 1$, $t = 2 - 1 = 1$

$$\begin{aligned} \text{Average rate of change} &= \frac{C(2)-C(1)}{2-1} = \frac{50.1-99.8}{1} = -49.7 \text{ (mg/L)/h} \end{aligned}$$

On $[1, 1.5]$, $a = 1$, $t = 1.5 - 1 = 0.5$

$$\begin{aligned} \text{Average rate of change} &= \frac{C(1.5)-C(1)}{0.5-0.5} = \frac{84.4-99.8}{0.5} = -30.8 \text{ (mg/L)/h} \end{aligned}$$

On $[1, 1.25]$, $a = 1$, $t = 1.25 - 1 = 0.25$

$$\begin{aligned} \text{Average rate of change} &= \frac{C(1.25)-C(1)}{0.25-0.25} = \frac{95.7-99.8}{0.25} = -16.4 \text{ (mg/L)/h} \end{aligned}$$

Chapter 2 Functions and Graphs f.

On $[1, 1+t]$,

$$\text{Con}(1+t)$$

$$25(1+t)^3 - 150(1+t)^2 + 225(1+t)$$

$$25(1+3t+3t^2+1(t)^3) - 150(1+2(t)+(t)^2)$$

$$225(1+t)$$

$$25+75(t)+75(t)^2+25(t)^3$$

$$-150-300(t)-150(t)^2+225+225(t)$$

$$100-75(t)^2+25(t)^3$$

$$\text{Con}(1) = 100$$

Average rate of change

$$\frac{\text{Con}(1+t)-\text{Con}(1)}{t}$$

$$\frac{100-75(t)^2+25(t)^3-100}{t}$$

$$\frac{-75(t)^2+25(t)^3}{t}$$

$$-75(t)+25(t)^2$$

As t approaches 0, the average rate of change over

$[1, 1+t]$ seems to approach 0 (mg/L)/h.

a. On $[2, 3]$, $a = 2$

$$t = 3 - 2 = 1$$

$$s(a+t) = s(3) = 6 \cdot 3^2 = 54$$

$$s(a) = s(2) = 6 \cdot 2^2 = 24$$

$$\text{Average velocity} = \frac{s(a+t)-s(a)}{t} = \frac{s(3)-s(2)}{1}$$

$$54 - 24 = 30 \text{ feet per second}$$

This is identical to the slope of the line

through $(2, s(2))$ and $(3, s(3))$ since

$$= \frac{s(3)-s(2)}{3-2} = s(3)-s(2).$$

On $[2, 2.5]$, $a = 2$,

$$t = 2.5 - 2 = 0.5$$

$$s(a+t) = s(2.5) = 6(2.5)^2 = 37.5$$

$$\text{Average velocity} = \frac{s(2.5)-s(2)}{0.5} = \frac{37.5-24}{0.5}$$

$$= \frac{13.5}{0.5} = 27 \text{ feet per second}$$

On $[2, 2.1]$, $a = 2$

$$t = 2.1 - 2 = 0.1$$

$$s(a+t) = s(2.1) = 6(2.1)^2 = 26.46$$

$$\text{Average velocity} = \frac{s(2.1)-s(2)}{0.1} = \frac{26.46-24}{0.1}$$

$$\frac{2.46}{0.1} = 24.6 \text{ feet per}$$

second d. On $[2, 2.01]$, $a = 2$

$$t = 2.01 - 2 = 0.01$$

$$s(a+t) = s(2.01) = 6(2.01)^2 = 24.2406$$

$$\text{Average velocity} = \frac{s(2.01)-s(2)}{0.01} = \frac{24.2406-24}{0.01}$$

$$\frac{0.2406}{0.01} = 24.06 \text{ feet per}$$

second e. On $[2, 2.001]$, $a = 2$

$$t = 2.001 - 2 = 0.001$$

$$s(a+t) = s(2.001) = 6(2.001)^2 = 24.024006$$

$$\text{Average velocity} = \frac{s(2.001)-s(2)}{0.001} = \frac{24.024006-24}{0.001}$$

$$\frac{0.024006}{0.001} = 24.006 \text{ feet per second}$$

f. On $[2, 2+t]$,

$$\text{Con } s(2+t) - s(2) = \frac{6(2+t)^2 - 24}{t}$$

$$\frac{6(4+4(t)+(t)^2) - 24}{t}$$

$$\frac{24+24(t)+6(t)^2-24}{(t)}$$

$$\frac{24t+6(t)^2}{t} = 24+6(t)$$

As t approaches 0, the average velocity seems to approach 24 feet per second.

Find the composite functions.

$$0 \quad (g \circ f)(x) = g[f(x)]$$

$$5 \quad g[3x+5]$$

$$2[3x+5]-7$$

$$6x+10-7$$

$$6x+3$$

$$(f \circ g)(x) = f[g(x)]$$

$$f[2x - 7]$$

$$3[2x - 7] + 5$$

$$6x - 21 + 5$$

$$6x - 16$$

Find the composite functions.

$$\begin{aligned}
 (g \circ f)(x) &= g[f(x)] & (f \circ g)(x) &= f[g(x)] \\
 &= g[2x - 7] & &= f[3x + 2] \\
 &= 3[2x - 7] + 2 & &= 2[3x + 2] - 7 \\
 &= 6x - 21 + 2 & &= 6x + 4 - 7 \\
 &= 6x - 19 & &= 6x - 3
 \end{aligned}$$

Find the composite functions.

$$\begin{aligned}
 (g \circ f)(x) &= g(x^2 + 4x - 1) \\
 &= (x^2 + 4x - 1) + 2 \\
 &= x^2 + 4x + 1
 \end{aligned}$$

$$\begin{aligned}
 (f \circ g)(x) &= f[x + 2] \\
 &= [x + 2]^2 + 4[x + 2] - 1 \\
 &= x^2 + 4x + 4 + 4x + 8 - 1 \\
 &= x^2 + 8x + 11
 \end{aligned}$$

Find the composite functions.

$$\begin{aligned}
 (g \circ f)(x) &= g(2x^2 - 11x + 3) \\
 &= 2(2x^2 - 11x + 3) + 3 \\
 &= 4x^2 - 22x + 6 + 3 \\
 &= 4x^2 - 22x + 9
 \end{aligned}$$

Find the composite functions.

$$\begin{aligned}
 (g \circ f)(x) &= g[f(x)] \\
 &= g(x^3 + 2x) \\
 &= (x^3 + 2x) - 5 \\
 &= x^3 - 10x
 \end{aligned}$$

$$\begin{aligned}
 (f \circ g)(x) &= f[g(x)] \\
 &= f[-5x] \\
 &= [-5x]^3 + 2[-5x] \\
 &= -125x^3 - 10x
 \end{aligned}$$

Find the composite functions.

$$\begin{aligned}
 (g \circ f)(x) &= g[f(x)] \\
 &= g(x^3 - 7) \\
 &= (x^3 - 7) - 6 \\
 &= x^3 - 13
 \end{aligned}$$

$$\begin{aligned}
 (f \circ g)(x) &= f[g(x)] \\
 &= f[x + 1] \\
 &= -(x + 1)^3 - 7 \\
 &= -x^3 - 3x^2 - 3x - 1 - 7 \\
 &= -x^3 - 3x^2 - 3x - 8
 \end{aligned}$$

Find the composite functions.

$$\begin{aligned}
 (g \circ f)(x) &= g\left[\frac{2}{x+1} - 5\right] \\
 &= 3\left(\frac{2}{x+1} - 5\right) + 1 \\
 &= \frac{6 - 5x - 5}{x+1} + 1 \\
 &= \frac{1 - 5x}{x+1} + 1
 \end{aligned}$$

$$\begin{aligned}
 (f \circ g)(x) &= f[g(x)] \\
 &= f\left[\frac{3x-5}{2}\right] \\
 &= \frac{3\left(\frac{3x-5}{2}\right) - 5}{3x-4} + 1
 \end{aligned}$$

Find the composite functions.

$$\begin{aligned}
 (g \circ f)(x) &= g\left[\frac{1}{\sqrt{x+4}}\right] & (f \circ g)(x) &= f\left[\frac{1}{\sqrt{x+4}}\right] \\
 &= \frac{1}{\sqrt{x+4}} - 5 & &= \frac{1}{\sqrt{x+4}} - 5 \\
 &= \frac{1}{\sqrt{x+4}} - 5 & &= \frac{1}{\sqrt{x+4}} - 5 \\
 &= \frac{1}{\sqrt{x+4}} - 5 & &= \frac{1}{\sqrt{x+4}} - 5
 \end{aligned}$$

$$-125x^3 - 10x$$

$$\frac{x+4}{x}$$








$$= \frac{1}{x} + \frac{4}{x}$$

$$= \frac{1+4x}{x}$$

$$\frac{\sqrt{\quad}}{\sqrt{\quad}}$$

$$\frac{\sqrt{\quad} \cdot \sqrt{\quad}}{\sqrt{\quad}}$$

$$= \frac{\sqrt{x+4x^2}}{x}$$



Evaluate the composite function.

$$(f \circ g)(x) = 2x^2 - 10x + 3$$

$$(f \circ g)(-3) = 2(-3)^2 - 10(-3) + 3$$

$$18 + 30 + 3$$

$$51$$

Evaluate the composite function.

$$(g \circ f)(x) = 4x^2 + 2x - 6$$

$$(g \circ f)(-1) = 4(-1)^2 + 2(-1) - 6$$

$$4 - 2 - 6$$

$$-4$$

Evaluate the composite function.

$$(g \circ h)(x) = 9x^4 - 9x^2 - 4$$

$$(g \circ h)(0) = 9(0)^4 - 9(0)^2 - 4$$

$$-4$$

Evaluate the composite function.

$$(h \circ g)(x) = -3x^4 + 30x^3 - 75x^2 + 4$$

$$(h \circ g)(0) = -3(0)^4 + 30(0)^3 - 75(0)^2 + 4$$

$$4$$

Evaluate the composite function.

$$(f \circ f)(x) = 4x + 9$$

$$(f \circ f)(8) = 4(8) + 9$$

$$41$$

Evaluate the composite function.

$$(f \circ f)(x) = 4x + 9$$

$$(f \circ f)(-8) = 4(-8) + 9$$

$$-23$$

Evaluate the composite function.

$$(h \circ g)(x) = -3x^4 + 30x^3 - 75x^2 + 4$$

$$(h \circ g)\left(\frac{2}{5}\right) = -3\left(\frac{2}{5}\right)^4 + 30\left(\frac{2}{5}\right)^3 - 75\left(\frac{2}{5}\right)^2 + 4$$

$$-\frac{48}{625} + 125\frac{240}{25} - \frac{300}{25} + 4$$

$$\frac{-48 + 1200 - 7500 + 2500}{625}$$

$$-\frac{3848}{625}$$

Evaluate the composite function.

$$(g \circ h)(x) = 9x^4 - 9x^2 - 4$$

$$(g \circ h)\left(-\frac{1}{3}\right) = 9\left(-\frac{1}{3}\right)^4 - 9\left(-\frac{1}{3}\right)^2 - 4$$

$$81^9 - \frac{9}{9} - 4$$

$$\frac{1}{9} - 1 - 4$$

$$-4\frac{8}{9} \text{ or } -\frac{44}{9}$$

Evaluate the composite function.

$$(g \circ f)(x) = 4x^2 + 2x - 6$$

$$(g \circ f)(\sqrt{3}) = 4(\sqrt{3})^2 + 2(\sqrt{3}) - 6$$

$$12 + 2\sqrt{3} - 6$$

$$6 + 2\sqrt{3}$$

Evaluate the composite function.

$$(f \circ g)(x) = 2x^2 - 10x + 3$$

$$(f \circ g)(\sqrt{2}) = 2(\sqrt{2})^2 - 10(\sqrt{2}) + 3$$

$$4 - 10\sqrt{2} + 3$$

$$7 - 10\sqrt{2}$$

Evaluate the composite function.

$$(g \circ f)(x) = 4x^2 + 2x - 6$$

$$(g \circ f)(2c) = 4(2c)^2 + 2(2c) - 6$$

$$16c^2 + 4c - 6$$

Evaluate the composite function.

$$2$$

$$(f \circ g)(x) = 2x^2 - 10x + 3$$

$$(f \circ g)(3k) = 2(3k)^2 - 10(3k) + 3$$

$$18k^2 - 30k + 3$$

Evaluate the composite function.

$$(g \circ h)(x) = 9x^4 - 9x^2 - 4$$

$$(g \circ h)(k+1)$$

$$9(k+1)^4 - 9(k+1)^2 - 4$$

$$9(k^4 + 4k^3 + 6k^2 + 4k + 1) - 9k^2 - 18k - 9 - 4$$

$$9k^4 + 36k^3 + 54k^2 + 36k + 9 - 9k^2 - 18k - 13$$

$$9k^4 + 36k^3 + 45k^2 + 18k$$

- 4

Evaluate the composite function.

$$\begin{aligned}
 (h \circ g)(x) &= -3x^4 + 30x^3 - 75x^2 + 4 \\
 (h \circ g)(k-1) &= -3(k-1)^4 + 30(k-1)^3 - 75(k-1)^2 + 4 \\
 &= -3k^4 + 12k^3 - 18k^2 + 12k - 3 \\
 &\quad + 30k^3 - 90k^2 + 90k - 30 - 75k^2 + 150k - 75 + 4 \\
 &= -3k^4 + 42k^3 - 183k^2 + 252k - 104
 \end{aligned}$$

Show $(g \circ f)(x) = (f \circ g)(x)$.

$(g \circ f)(x)$	$(f \circ g)(x)$
$= g[f(x)]$	$= f[g(x)]$
$= g[2x + 3]$	$= f[5x + 12]$
$= 5(2x + 3) + 12$	$= 2(5x + 12) + 3$
$= 10x + 15 + 12$	$= 10x + 24 + 3$
$= 10x + 27$	$= 10x + 27$

$(g \circ f)(x) = (f \circ g)(x)$

Show $(g \circ f)(x) = (f \circ g)(x)$.

$(g \circ f)(x) = g[f(x)]$	$(f \circ g)(x) = f[g(x)]$
$= g[4x - 2]$	$= f[7x - 4]$
$= 7(4x - 2) - 4$	$= 4(7x - 4) - 2$
$= 28x - 14 - 4$	$= 28x - 16 - 2$
$= 28x - 18$	$= 28x - 18$

$(g \circ f)(x) = (f \circ g)(x)$

Show $(g \circ f)(x) = (f \circ g)(x)$.

$(g \circ f)(x)$	$(f \circ g)(x)$
$= g[f(x)]$	$= f[g(x)]$
$= g\left(\frac{6x}{x-1}\right)$	$= f\left(\frac{5x}{x-2}\right)$
$= \frac{5\left(\frac{6x}{x-1}\right)}{\frac{6x}{x-1} - 2}$	$= \frac{6\left(\frac{5x}{x-2}\right)}{\frac{5x}{x-2} - 1}$
$= \frac{6x}{x-1} \cdot \frac{x-1}{30x - 2(x-1)}$	$= \frac{5x}{x-2} \cdot \frac{x-2}{30x - (x-2)}$
$= \frac{6x-2x+2}{4x+2}$	$= \frac{5x-2x+2}{30x-x+2}$

Show $(g \circ f)(x) = (f \circ g)(x)$.

$(g \circ f)(x)$	$(f \circ g)(x)$
$= g[f(x)]$	$= f[g(x)]$
$= g\left(\frac{5x}{x+3}\right)$	$= f\left(\frac{2x}{x-4}\right)$
$= \frac{10x}{5x+3} - 4$	$= \frac{-10x}{-2x+3x-12} = \frac{-10x}{x-12}$
$= \frac{10x}{x+3} - \frac{4(x+3)}{x+3} = \frac{10x - 4x - 12}{x+3} = \frac{6x-12}{x+3}$	$= \frac{-10x}{x-12}$
$= \frac{6(x-2)}{x+3}$	$= \frac{-10x}{x-12}$
$= \frac{6x-12}{x+3}$	$= \frac{-10x}{x-12}$

$(g \circ f)(x) = (f \circ g)(x)$

Show $(g \circ f)(x) = x$ and $(f \circ g)(x) = x$.

$(g \circ f)(x) = g[f(x)]$	$(f \circ g)(x) = f[g(x)]$
$= g[2x + 3]$	$= f\left(\frac{x-3}{2}\right)$
$= \frac{2x+3}{2} - 3$	$= 2\left(\frac{x-3}{2}\right) + 3$
$= \frac{2x}{2}$	$= x - 3 + 3$
$= x$	$= x$

Show $(g \circ f)(x) = x$ and $(f \circ g)(x) = x$.

$(g \circ f)(x) = g[f(x)]$	$(f \circ g)(x) = f[g(x)]$
$= g[4x - 5]$	$= f\left(\frac{x+5}{4}\right)$
$= \frac{4x-5}{4} + 5$	$= 4\left(\frac{x+5}{4}\right) - 5$
$= \frac{4x}{4}$	$= x + 5 - 5$
$= x$	$= x$

$$\begin{aligned}
 &= \frac{30x}{x-1} \cdot \frac{x-1}{2(2x+1)} \\
 &= \frac{15x}{2x+1}
 \end{aligned}
 \qquad
 \begin{aligned}
 &= \frac{30x^{x-2}}{x-2} \cdot \frac{x-2}{2(2x+1)} \\
 &= \frac{15x}{2x+1}
 \end{aligned}$$

$$(g \circ f)(x) = (f \circ g)(x)$$

$$X_{\min} = 0, X_{\max} = 14, X_{\text{scl}} = 2,$$

$$Y_{\min} = -1, Y_{\max} = 6, Y_{\text{scl}} = 2$$

$$r = 1.5t$$

a. Since

$$d^2 + 4^2 = s^2,$$

$$d^2 = s^2 - 16$$

$$d = \sqrt{s^2 - 16}$$

$$d = \sqrt{(48-t)^2 - 16} \quad \text{Substitute } 48-t \text{ for } s$$

$$\sqrt{2304 - 96t + t^2 - 16}$$

$$\sqrt{t^2 - 96t + 2288}$$

b. $s(35) = 48 - 35 = 13$ ft

$$d(35) = \sqrt{35^2 - 96(35) + 2288}$$

$$\sqrt{153} \approx 12.37 \text{ ft}$$

The sides of the triangles are proportional, so we have

$$\frac{22}{16} = \frac{12}{t}$$

Solving for x

$$x = \frac{12(22)}{16t}, \text{ or } \frac{33}{2t}$$

Prepare for Section 2.7

$$\frac{1}{1}$$

P1. Slope: $-\frac{3}{3}$; y-intercept: (0, 4)

P2. $3x - 4y = 12$

$$= 4\frac{3}{4}x - 3$$

$$\frac{3}{4}$$

Slope: $\frac{3}{4}$; y-intercept: (0, -3)

P3. $y = -0.45x + 2.3$

$$\frac{2}{3}$$

P4. $y + 4 = -\frac{3}{3}(x - 3)$

$$= -\frac{2}{3}x - 2$$

P5. $f(2) = 3(2)^2 + 4(2) - 1 = 12 + 8 - 1 = 19$

P6. $|f(x_1) - y_1| + |f(x_2) - y_2|$

$$|(2)^2 - 3(-1)| + |(4)^2 - 3(-14)|$$

$$|4 - 3 + 1| + |16 - 3 - 14|$$

$$2 + 1$$

$$3$$

Section 2.7 Exercises

The scatter diagram suggests no relationship between x and y .

The scatter diagram suggest a nonlinear relationship between x and y .



The scatter diagram suggests a linear relationship between x and y .

The scatter diagram suggests a linear relationship between x and y .

Figure A better approximates a graph that can be modeled by an equation than does Figure B. Thus Figure A has a coefficient of determination closer to 1.



Figure A better approximates a graph that can be modeled by an equation than does Figure B. Thus Figure A has a coefficient of determination closer to 1.

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.



$$y = 2.00862069x + 0.5603448276$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.



$$= 1.918918919x + 0.4594594595$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

EDIT	TESTS	L1	L2	L3	2
1:Edit...		-3	11.8	-----	
2:SortA(-1	9.5		
3:SortD(0	8.6		
4:ClrList		1	8.7		
5:SetUpEditor		2	5.4		

		L2(G) =			

EDIT	TESTS	LinReg
1:1-Var Stats		y=ax+b
2:2-Var Stats		a=-.7231182796
3:Med-Med		b=9.233870968
4:LinReg(ax+b)		r ² =.9218901289
5:QuadReg		r=-.9601510969
6:CubicReg		
7:QuartReg		

$$y = 0.7231182796x + 9.233870968$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

EDIT	TESTS	L1	L2	L3	2
1:Edit...		-7	-11.7	-----	
2:SortA(-5	8.8		
3:SortD(1	8.1		
4:ClrList		1	8.9		
5:SetUpEditor		2	-5.7		

		L2(G) =			

EDIT	TESTS	LinReg
1:1-Var Stats		y=ax+b
2:2-Var Stats		a=.6591216216
3:Med-Med		b=-6.658108108
4:LinReg(ax+b)		r ² =.9759762324
5:QuadReg		r=.9879150937
6:CubicReg		
7:QuartReg		

$$y = 0.6591216216x - 6.658108108$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

EDIT	TESTS	L1	L2	L3	2
1:Edit...		1.3	-4.1	-----	
2:SortA(2.6	-9		
3:SortD(5.4	1.2		
4:ClrList		6.2	7.6		
5:SetUpEditor		7.5	10.5		

		L2(G) =			

EDIT	TESTS	LinReg
1:1-Var Stats		y=ax+b
2:2-Var Stats		a=2.222641509
3:Med-Med		b=-7.364150943
4:LinReg(ax+b)		r ² =.8956132837
5:QuadReg		r=.9463684714
6:CubicReg		
7:QuartReg		

$$y = 2.222641509x - 7.364150943$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

EDIT	TESTS	L1	L2	L3	2
1:Edit...		-1.5	8.1	-----	
2:SortA(-5	6.2		
3:SortD(3	2.3		
4:ClrList		5.4	-7.1		
5:SetUpEditor		6.1	-8.6		

		L2(G) =			

EDIT	TESTS	LinReg
1:1-Var Stats		y=ax+b
2:2-Var Stats		a=2.301587302
3:Med-Med		b=4.813968254
4:LinReg(ax+b)		r ² =.9978517183
5:QuadReg		r=-.9989252816
6:CubicReg		
7:QuartReg		

$$y = 2.301587302x + 4.813968254$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

EDIT	TESTS	L1	L2	L3	2
1:Edit...		1	-1	-----	
2:SortA(1	0		
3:SortD(1	4		
4:ClrList		2	2		
5:SetUpEditor					

		L2(G) =			

EDIT	TESTS	QuadReg
1:1-Var Stats		y=ax ² +bx+c
2:2-Var Stats		a=1.095779221
3:Med-Med		b=-2.696428571
4:LinReg(ax+b)		c=1.136363636
5:QuadReg		R ² =.9943480823
6:CubicReg		
7:QuartReg		

$$y = 1.095779221x^2 - 2.696428571x + 1.136363636$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

EDIT	TESTS	L1	L2	L3	2
1:Edit...		-2	-5	-----	
2:SortA(-1	0		
3:SortD(0	1		
4:ClrList		1	4		
5:SetUpEditor		2	4		

		L2(G) =			

EDIT	TESTS	QuadReg
1:1-Var Stats		y=ax ² +bx+c
2:2-Var Stats		a=-.5714285714
3:Med-Med		b=2.2
4:LinReg(ax+b)		c=1.942857143
5:QuadReg		R ² =.9666319082
6:CubicReg		
7:QuartReg		

$$y = 0.5714285714x^2 + 2.2x + 1.942857143$$

Enter the data on your calculator. The technique for a


TI-83 calculator is illustrated here. Press STAT.

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

L1	L2	L3	3
44	19		
12	19		
15	19		
18	19		
21	19		
24	19		
27	19		
30	19		
33	19		
36	19		
39	19		
42	19		
45	19		
48	19		
51	19		
54	19		
57	19		
60	19		
63	19		
66	19		
69	19		
72	19		
75	19		
78	19		
81	19		
84	19		
87	19		
90	19		
93	19		
96	19		
99	19		
102	19		
105	19		
108	19		
111	19		
114	19		
117	19		
120	19		
123	19		
126	19		
129	19		
132	19		
135	19		
138	19		
141	19		
144	19		
147	19		
150	19		
153	19		
156	19		
159	19		
162	19		
165	19		
168	19		
171	19		
174	19		
177	19		
180	19		
183	19		
186	19		
189	19		
192	19		
195	19		
198	19		
201	19		
204	19		
207	19		
210	19		
213	19		
216	19		
219	19		
222	19		
225	19		
228	19		
231	19		
234	19		
237	19		
240	19		
243	19		
246	19		
249	19		
252	19		
255	19		
258	19		
261	19		
264	19		
267	19		
270	19		
273	19		
276	19		
279	19		
282	19		
285	19		
288	19		
291	19		
294	19		
297	19		
300	19		
303	19		
306	19		
309	19		
312	19		
315	19		
318	19		
321	19		
324	19		
327	19		
330	19		
333	19		
336	19		
339	19		
342	19		
345	19		
348	19		
351	19		
354	19		
357	19		
360	19		
363	19		
366	19		
369	19		
372	19		
375	19		
378	19		
381	19		
384	19		
387	19		
390	19		
393	19		
396	19		
399	19		
402	19		
405	19		
408	19		
411	19		
414	19		
417	19		
420	19		
423	19		
426	19		
429	19		
432	19		
435	19		
438	19		
441	19		
444	19		
447	19		
450	19		
453	19		
456	19		
459	19		
462	19		
465	19		
468	19		
471	19		
474	19		
477	19		
480	19		
483	19		
486	19		
489	19		
492	19		
495	19		
498	19		
501	19		
504	19		
507	19		
510	19		
513	19		
516	19		
519	19		
522	19		
525	19		
528	19		
531	19		
534	19		
537	19		
540	19		
543	19		
546	19		
549	19		
552	19		
555	19		
558	19		
561	19		
564	19		
567	19		
570	19		
573	19		
576	19		
579	19		
582	19		
585	19		
588	19		
591	19		
594	19		
597	19		
600	19		
603	19		
606	19		
609	19		
612	19		
615	19		
618	19		
621	19		
624	19		
627	19		
630	19		
633	19		
636	19		
639	19		
642	19		
645	19		
648	19		
651	19		
654	19		
657	19		
660	19		
663	19		
666	19		
669	19		
672	19		
675	19		
678	19		
681	19		
684	19		
687	19		
690	19		
693	19		
696	19		
699	19		
702	19		
705	19		
708	19		
711	19		
714	19		
717	19		
720	19		
723	19		
726	19		
729	19		
732	19		
735	19		
738	19		
741	19		
744	19		
747	19		
750	19		
753	19		
756	19		
759	19		
762	19		
765	19		
768	19		
771	19		
774	19		
777	19		
780	19		
783	19		
786	19		
789	19		
792	19		
795	19		
798	19		
801	19		
804	19		
807	19		
810	19		
813	19		
816	19		
819	19		
822	19		
825	19		
828	19		
831	19		
834	19		
837	19		
840	19		
843	19		
846	19		
849	19		
852	19		
855	19		
858	19		
861	19		
864	19		
867	19		
870	19		
873	19		
876	19		
879	19		
882	19		
885	19		
888	19		
891	19		
894	19		
897	19		
900	19		
903	19		
906	19		
909	19		
912	19		
915	19		
918	19		
921	19		
924	19		
927	19		
930	19		
933	19		
936	19		
939	19		
942	19		
945	19		
948	19		
951	19		
954	19		
957	19		
960	19		
963	19		
966	19		
969	19		
972	19		
975	19		
978	19		
981	19		
984	19		
987	19		
990	19		
993	19		
996	19		
999	19		

L1	L2	L3	2
44	19		
12	19		
15	19		
18	19		
21	19		
24	19		
27	19		
30	19		
33	19		
36	19		
39	19		
42	19		
45	19		
48	19		
51	19		
54	19		
57	19		
60	19		
63	19		
66	19		
69	19		
72	19		
75	19		
78			


Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.



$$y = 0.1628623408x - 0.6875682232$$

$$y = 0.1628623408(158) - 0.6875682232 \quad 25$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.




$$y = 0.6800298805x + 69.05129482$$

5 feet 8 inches = 68 inches;

$$y = 0.6800298805(68) + 69.05129482 \quad 23$$


Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.



The value of r is close to 0. Therefore, no, there is not


a strong linear relationship between the current and the torque.

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.



Yes, because the linear correlation coefficient is close to -1 .

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.




Yes, there is a strong linear correlation.

$$y = -0.9116x + 79.783$$


$$y = -0.9116(25) + 79.783 \approx 57 \text{ years}$$

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.





Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT.



TI-83 calculator is illustrated here. Press STAT.



$$= 113.3111246x + 21.83605895$$



Positively

$$y = 113.3111246(9.5) + 21.83605895$$

1098 calories

Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT.




$$y = -0.0074642857x^2 + 1.148214286x + 4.807142857$$

$$y = -0.0074642857(65)^2 + 1.148214286(65) + 4.807142857$$


47.9 ft

Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT.





a strong linear relationship between the current and the torque.



$$y = 0.6328671329x^2 + 33.61608329x - 379.4405594$$

Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT.



$$= -0.75x^2 + 10.66x - 17.91$$



For August 2011, $x = 8$,

$$= -0.75(8)^2 + 10.66(8) - 17.91$$

»19.4

Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT.



$$= 198.2272727x^2 + 10,708.60909x + 52,885.98182$$



52,885.98182 For 2006, $x = 1$,

$$= 198.2272727(1)^2 + 10,708.60909(1) + 52,885.98182$$

»63,793 thousand gallons

Enter the data on your calculator. The technique for a



TI-83 calculator is illustrated here. Press STAT.



$$= 107.4664502x^2 - 137.1482684x + 6909.909091$$

Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT.





$$y = 0.0165034965x^2 + 1.366713287x + 5.685314685$$

$$y = 0.0165034965(50)^2 + 1.366713287(50) + 5.685314685$$

32.8 mpg

Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT.





$$y = 0.05208x^2 - 3.56026x + 82.32999$$

$$-2 \frac{b}{a} = -2 \frac{-3.56026}{2(0.05208)} \approx 34 \text{ kilometers per hour}$$

a. Enter the data on your calculator. The technique for



TI-83 calculator is illustrated here. Press STAT.

5-lb ball





$$= 0.6130952381t^2 - 0.0714285714t + 1.071428571$$

10-lb ball



$$= 0.6091269841t^2 - 0.0011904762t - 0.3$$

15-lb ball



$$y = 0.5922619048t^2 + 0.3571428571t - 1.520833333$$

Enter the data on your calculator. The technique for a

b. All the regression equations are approximately the same. Therefore, there is one equation of motion.

Enter the data on your calculator. The technique for a TI-83 calculator is illustrated here. Press STAT.

<pre> EDIT CALC TESTS 1:Edit... 2:SortA(3:SortD(4:ClrList 5:SetUpEditor </pre>	<table border="1"> <tr> <th>L1</th> <th>L2</th> <th>L3</th> <th>3</th> </tr> <tr> <td>.032</td> <td>170</td> <td></td> <td></td> </tr> <tr> <td>.034</td> <td>290</td> <td></td> <td></td> </tr> <tr> <td>.214</td> <td>-130</td> <td></td> <td></td> </tr> <tr> <td>.263</td> <td>-70</td> <td></td> <td></td> </tr> <tr> <td>.275</td> <td>-185</td> <td></td> <td></td> </tr> <tr> <td>.45</td> <td>200</td> <td></td> <td></td> </tr> </table>	L1	L2	L3	3	.032	170			.034	290			.214	-130			.263	-70			.275	-185			.45	200		
L1	L2	L3	3																										
.032	170																												
.034	290																												
.214	-130																												
.263	-70																												
.275	-185																												
.45	200																												

<pre> EDIT CALC TESTS 1:1-Var Stats 2:2-Var Stats 3:Med-Med 4:LinReg(ax+b) 5:QuadReg 6:CubicReg 7:QuartReg </pre>	<pre> LinReg y=ax+b a=454.1584409 b=-40.7836491 r^2=.6235305209 r=.7896394879 </pre>
---	--

$$y = 454.1584409x - 40.78364910$$

$$y = 454.1584409(1.5) - 40.78364910 \text{ 640}$$

kilometers per second

Chapter 2 Review Exercises

Finding the distance. [2.1]

$$= \sqrt{7-(-3))^2 + (11-2)^2}$$

$$= \sqrt{0^2 + 9^2} = 10\sqrt{0+81} = 18\sqrt{\quad}$$

Finding the distance. [2.1]

$$= \sqrt{5-(-3))^2 + (-4-(-8))^2}$$

$$= \sqrt{8^2 + 4^2} = \sqrt{64+16} = \sqrt{80} = 4\sqrt{5}$$


Finding the midpoint: (2, 8), (-3, 12). [2.1]

$$M = \left(\frac{2 + (-3)}{2}, \frac{8 + 12}{2} \right) = \left(\frac{-1}{2}, 10 \right)$$

Finding the midpoint: (-4, 7), (8, -11). [2.1]


$$M = \left(\frac{-4 + 8}{2}, \frac{7 + (-11)}{2} \right) = (2, -2)$$

Graph the equation: $2x - y = -2$. [2.1]




$$y = 0.05208x^2 - 3.56026x + 82.32999$$


Graph the equation: $2x^2 + y = 4$. [2.1]



Graph the equation: $y = x|-2 + |$. [2.1]



Graph the equation: $y = -2x$. [2.1]



Finding x- and y-intercepts and graph: $x = y^2 - 1$ [2.1]

For the y-intercept, let $x = 0$ and solve for y .


$$0 = y^2 - 1$$

$$= 1, \text{ y-intercepts: } (0, -1), (0, 1)$$

For the x-intercept, let $y = 0$ and solve for x .

$$= (0)^2 - 1$$

$$x = -1, \text{ x-intercept: } (-1, 0)$$



Finding x- and y-intercepts and graph: $|x - y| = 4$ [2.1]

For the y-intercept, let $x = 0$ and solve for y .

$$|0 - y| = 4$$


$$= 4, \text{ y-intercepts: } (0, -4), (0, 4)$$

the x-intercept, let $y = 0$ and solve for x .

$$|x - (0)| = 4$$

$x =$

4, x -intercepts: $(-4, 0), (4, 0)$



Finding x - and y -intercepts and graph: [2.1]

$$3x + 4y = 12$$

For the y -intercept, let $x = 0$ and solve for y .


$$3(0) + 4y = 12$$

$$y = 3, \text{ y-intercept: } (0, 3)$$

For the x -intercept, let $y = 0$ and solve for x .

$$3x + 4(0) = 12$$

$$x = 4, \text{ x-intercept: } (4, 0)$$



Finding x - and y -intercepts and graph:

$$= y - 1 + 1 \text{ [2.1]}$$

For the y -intercept, let $x = 0$ and solve for y .


$$0 = y - 1 + 1$$

$-1 = y - 1$, since this statement is false, there is no y -intercept.

For the x -intercept, let $y = 0$ and solve for x .

$$= 0 - 1 + 1$$

$$x = 2, \text{ x-intercept: } (2, 0)$$



Finding the center and radius. [2.1]

$$(x - 3)^2 + (y + 4)^2 = 81$$

center $(3, 4)$, radius 9

Finding the center and radius. [2.1]

$$x^2 + 10x + y^2 + 4y = -20$$

$$x^2 + 10x + 25 + y^2 + 4y + 4 = -20 + 25 + 4$$

$$4(x + 5)^2 + (y + 2)^2 = 9$$

center $(-5, -2)$, radius 3

Finding the equation. [2.1]

Center: $(2, -3)$, radius 5

$$(x - 2)^2 + (y + 3)^2 = 5^2$$

Finding the equation. [2.1]

Center: $(-5, 1)$, passing through $(3, 1)$

$$(x + 5)^2 + (y - 1)^2 = r^2$$

$$(3 + 5)^2 + (1 - 1)^2 = r^2$$

$$8^2 + 0^2 = r^2$$

$$8^2 = r^2$$

$$(x + 5)^2 + (y - 1)^2 = 8^2$$

Is y a function of x ? [2.2]

$$x - y = 4$$

$$y = x - 4, \text{ y is a function of x.}$$

Is y a function of x ? [2.2]

$$x + y^2 = 4$$

$$y^2 = -x + 4$$

$$y = \pm\sqrt{-x + 4}, \text{ y is a not function of x.}$$

Is y a function of x ? [2.2]

$$|x| + |y| = 4$$

$$|y| = -|x| + 4$$

$$y = (-|x| + 4), \text{ y is a not function of x.}$$

Is y a function of x ? [2.2]

$$|x| + y = 4$$

$$y = -|x| + 4, \text{ y is a function of x.}$$

21. Evaluate the function $f(x) = 3x^2 + 4x - 5$, [2.2]

$$f(1) = 3(1)^2 + 4(1) - 5$$

$$3(1) + 4 - 5$$

$$3 + 4 - 5$$

$$2$$

Chapter 2 Functions and Graphs

$$f(-3) = 3(-3)^2 + 4(-3) - 5$$

$$= 3(9) - 12 - 5$$

$$= 27 - 12 - 5$$

$$= 10$$

$$f(t) = 3t^2 + 4t - 5$$

$$f(x+h) = 3(x+h)^2 + 4(x+h) - 5$$

$$= 3(x^2 + 2xh + h^2) + 4x + 4h - 5$$

$$= 3x^2 + 6xh + 3h^2 + 4x + 4h - 5$$

$$3f(t) = 3(3t^2 + 4t - 5)$$

$$= 9t^2 + 12t - 15$$

$$f(3t) = 3(3t)^2 + 4(3t) - 5$$

$$= 3(9t^2) + 12t - 5$$

$$= 27t^2 + 12t - 5$$

22. Evaluate the function $g(x) = \sqrt{64 - x^2}$, [2.2]

$$g(3) = \sqrt{64 - 3^2}$$

$$= \sqrt{64 - 9}$$

$$= \sqrt{55}$$

$$g(-5) = \sqrt{64 - (-5)^2}$$

$$= \sqrt{64 - 25}$$

$$= \sqrt{39}$$

$$g(8) = \sqrt{64 - 8^2}$$

$$= \sqrt{64 - 64}$$

$$= \sqrt{0}$$

$$= 0$$

$$g(-x) = \sqrt{64 - (-x)^2}$$

$$= \sqrt{64 - x^2}$$

$$2g(t) = 2\sqrt{64 - t^2}$$

$$g(2t) = \sqrt{64 - (2t)^2}$$

$$= \sqrt{64 - 4t^2}$$

$$= \sqrt{4(16 - t^2)}$$

$$= 2\sqrt{16 - t^2}$$

Evaluate the function. [2.2]

a. Since $x = 3 \geq 0$, use $f(x) = x^2 - 3$.

$$f(3) = (3)^2 - 3 = 9 - 3 = 6$$

Since $x = -2 < 0$, use $f(x) = 3x + 2$.

$$f(-2) = 3(-2) + 2 = -6 + 2 = -4$$

Since $x = 0 \geq 0$, use $f(x) = x^2 - 3$.

$$f(0) = (0)^2 - 3 = 0 - 3 = -3$$

Evaluate the function. [2.2]

Since $x = 0$ and $-3 \leq x < 5$, use $f(x) = x^2 + 1$. $f(0)$

$$f(0) = (0)^2 + 1 = 1$$

b. Since $x = -3$ and $-3 \leq x < 5$, use $f(x) = x^2 + 1$.

$$f(-3) = (-3)^2 + 1 = 9 + 1 = 10$$

Since $x = 5 \geq 5$, use $f(x) = x - 7$.

$$f(5) = 5 - 7 = -2$$

Find the domain of $f(x) = -2x^2 + 3$. [2.2]

Domain $\{x \mid x \text{ is a real number}\}$

26. Find the domain of $f(x) = \sqrt{6 - x}$. [2.2]

Domain $\{x \mid x \leq 6\}$

27. Find the domain of $f(x) = \sqrt{25 - x^2}$. [2.2]

Domain $\{x \mid -5 \leq x \leq 5\}$

28. Find the domain of $f(x) = \frac{3}{x^2 - 2x - 15}$. [2.2]

Domain $\{x \mid x \neq -3, x \neq 5\}$

Find the values of a in the domain of

$$f(x) = x^2 + 2x - 4 \text{ for which } f(a) = -1. [2.2]$$

$$a^2 + 2a - 4 = -1 \text{ Replace } f(a) \text{ with } a^2 + 2a - 4$$

$$a^2 + 2a - 3 = 0$$

$$(a + 3)(a - 1) = 0$$


$$\begin{array}{ll} a+3=0 & a-1=0 \\ a=-3 & a=1 \end{array}$$

30. Find the values of a in the domain of $f(x) = \frac{4}{x+1}$ for


which $f(a) = 2$. [2.2]

$$\begin{aligned} \frac{4}{a+1} &= 2 \quad \text{Replace } f(a) \text{ with } \frac{4}{a+1} \\ &= 2(a+1) \\ &= 2a+2 \\ &= 2a \\ &= a \end{aligned}$$

Graph $f(x) = |x-1| + 1$ [2.2]



Graph $f(x) = 4 - \sqrt{x}$ [2.2]



33. Find the zeros of f for $f(x) = 2x + 6$. [2.2]

$$\begin{aligned} f(x) &= 0 \\ 2x + 6 &= 0 \\ 2x &= -6 \\ x &= -3 \end{aligned}$$

Find the zeros of f for $f(x) = x^2 - 4x - 12$. [2.2]

$$\begin{aligned} f(x) &= 0 \\ x^2 - 4x - 12 &= 0 \\ (x+2)(x-6) &= 0 \\ x+2 = 0 \quad x-6 = 0 \\ x = -2 \quad x = 6 \end{aligned}$$

Evaluate the function $g(x) = 2x$. [2.2]

a. $g(2) = 2 \cdot 2 = 4$

b. $g\left(\frac{2}{3}\right) = 2 \cdot \frac{2}{3} = \frac{4}{3} \approx 1.333333 = 1.33$

c. $g(-2) = 2(-2) = -4$

36. Evaluate the function $f(x) = 1 - x$. [2.2]

a. $f(\sqrt{2}) = 1 - \sqrt{2} \approx 1 - 1.4142 = -0.4142$

$f(0.5) = 1 - 0.5 = 0.5$

$f(-1) = 1 - (-1) = 2$

Find the slope. [2.3]

$$m = \frac{-1 - 6}{4 - 3} = \frac{-7}{1} = -7$$

$4 + 3 = 7$

Find the slope. [2.3]

$$m = \frac{4 - 2}{-5 - 5} = \frac{2}{-10} = -\frac{1}{5}$$


Find the slope. [2.3]

$$m = \frac{-2 + 2}{-3 - 4} = \frac{0}{-7} = 0$$

Find the slope. [2.3]


$$m = \frac{-1 + 3}{-4 - 6} = \frac{2}{-10} = -\frac{1}{5}$$

Graph $f(x) = -\frac{3}{4}x + 2$. [2.3]




Graph $f(x) = 2 - x$. [2.3]

$m = -1$, y-intercept (0, 2)




43. Graph $3x - 4y = 8$. [2.3]
 $-4y = -3x + 8$
 $y = \frac{3}{4}x - 2$

x-intercept $(\frac{8}{3}, 0)$, y-intercept $(0, -2)$



44. Graph $2x + 3y = 9$. [2.3]
 $3y = -2x + 9$
 $y = -\frac{2}{3}x + 3$

x-intercept $(\frac{9}{2}, 0)$, y-intercept $(0, 3)$



Find the equation. [2.3]

$$y - 2 = -\frac{2}{3}(x + 3)$$

$$y - 2 = -\frac{2}{3}x - 2$$

$$y = -\frac{2}{3}x$$

Find the equation. [2.3]

$$+ 4 = -2(x - 1)$$

$$+ 4 = -2x + 2$$

$$y = -2x - 2$$

Find the equation. [2.3]

$$m = 1 \cdot \frac{6}{-2} = -\frac{3}{1} = -3$$

$$- 6 = 1(x - 1)$$

Find the equation. [2.3]

$$m = \frac{15+6}{8-4} = \frac{21}{4} = \frac{7}{4}$$

$$y - 15 = \frac{7}{4}(x - 8)$$

$$y - 15 = \frac{7}{4}x - 14$$

$$y = \frac{7}{4}x + 1$$

Find the equation. [2.3]

$$y = \frac{2}{3}x - 1 \text{ has slope } m = \frac{2}{3}$$

$$y - y_1 = \frac{2}{3}(x - x_1)$$

$$y - 5 = \frac{2}{3}(x - 7)$$

$$y - 5 = \frac{2}{3}x - \frac{14}{3}$$

$$y + 5 = \frac{2}{3}x - 7$$

$$y = \frac{2}{3}x - 12$$

Find the equation. [2.3]

$$2x - 5y = 2$$

$$-5y = -2x + 2$$

$$y = \frac{2}{5}x - \frac{2}{5} \text{ has slope } m = \frac{2}{5}$$

$$y - y_1 = \frac{2}{5}(x - x_1)$$

$$-(-5) = \frac{2}{5}(x - (-1))$$

$$y + 5 = \frac{2}{5}x + \frac{2}{5}$$

$$y = \frac{2}{5}x - \frac{23}{5}$$

Find the equation. [2.3]

$$y - 2 = -\frac{3}{2}(x - 2) \text{ has perpendicular slope } m = \frac{2}{3}$$

$$y - 2 = \frac{2}{3}(x - 2)$$

$$y - 2 = \frac{2}{3}x - \frac{4}{3}$$

$$y - (-1) = \frac{2}{3}(x - 3)$$

$$y + 1 = \frac{2}{3}x - 2$$

$$y - 6 = x - 1$$

$$= x + 5$$

$$= \frac{2}{3}x - 3$$

Find the equation. [2.3]

$$2x - 5y = 10$$

$$-5y = -2x + 10$$

$$y = \frac{2}{5}x - 2 \text{ has perpendicular slope } m = -\frac{5}{2}$$

$$y - y_1 = -\frac{5}{2}(x - x_1)$$

$$y - 6 = -\frac{5}{2}(x - 2)$$

$$y - 6 = -\frac{5}{2}x + 5$$

$$y = -\frac{5}{2}x + 11$$

Find the function. [2.3]

$$m = \frac{175 - 155}{118 - 106} = \frac{20}{12} = \frac{5}{3}$$

$$(x) - 175 = \frac{5}{3}(x - 118)$$

$$(x) - 175 = \frac{5}{3}x - \frac{590}{3}$$

$$(x) = \frac{5}{3}x - \frac{65}{3}$$

Find the function. [2.3]

$$m = \frac{350 - 122}{10 - 2} = \frac{228}{8} = 28.5$$

$$(t) - 350 = 28.5(t - 10)$$

$$(t) - 350 = 28.5t - 285$$

$$f(t) = 28.5t + 65$$

Write the quadratic equation in standard form. [2.4]

$$(x) = (x^2 + 6x) + 10$$

$$(x) = (x^2 + 6x + 9) + 10 - 9$$

$$(x) = (x + 3)^2 + 1$$

Write the quadratic equation in standard form. [2.4]

$$(x) = (2x^2 + 4x) + 5$$

$$(x) = 2(x^2 + 2x) + 5$$

$$(x) = 2\left(x^2 + 2x + 1\right) + 5 - 2$$

$$(x) = 2(x + 1)^2 + 3$$

Write the quadratic equation in standard form. [2.4]

$$(x) = -x^2 - 8x + 3$$

Write the quadratic equation in standard form. [2.4]

$$f(x) = (4x^2 - 6x) + 1$$

$$f(x) = 4\left(x^2 - 2\frac{3}{4}x\right) + 1$$

$$(x) = 4\left(x^2 - 2\frac{3}{4}x + \frac{9}{16}\right) + 1 - \frac{9}{4}$$

$$(x) = 4\left(x - 4\frac{3}{4}\right)^2 + \frac{4}{4} - \frac{9}{4}$$

$$(x) = 4\left(x - 4\frac{3}{4}\right)^2 - 4\frac{5}{4}$$

Write the quadratic equation in standard form. [2.4]

$$(x) = -3x^2 + 4x - 5$$

$$(x) = -3\left(x^2 - \frac{4}{3}x\right) - 5$$

$$(x) = -3\left(x^2 - \frac{4}{3}x + 9\frac{4}{9}\right) - 5 + \frac{4}{3}$$

$$(x) = -3\left(x - \frac{2}{3}\right)^2 - \frac{11}{3}$$

Write the quadratic equation in standard form. [2.4]

$$(x) = x^2 - 6x + 9$$

$$(x) = (x^2 - 6x) + 9$$

$$(x) = (x^2 - 6x + 9) + 9 - 9$$

$$(x) = (x - 3)^2 + 0$$

Find the vertex. [2.4]

$$\frac{-b}{2a} = \frac{-(-6)}{2(3)} = \frac{6}{6} = 1$$

$$(1) = 3(1)^2 - 6(1) + 11$$

$$3(1) - 6 + 11$$

$$3 - 6 + 11$$

$$8$$

Thus the vertex is (1, 8).

Find the vertex. [2.4]

$$\frac{-b}{2a} = \frac{0}{-2(4)} = 0$$

$$(0) = 4(0)^2 - 10$$

$$0 - 10$$

$$-10$$

$$f(x) = -(x^2 + 8x) + 3$$

$$f(x) = -(x^2 + 8x + 16) + 3 + 16$$

$$f(x) = -(x + 4)^2 + 19$$

Thus the vertex is $(-4, 19)$. Chapter Review Exercises

Chapter 2 Functions and Graphs

Find the vertex. [2.4]

$$\frac{-b}{2a} = \frac{-(-60)}{2(-6)} = \frac{-60}{-12} = 5$$

$$\begin{aligned} (5) &= -6(5)^2 + 60(5) + 11 \\ &= -6(25) + 300 + 11 \\ &= -150 + 300 + 11 \\ &= 161 \end{aligned}$$

Thus the vertex is (5, 161).

Find the vertex. [2.4]

$$\frac{-b}{2a} = \frac{-(-8)}{2(-1)} = \frac{8}{-2} = -4$$

$$\begin{aligned} (-4) &= 14 - 8(-4) - (-4)^2 \\ &= 14 + 32 - 16 \\ &= 30 \end{aligned}$$

Thus the vertex is (-4, 30).

Find the value. [2.4]

$$\begin{aligned} (x) &= -x^2 + 6x - 3 \\ &= -(x^2 - 6x) - 3 \\ &= -(x^2 - 6x + 9) - 3 + 9 \\ &= -(x - 3)^2 + 6 \end{aligned}$$

maximum value of 6

Find the value. [2.4]

$$\begin{aligned} (x) &= 2x^2 + 3x - 4 \\ &= 2\left(x^2 - 2\frac{3}{2}x\right) - 4 \\ &= 2\left(x^2 - 2\frac{3}{2}x + 16\frac{9}{16}\right) - 4 - \frac{9}{8} \\ &= 2\left(x - 4\frac{3}{4}\right)^2 - 5.125 \end{aligned}$$

minimum value of -5.125

Find the maximum height. [2.4]

$$h(t) = -16t^2 + 50t + 4$$

$$\frac{-b}{2a} = \frac{50}{2(-16)} = \frac{25}{16}$$

$$\left(16\frac{25}{16}\right) = -16\left(16\frac{25}{16}\right)^2 + 50\left(16\frac{25}{16}\right) + 4 = 43.0625$$

The ball reaches a maximum height of 43.0625 ft.

68. a. Revenue = 13x [2.5]

$$\begin{aligned} \text{Profit} &= \text{Revenue} - \text{Cost} \\ &= 13x - (0.5x + 1050) \end{aligned}$$

$$\begin{aligned} P &= 13x - 0.5x - 1050 \\ &= 12.5x - 1050 \end{aligned}$$

Break even Revenue = Cost

$$\begin{aligned} 13x &= 0.5x + 1050 \\ 12.5x &= 1050 \\ x &= 84 \end{aligned}$$

The company must ship 84 parcels.

Find the maximum area. [2.4]

Let x be the width. Using the formula for perimeter for three sides, $P = 2w + l$ $700 = 2x + l$

$$l = 700 - 2x$$

Using the formula for area, $A = lw$. Then

$$A(x) = x(700 - 2x)$$

$$A(x) = -2x^2 + 700x$$


$$\frac{b}{2a} = \frac{700}{2(-2)} = -175$$

$$2a2(-2)$$


$$\frac{700}{2} = 350$$

$$A(175) = -2(175)^2 + 700(175) = 61,250 \text{ ft}^2$$

Sketch a graph with different kinds of symmetry. [2.5]



Sketch a graph with different kinds of symmetry. [2.5]



The graph of $y = x^2 - 7$ is symmetric with respect to

the y-axis. [2.5]

The graph of $x = y^2 + 3$ is symmetric with respect to

the x-axis. [2.5]

The graph of $y = x^3 - 4x$ is symmetric with respect to the origin. [2.5]

The graph of $y^2 = x^2 + 4$ is symmetric with respect to the x -axis, y -axis, and the origin. [2.5]


76. The graph of $\frac{x^2}{3^2} + \frac{y^2}{4^2} = 1$ is symmetric with respect to the x -axis, y -axis, and the origin. [2.5]

The graph of $xy = 8$ is symmetric with respect to the origin. [2.5]

The graph of $y = x^3$ is symmetric with respect to the x -axis, y -axis, and the origin. [2.5]

The graph of $x + y = 4$ is symmetric with respect to the origin. [2.5]

Sketch the graph $g(x) = -x^2 + 4$. [2.5]




Domain all real numbers

Range $\{y \mid y \leq 4\}$

g is an even function

Sketch the graph $g(x) = -2x - 4$. [2.5]




Domain all real numbers

Range all real numbers

g is neither even nor odd

Sketch the graph $g(x) = |x - 2| + 2$. [2.5]




Domain all real numbers

Range $\{y \mid y \geq 4\}$

g is an even function

Sketch the graph $g(x) = \sqrt{6 - x^2}$. [2.5]




Domain $\{x \mid -\sqrt{6} \leq x \leq \sqrt{6}\}$

Range $\{y \mid 0 \leq y \leq \sqrt{6}\}$

g is an even function

Sketch the graph $g(x) = x^3 - x$. [2.5]




Domain all real numbers

Range all real numbers

g is an odd function

Sketch the graph $g(x) = 2x$. [2.5]




Domain all real numbers Range


$\{y \mid y \text{ is an even integer}\}$

g is neither even nor odd


$g(x) = f(x) - 2$ [2.5]




87. $g(x) = f(x + 3)$ [2.5]




$g(x) = f(x - 1) - 3$ [2.5]




$g(x) = f(x + 2) - 1$ [2.5]




$g(x) = f(-x)$ [2.5]




$g(x) = -f(x)$ [2.5]




92. $g(x) = 2f(x)$ [2.5]




$g(x) = \frac{1}{2}f(x)$ [2.5]



$g(x) = f(2x)$ [2.5]



$g(x) = f\left(\frac{1}{2}x\right)$ [2.5]



Perform the operations. [2.6]

a. $(f + g)(2) = 2^2 + 2 - 2 + [3(2) + 1]$

$4 + 7 = 11$

b. $\frac{3x^2 + 4x - 2}{3x - 1} - \frac{x^2 - 1}{x + 1} = 1$

$(f - g)(x) = (x^2 + x - 2) - (3x + 1) = x^2 - 2x - 3$

$(f \cdot g)(x) = (x^2 + x - 2)(3x + 1)$
 $3x^2 + 4x^2 - 5x - 2$

Find the difference quotient. [2.6]

$$\begin{aligned} & \frac{f(x+h) - f(x)}{h} \\ &= \frac{4(x+h)^2 - 3(x+h) - 1 - (4x^2 - 3x - 1)}{h} \\ &= \frac{4x^2 + 8xh + 4h^2 - 3x - 3h - 1 - 4x^2 + 3x + 1}{h} \\ &= \frac{8xh + 4h^2 - 3h}{h} \\ &= 8x + 4h - 3 \end{aligned}$$

Find the difference quotient. [2.6]

$$\begin{aligned} & \frac{g(x+h) - g(x)}{h} \\ &= \frac{(x+h)^3 - (x^3)}{h} \\ &= \frac{x^3 + 3x^2h + 3xh^2 + h^3 - x^3}{xh} \\ &= \frac{3x^2h + 3xh^2 + h^3}{h} \\ &= 3x^2 + 3xh + h^2 \end{aligned}$$

99. $s(t) = 3t^2$ [2.4]

a. Average velocity = $\frac{3(4) - 3(2)}{4 - 2}$

$$\begin{aligned} &= \frac{3(16) - 3(4)}{2} \\ &= \frac{48 - 12}{2} \\ &= \frac{36}{2} \\ &= 18 \text{ ft/sec} \end{aligned}$$

b. Average velocity = $\frac{3(3) - 3(2)}{3 - 2}$

$$\begin{aligned} &= \frac{3(9) - 3(4)}{1} \\ &= \frac{27 - 12}{1} = 15 \text{ ft/sec} \end{aligned}$$

c. Average velocity = $\frac{3(2.5) - 3(2)}{2.5 - 2}$

$$\begin{aligned} &= \frac{3(6.25) - 3(4)}{0.5} \\ &= \frac{18.75 - 12}{0.5} \\ &= \frac{6.75}{0.5} = 13.5 \text{ ft/sec} \end{aligned}$$

d. Average velocity = $\frac{3(2.01) - 3(2)}{2.01 - 2}$

$$\begin{aligned} &= \frac{3(4.0401) - 3(4)}{0.01} \\ &= \frac{12.1203 - 12}{0.01} \\ &= \frac{0.1203}{0.01} \\ &= 12.03 \end{aligned}$$

100. Evaluate the composite functions. [2.6]

$$(f \circ g)(3) = f(g(3)) = f(3 - 8)$$

$$\begin{aligned} f(-5) &= (-5)^2 + 4(-5) \\ 25 - 20 &= 5 \end{aligned}$$

$$\begin{aligned} (g \circ f)(-3) &= g(f(-3)) = g((-3)^2 + 4(-3)) \\ g(-3) &= -3 - 8 \\ &= -11 \end{aligned}$$

$$(f \circ g)(x) = f(g(x))$$

$$\begin{aligned} & (x - 8)^2 + 4(x - 8) \\ &= x^2 - 16x + 64 + 4x - 32 \\ &= x^2 - 12x + 32 \end{aligned}$$

$$\begin{aligned} (g \circ f)(x) &= g(f(x)) \\ &= (x^2 + 4x) - 8 \\ &= x^2 + 4x - 8 \end{aligned}$$

101. Evaluate the composite functions. [2.6]

$$\begin{aligned} (f \circ g)(-5) &= f(g(-5)) = f(-5 - 1) = f(-6) \\ &= (-6)^2 + 7 \\ &= 72 + 7 = 79 \end{aligned}$$

$$\begin{aligned} (g \circ f)(-5) &= g(f(-5)) = g(2(-5)^2 + 7) \\ &= g(57) = 57 - 1 \\ &= 56 \end{aligned}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) \\ &= 2|x - 1|^2 + 7 \\ &= 2x^2 - 4x + 2 + 7 \\ &= 2x^2 - 4x + 9 \end{aligned}$$



$$\begin{aligned} (g \circ f)(x) &= g(f(x)) \\ &= |2x^2 + 7 - 1| \\ &= |2x^2 + 6| \\ &= 2x^2 + 6 \end{aligned}$$

$$= 0.01 = 12.03 \text{ ft/sec}$$

It appears that the average velocity of the ball approaches 12 ft/sec.

102. Enter the data on your calculator. The technique for a

TI-83 calculator is illustrated here. Press STAT. [2.7]





$$y = 1.171428571x + 5.19047619$$

$$y = 1.171428571(12) + 5.19047619 \approx 19 \text{ m/s}$$

a. Enter the data on your calculator. The technique for

TI-83 calculator is illustrated here. Press STAT. [2.7]




$$= 0.0047952048t^2 - 1.756843157t + 180.4065934$$

b. Empty $y = 0$ the graph intersects the x -axis.

Graph the equation, and notice that it never intersects

the x -axis.



$$X_{\min} = 0, X_{\max} = 400, X_{\text{scl}} = 100$$

$$Y_{\min} = 0, Y_{\max} = 200, X_{\text{scl}} = 50$$

Thus, no, on the basis of this model, the can never empties.

The regression line is a model of the data and is not based on physical principles.

Chapter 2 Test

Finding the midpoint and length. [2.1]

$$\text{midpoint} = \left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$$

$$= \left(\frac{-2 + 4}{2}, \frac{3 + (-1)}{2} \right)$$

$$= \left(\frac{2}{2}, \frac{2}{2} \right) = (1, 1)$$

$$\text{length} = d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

$$\sqrt{(-2 - 4)^2 + (3 - (-1))^2}$$

$$= \sqrt{(-6)^2 + 4^2} = \sqrt{36 + 16} = \sqrt{52}$$

$$= 2\sqrt{13}$$

Finding the x - and y -intercepts and graphing. [2.1]

$$= 2y^2 - 4$$

$$= 0 \quad x = 2(0)^2 - 4 = -4$$

Thus the x -intercept is $(-4, 0)$.


$$= 0 \quad 0 = 2y^2 - 4$$

$$= 2y^2$$


$$2 = y^2$$

$$2 = \sqrt{y}$$

Thus the y -intercepts are $(0, -\sqrt{2})$ and $(0, \sqrt{2})$.



Graphing $y = |x - 2| - 1$. [2.1]



Finding the center and radius. [2.1]

$$x^2 - 4x + y^2 + 2y - 4 = 0$$

$$(x^2 - 4x) + (y^2 + 2y) = 4$$

$$(x^2 - 4x + 4) + (y^2 + 2y + 1) = 4 + 4$$

$$+1(x - 2)^2 + (y + 1)^2 = 9$$

center (2, -1), radius 3


Determining the domain of the function. [2.2]

$$x^2 - 16 \geq 0$$

$$(x - 4)(x + 4) \geq 0$$

The product is positive or zero.

The critical values are -4 and 4.



The domain is $\{x | x \leq -4 \text{ or } x \geq 4\}$.

Find the values of a in the domain of

$$f(x) = x^2 + 6x - 17 \text{ for which } f(a) = -1. [2.2]$$

$$a^2 + 6a - 17 = -1 \text{ Replace } f(a) \text{ with } a^2 + 6a - 17$$

$$a^2 + 6a - 16 = 0$$

$$(a + 8)(a - 2) = 0$$

$$a + 8 = 0 \quad a - 2 = 0$$

$$a = -8 \quad a = 2$$

Find the slope. [2.3]

$$m = \frac{3 - (-2)}{-1 - 5} = \frac{5}{-6} = -\frac{5}{6}$$

Find the equation. [2.3]

$$-(-3) = -2(x - 5) + y$$

$$+3 = -2x + 10$$

$$y = -2x + 7$$

Finding the equation in slope-intercept form. [2.3]

$$3x - 2y = 4$$

$$-2y = -3x + 4$$

$$= 2\frac{3}{2}x - 2$$

Slope of perpendicular line is $-\frac{2}{3}$.

$$y - y_1 = m(x - x_1)$$

$$\frac{2}{3}$$

$$y + 2 = -\frac{2}{3}(x - 4)$$

$$y + 2 = -\frac{2}{3}x + \frac{8}{3}$$

$$= -\frac{2}{3}x + \frac{8}{3} - \frac{6}{3}$$

$$= -\frac{2}{3}x + \frac{2}{3}$$

Write in standard form, find the vertex and the axis of symmetry. [2.4]

$$f(x) = x^2 + 6x - 2$$

$$(x^2 + 6x + 9) - 2 - 9$$

$$= (x + 3)^2 - 11 \text{ standard form,}$$

vertex (3, -11), axis of symmetry $x = -3$

Finding the maximum or minimum value. [2.4]

$$\frac{b}{2a} = \frac{-6}{2(1)} = -3$$

$$(2) = 2^2 - 4(2) - 8$$

$$4 - 8 - 8$$

$$-12$$

The minimum value of the function is -12.

Classifying the functions as even, odd or neither. [2.5]

$$f(x) = x^4 - x^2$$

$$f(-x) = (-x)^4 - (-x)^2 = x^4 - x^2 = f(x)$$

$f(x)$ is an even function.

$$f(x) = x^3 - x$$

$$f(-x) = (-x)^3 - (-x) = -x^3 + x$$

$$= -(x^3 - x) = -f(x)$$

$f(x)$ is an odd function.

$$f(x) = x - 1$$

$$f(-x) = -x - 1 \neq f(x) \text{ not an even function}$$

$$(-x) = -x - 1 \neq -f(x) \text{ not an odd function}$$

neither

Identify the type of symmetry. [2.5]

$$\mathbf{a.} (-y)^2 = x + 1$$


$$y^2 = x + 1 \text{ symmetric with respect to } x\text{-axis}$$

b. $y = 2(-x)^3 + 3(-x)$


$y = 2x^3 + 3x$ symmetric with respect to origin

$y = 3(-x)^2 - 2$


14.




15. $g(x) = f\left(\frac{1}{2}x\right)$ [2.5]




16. $g(x) = -f(x)$ [2.5]



17. $g(x) = f(x-1) + 3$ [2.5]



18.



19. Perform the operations. [2.6]

a. $(f-g)(x) = (x^2 - x + 2) - (2x - 1) = x^2 - 3x + 3$

b. $(f \cdot g)(-2) = ((-2)^2 - (-2) + 2)((-2) - 1)$
 $= (8)(-5) = -40$

c. $(f \circ g)(3) = f(g(3)) = f(2(3)-1)$
 $= f(5) = 5^2 - 5 + 2$
 $= 22$

d. $(g \circ f)(x) = g(f(x)) = 2(x^2 - x + 2) - 1$
 $= 2x^2 - 2x + 3$

20. Finding the difference quotient of the function. [2.6]

$f(x) = x^2 + 1$

$$\frac{f(x+h) - f(x)}{h} = \frac{(x+h)^2 + 1 - (x^2 + 1)}{h}$$

$$= \frac{x^2 + 2xh + h^2 + 1 - x^2 - 1}{h}$$

$$= \frac{2xh + h^2}{h} = \frac{h(2x+h)}{h}$$

$$= 2x + h$$

Find the maximum area. [2.4]

Using the formula for perimeter for three sides,

$$P = 2w + l \quad 80 = 2x + y$$

$$= 80 - 2x$$

Using the formula for area, $A = xy$. Then

$$A(x) = x(80 - 2x)$$

$$A(x) = -2x^2 + 80x$$

$$\frac{b}{2a} = -\frac{80}{2(-2)} = 20$$

$$2a2(-2)$$

$$= 80 - 2(20) = 40$$

$$= 20 \text{ ft and } y = 40 \text{ ft}$$

Evaluating the function, $s(t) = 5t^2$. [2.6]

Average velocity = $\frac{5(3)^2 - 5(2)^2}{3-2} = \frac{5(9) - 5(4)}{3-2}$

$$= \frac{45 - 20}{1} = 25 \text{ ft/sec}$$

b. Average velocity = $\frac{5(2.5)^2 - 5(2)^2}{2.5-2}$

$$= \frac{5(6.25) - 5(4)}{0.5}$$

$$= \frac{31.25 - 20}{0.5} = 22.5 \text{ ft/sec}$$

c. Average velocity = $\frac{5(2.01)^2 - 5(2)^2}{2.01-2}$

$$= \frac{5(4.0401) - 5(4)}{0.01}$$


$$= \frac{20.2005 - 20}{0.01} = 20.05 \text{ ft/sec}$$

a. Enter the data on your calculator. The technique for

a TI-83 calculator is illustrated here. Press STAT. [2.7]



L1	L2	L3	3
93.2	28		
92.3	26		
91.9	39		
89.5	56		
89.6	56		
90.5	36		
L3(1)=			



LinReg
y=ax+b
a=-7.98245614
b=767.122807
r2=.805969575
r=-.8977580826

Evaluating the equation from part (a) at 89.

$$y = 7.98245614(89) + 767.122807 = 57$$

calories

Cumulative Review Exercises

Determine the property for $3(a + b) = 3(b + a)$. [P.1]

Commutative Property of Addition

$\frac{6}{-}$, $\sqrt{2}$ are not rational numbers [P.1]

Simplifying. [P.1]

$$3 + 4(2x - 9)$$

$$= 3 + 8x - 36$$

$$= 8x - 33$$

Simplifying. [P.2]

$$(-4xy^2)^3 (-2x^2y^4) = (-64x^3y^6)(-2x^2y^4)$$

$$= (-64)(-2)(x^{3+2}y^{6+4})$$

$$= 128x^5y^{10}$$

Simplifying. [P.2]

$$\frac{24a^4b^3}{-2} = \frac{4a^{4-4}b^{3-5}}{-2} = \frac{4b^{-2}}{-2} = -2b^2$$

$$\frac{18a^4b^5}{3} = \frac{6a^4b^5}{3} = 2a^4b^5$$

Simplifying. [P.3]

$$(2x + 3)(3x - 7) = 6x^2 - 5x - 21$$

Simplifying. [P.5]

$$\frac{x^2 + 6x - 27}{x^2 - 9} = \frac{(x + 9)(x - 3)}{(x + 3)(x - 3)} = \frac{x + 9}{x + 3}$$

Simplifying. [P.5]

$$\frac{4}{2x-1} - \frac{2}{x-1} = \frac{4(x-1)}{(2x-1)(x-1)} - \frac{2(2x-1)}{(2x-1)(x-1)}$$

$$= \frac{4x - 4 - 4x + 2}{(2x-1)(x-1)}$$

$$= \frac{-2}{(2x-1)(x-1)}$$

Solving for x. [1.1]

$$6 - 2(2x - 4) = 14$$

$$6 - 4x + 8 = 14$$

$$-4x = 0$$

$$x = 0$$

$$y = 7.98245614x - 767.122807$$

Chapter 2 Functions and Graphs

Solving for x . [1.3]

$$x^2 - x - 1 = 0$$

$$x = \frac{-(-1) \pm \sqrt{(-1)^2 - 4(1)(-1)}}{2(1)}$$

$$x = \frac{1 \pm \sqrt{4+1}}{2} = \frac{1 \pm \sqrt{5}}{2}$$

Solving for x . [1.3]

$$(2x - 1)(x + 3) = 4$$

$$2x^2 + 5x - 3 = 4$$

$$2x^2 + 5x - 7 = 0$$

$$(2x + 7)(x - 1) = 0$$

$$x = -\frac{7}{2} \text{ or } x = 1$$

$$3x + 2y = 15$$

$$3x = -2y + 15$$

$$\frac{2}{3}x = -\frac{2}{3}y + 5$$

$$x = -\frac{2}{3}y + 5$$

Solving for x . [1.4]

$$x^4 - x^2 - 2 = 0$$

$$\text{Let } u = x^2.$$

$$u^2 - u - 2 = 0$$

$$(u - 2)(u + 1) = 0$$

$$u - 2 = 0 \quad \text{or} \quad u + 1 = 0$$

$$u = 2 \quad \quad \quad u = -1$$

$$x^2 = 2 \quad \quad \quad x^2 = -1$$

$$x = \sqrt{2} \quad \quad \quad x = i$$

Solving for x . [1.5]

$$3x - 1 < 5x + 7$$

$$-2x < 8$$

$$x > -4$$

Finding the distance. [2.1]

$$\text{distance} = \sqrt{[-2 - 2]^2 + [-4 - (-3)]^2}$$

$$= \sqrt{(-4)^2 + (-1)^2} = \sqrt{16 + 1} = \sqrt{17}$$

Finding $G(-2)$. [2.2]

$$G(x) = 2x^3 - 4x - 7$$

$$G(-2) = 2(-2)^3 - 4(-2) - 7 = 2(-8) + 8 - 7 = -15$$

Finding the equation of the line. [2.3]

$$\text{The slope is } m = \frac{-1 - (-3)}{-2 - 2} = \frac{-1 + 3}{-4} = \frac{2}{-4} = -\frac{1}{2}$$

$$\text{The equation is } y - (-3) = -\frac{1}{2}(x - 2)$$

Solving a mixture problem. [1.1]

	x
0.08	60
0.03	x
60 + x	

$$0.08(60) + 0x = 0.03(60 + x)$$

$$4.8 = 1.8 + 0.03x$$

$$3.0 = 0.03x$$

$$x = 100$$

100 ounces of water

Evaluating a quadratic function. [2.4]

$$h(x) = -0.002x^2 - 0.03x + 8$$

$$h(39) = -0.002(39)^2 - 0.03(39) + 8 = 3.788 \text{ ft}$$

Yes.

Finding the rate, or slope. [2.3]

$$0.04^\circ\text{F/min}$$