## Solution Manual for Natural Disasters Canadian 4th Edition Abbott 1259451682

### Full link download:

Test Bank:

https://testbankpack.com/p/test-bank-for-natural-disasters-canadian-4th-edition-abbott-1259451682/

Solution Manual:

https://testbankpack.com/p/solution-manual-for-natural-disasters-canadian-4th-edition-abbott-1259451682/

### **Chapter 2 Earth's Internal Energy and Plate Tectonics**

### **Overview**

Natural disasters necessarily involve the concentration and release of energy. Sources of energy include the Earth's internal heat generated from radioactive decay, solar energy, gravity, and the impact of extraterrestrial bodies.

The most basic divisions of the Earth are, based on density, core, mantle, and crust, or, based on material strength, core, mesosphere, asthenosphere, and lithosphere. The core is the innermost and densest region, composed mainly of iron with a solid inner part and a liquid outer part. Circulation in the outer core creates the Earth's magnetic field. The rocky mantle, extending from roughly half the Earth's radius to close to the Earth's surface, is the second densest region. The shallowest and least dense part of the solid Earth is the rocky crust.

With the exception of the uppermost part, the mantle is hot enough that it flows even though it is mostly solid. This soft plastic layer is called the asthenosphere. The lithosphere floats buoyantly on top of the asthenosphere like icebergs float in seawater. The Earth's lithosphere is cold and solid, and is divided into a number of thin but very wide plates. According to the theory of plate tectonics developed principally by Professor John Tuzo Wilson of the University of Toronto in the 1960s, the relative motions of these plates are responsible for most earthquakes, volcanic eruptions, and mountain-building on Earth. A new plate is formed by upwelling and solidification of magma at divergent

zones, and old plates are recycled into the asthenosphere at subduction zones, a type of convergent zone. Evidence of plate tectonics includes magnetization patterns on the seafloor, age of the ocean basins, the distribution of ocean water depth, epicentre and hypocentre locations, and fossil correlations.

Most earthquakes and volcanoes occur at plate boundaries. At spreading centers (divergent zones), the high temperatures near the Earth's surface preclude the occurrence of very powerful or deep earthquakes, though numerous moderate and shallow earthquakes occur. In contrast, the deepest earthquakes and the most powerful earthquakes occur at convergent zones (subduction zones and continent-continent collision zones). Large earthquakes can also occur at transform plate boundaries.

Most volcanic activity on Earth takes place at spreading centres where large amounts of lava are extruded peacefully underwater. Dangerous, explosive volcanic eruptions mostly occur at subduction zones.


#### © 2017 McGraw-Hill Education Limited 1


#### Learning Objectives

- 1. List the sources of energy fueling natural hazards.
- 2. Describe the Earth's internal structure.
- 3. Explain the behavious of materials under stress.
- 4. Explain how plate tectonics operates.
- 5. Evaluate the evidence for plate tectonics.
- 6. Explain the relationship between plate tectonics and the location of earthquakes and volcanoes.

### **Critical Thinking Questions**

- 1. There are three different styles of rock deformation: elastic, ductile, brittle, and plastic. Can you identify various common objects that show these styles of deformation?
- 2. The lithospheric plates move around the surface of the Earth, but the surface area of the Earth has not changed over geological time. If a new spreading center (divergent zone) were to develop, what other plate changes might also occur at the same time?






# NATURAL

Fourth Canadian Edition

Patrick Abbott Claire Samson

ABBOTT | SAMSON



# Chapter 2

Earth's Internal Energy and Plate Tectonics

Prepared by Bill Buhay University of Winnipeg

# **Learning Outcomes**

- LO1 List the sources of energy fueling natural hazards
- LO2 Describe the Earth's internal structure
- LO3 Explain the behaviour of materials under stress
- LO4 Explain how plate tectonics operates
- LO5 Evaluate the evidence for plate tectonics
- LO6 Explain the relationship between plate tectonics and the location of earthquakes and volcanoes

### **Outline**

Energy Sources of Natural Hazards

Origin of the Sun and Planets

The Earth's Early History

The Layered Earth

Plate Tectonics: Recycling the Earth's Outer Layers

**Evidence of Plate Tectonics** 

Plate-Tectonic Setting of Earthquakes and Volcanoes

Summary

## **Energy Sources of Natural Hazards**

### **Primary energy sources:**

### Earth's internal energy

- Primarily from the radioactive decay of elements
- Smaller contributions from impacts/gravitational compaction (Earth's early history)

### Gravity

- Mass movements
- Snow avalanche

## **Energy Sources of Natural Hazards**

### **Primary energy sources:**

Solar energy


- Hydrologic cycle
- With gravity powers the agents of erosion
- Stored in plants

The impact of extraterrestrial bodies

Asteroids and comets

## **Energy Sources of Natural Hazards**

Figure 2.2



The rotations and orbits of the Earth-Moon-Sun system result in tremendous amounts of energy.

## Origin of the Sun and Planets

### Initial stage

Rotating spherical cloud (ice, gas, debris)





Figure 2.3

### Spinning mass

Contracts into flattened disk (planets grow as particles collide and stick together)



## Ignited Sun

Surrounded by planets –Earth is the third planet from the Sun

## Origin of the Sun and Planets

### AGE OF THE EARTH

The oldest Solar System materials are about 4.57 billion years old

The oldest Earth rocks are 4.03 billion years old (Acasta gneiss, Northwest Territories, Canada)

The oldest ages obtained from Earth materials (zircons) are 4.37 billion years old (extracted from a 3.1 billion year old sandstone –western Australia)

Our planet has existed for about 4.5 billion years

## The Earth's Early History

Figure 2.5

Heat-Transforming the Early Earth

Impact energy


Decay of radioactive isotopes Gravitational energy

Massive amount of heat produced widespread melting that caused the formation of:

Primitive crust (low –density rock)

Large oceans

Dense atmosphere



# The Layered Earth

### **DENSITY LAYERS**

### Layers based on:

Density due to different chemical and mineral compositions

Core

Inner core

Outer core

Mantle

Crust

Different strengths

Mesosphere

Asthenosphere

Lithosphere




Figure 2.10

# The Layered Earth


### STRENGTH LAYERS

Figure 2.11

Mesosphere –stiff plastic solid -extends from the core-mantle boundary to ~ 360 km below the Earth's surface

Asthenosphere —soft plastic —surrounds the mesosphere

Lithosphere – rigid outer layer – varies in thickness between a few km under oceans to 100 km under continents.



## The Layered Earth


**ISOSTASY** 

**Buoyancy principle** 

low density continents and mountain ranges "float" on the denser mantle below

post-glacial rebound effect in Canada

Figure 2.13



## In Greater Depth


# MATERIAL DEFORMATION

Figure 2.7

materials subjected to stress, they deform or undergo strain.

Stress can be applied perpendicular to the surface of a body, causing it to stretch under tension or to contract under compression.

On the other hand, shear stress, which is applied parallel to the surface, tends to deform a body along internal planes slipping past one another.



# **In Greater Depth**

# MATERIAL DEFORMATION

### Figure 2.8

### **Behaviour of materials**

- a) **Elastic**: bend thin board; let it go and board recovers its original shape.
- b) **Ductile**: squeeze a wad of bubblegum or Silly Putty<sup>®</sup>; let it go but it stays in the deformed shape.
- c) **Erittle**: bend thin board sharply and it breaks.




The lithosphere is broken into pieces called plates.

Plate tectonics
Greek word
"tekton" = to
build tectonics.

Dozen larger plates (Pacific plate) and several small plates (Juan de Fuca).

Figure 2.16



Tectonic forces form:


Figure 2.18

Mountains/plateaus

Mid-ocean ridges

Deep ocean trenches


Chains of volcanic islands



~ 250 million years to complete the tectonic cycle (seafloor spreading –subduction).

Ocean basins < 200 million years old.

Figure 2.19



Magma rises from the asthenosphere –surfaces at oceanic volcanic ridges.

As the magma cools, the plates move laterally away (continue to cool).

Plates grow thicker, become denser, collide with less dense plates —then subductand assimilate in the asthenosphere.




Figure 2.20

Figure 2.21

Convection cells in the mantle drive the movement of the overlying tectonic plates.





Figure 2.22

Divergent zones

Convergent zones

Transform plate boundaries

Hot spots



**DIVERGENT ZONES** 


Figure 2.23

Centering: lithosphere moves over centresof anomalously hot mantle regions

**Doming:** mantle heat causes surface doming through uplifting, stretching and fracturing

Riffing: dome's central area sages – forms valley

**Spreading:** Advanced pulling apart – forming new seafloor



# Plate Tectonics: Recycling the Earth's DIVERGENT ZONES Outer Layers Figure 2.24

Triple Junction (Afar Triangle)

Northeast Africa being torn apart by three spreading centres:

Red Sea

Gulf of Aden

East African rift system





Figure 2.24

Triple junction. (a) Schematic example. (b) Northeast Africa is being torn apart by three spreading centres: the Red Sea, the Guif of Aden, and the East African rift system. The spreading centres meet at a triple junction in the Afar Triangle.



### **CONVERGENT ZONES**

Plates deform under compression at convergent zones.

### Three classes of collisions:


Oceanic plate-Oceanic plate

Oceanic plate-Continent plate

Continent plate-Continent plate

### **CONVERGENT ZONES**

- a) Oceanic-oceanic convergence formation of a deep offshore trench and seafloor volcanoes that can eventually form an island arc
- b) Oceanic-continental convergence formation of a coastal trench and adjacent chain of volcanoes



### **CONVERGENT ZONES**

Bonne Bay, GrosMorne National Park, Newfoundland.

500 million years ago –piece of oceanic plate was scraped off the subductingplate.

Provides geologists with exposed oceanic plate.

Figure 2.26



### **CONVERGENT ZONES**

Figure 2.27

When Godwanalandbegan separating —India moved northward towards Asia.

Once the seafloor between then had been subducted(under Asia) —India collided with Asia (~2000 km of crustal shortening/mountain building so far).





Figure 2.29

### **CONVERGENT ZONES**

Himalaya Mountains/Tibetan Plateau.

Tectonic Map –India pushing north and colliding with Asia.

Collision resulting in crustal shortening and the uplift of the Himalayan mountains and the Tibetan Plateau.




Transform plate boundaries

Figure 2.31

Shear stress at transform plate boundaries.

To accommodate the Earth's curvature the plates must fracture – transform faults.




# Human Focus: John Tuzo Wilson (1908-1993)

Figure 2.33

1963 -1966: Tuzo Wilson (U of Toronto)

Father of Plate Tectonics



**Hot Spots** 

Figure 2.32a

hot mantle rock (from the mesosphere) rises up through the asthenosphere and lithosphere as a plume of magma supplying a volcano.

lithospheric plate keeps moving -new volcanoes are formed.





Figure 2.32b


A chain of hot spot fed volcanoes formed as the Pacific plate moved (changed direction between 47 and 63 million years ago).



# **Evidence of Plate Tectonics**MAGNETIZATION PATTERNS ON THE SEAFLOORS

Figure 2.34

Magnetic polarity
measurements in volcanic
rocks combined with
radiometric ages
determined from the same
rocks allow the building of
a timescale based on
magnetic polarity reversals.



#### **MAGNETIZATION PATTERNS ON THE SEAFLOORS**

magnetically striped Pacific Ocean floor.

black areas are magnetized pointing to a north pole, yellow areas to a south pole.


mirror-image patterns on each side of the Juan de Fuca ridge.



#### **MAGNETIZATION PATTERNS ON THE SEAFLOORS**

Cross-section of magnetically striped seafloor.


mirror-image pattern documents movements away from oceanic ridges.



#### **EARTHQUAKE EPICENTRESAND HYPOCENTRES**

Shallow hypocentres - earthquakes generated in brittle rocks in both subducting(brown) and overriding (orange) plates


Deeper hypocentres - only the interior of the subductingPacific plate is cold enough to maintain the rigidity necessary to produce earthquakes



#### CONTINUITY OF GEOLOGICAL FEATURES

Figure 2.38

distribution of several fossils is continuous when the continents are restored to the position they occupied when these life forms existed.




## In Greater Depth

#### THE DANCE OF THE CONTINENTS

continuity of mountain ranges and fossils across oceans.

suggest that all the continents were once united in a supercontinent (Pangaea -220 million years ago).

Figure 2.40



## In Greater Depth

#### THE DANCE OF THE CONTINENTS

positions of the continents:

- (a) 180 million years ago
- (b) 135 million years ago
- (c) 65 million years ago
- (d) Today




Figure 2.39

idealized tectonic plate and the varying earthquake hazards that are concentrated at plate edges




Table 2.3

TABLE 2.3

TECTONIC ENVIRONMENTS AND EARTHQUAKE CHARACTERISTICS


		Earthquake Characteristics		
		Eartifquake Characteristics		
Tectonic	Deformation	Frequency	Maximum	Maximum Hypocentre
Environment	Force		Size	Depth
Divergent zone	Tension	Frequent	Strong	Shallow
Convergent zone	Compression	Infrequent	Great	Deep
Transform fault	Shear	Infrequent	Major	Shallow
Hot spot	Tension	Frequent	Strong	Shallow
	Compression			

TABLE 2.4		THE EARTH'S GREATEST EARTHQUAKES, 1900-2013		
Year	Location	Magnitude	Cause	
1960	Chile	9.5	<u>Subduction</u> —Nazca plate under South American plate	
1964	Alaska	9.2	<u>Subduction</u> —Pacific plate under North American plate	
2004	Indonesia	9.1	<u>Subduction</u> — <u>Indo</u> -Australian plate under Eurasian plate	
2011	Japan	9.0	Subduction—Pacific plate under North American plate	
1952	Kamchatka	9.0	<u>Subduction</u> —Pacific plate under North American plate	
2010	Chile	8.8	<u>Subduction</u> —Nazca plate under South American plate	
1906	Ecuador	8.8	<u>Subduction</u> —Nazca plate under South American plate	
1965	Alaska	8.7	<u>Subduction</u> —Pacific plate under North American plate	
2012	Indonesia	8.6	Subduction—Indo-Australian plate under Eurasian plate	
2005	Indonesia	8.6	Subduction—Indo-Australian plate under Eurasian	

Table 2.4

Figure 2.42

Volcanic hazards associated with an idealized tectonic plate.



## Summary

The Solar system formed over a short period –earth ~ 4.57 billion years old.

Gravity pulled the Earth into layers of differing density – stratified –solid inner core surrounded by a liquid outer core.

Convection within the plastic mesosphere and asthenoshpere.

Earth's outer layers –ridged lithosphere.

Earth's internal energy is heat generated by ongoing radioactive decay —contributions from impact energy and gravitational attraction.

Earth's internal energy drives earthquakes and volcanos.

Earth's outer layer is broken into tectonic plates (12 large, several smaller ones).

The boundary between the rigid lithosphere and the underlying plastic asthenosphere provides a sliding surface for plates to move.

Convection in the asthenosphere and gravity are the driving mechanisms of plate movement.

Different processes take place in the four different tectonic environments: (1) divergent zones, (2) convergent zones, (3) transform plate boundaries, and (4) hot spots.

New lithosphere is created by seafloor spreading when plates are pulled apart at divergent zones. Old lithosphere is being reabsorbed into the asthenosphere by subduction where two plates of differing densities collide.

Transform faults accommodate the along-side movements of plates without creation or destruction of lithosphere. Hot spots originate deep

Hot spots originate deep in the Earth and are the source of isolated plumes of partially molten rock rising through the asthenosphere and lithosphere.

Ocean studies have contributed solid evidence attesting to the movement of tectonic plates. Magnetization patterns on the seafloor delineate the newly formed lithosphere in space and time. Bathymetric surveys have mapped deep subduction trenches and relatively shallow volcanic ridges.

Most earthquakes are caused by fault movements and occur preferentially along the edges of tectonic plates. Magma movement in the shallow subsurface at spreading centresand hot spots tends to induce swarms of small earthquakes.

The dominantly horizontal movements at transform faults produce large earthquakes. The compressional movements at subduction zones and continent—continent collisions generate the largest earthquakes.

Most volcanic eruptions occur underwater at spreading centreswhere an enormous volume of lava solidifies to form new seafloor.

## END OF CHAPTER 2