

**Solution Manual for Precalculus Enhanced with Graphing Utilities 6th
Edition Sullivan 0321795466 9780132854351**

Full link download:

Test Bank:

<https://testbankpack.com/p/test-bank-for-prec calculus-enhanced-with-graphing-utilities-6th-edition-sullivan-0321795466-9780132854351/>

Solution Manual:

<https://testbankpack.com/p/solution-manual-for-prec calculus-enhanced-with-graphing-utilities-6th-edition-sullivan-0321795466-9780132854351/>

Chapter 2

Functions and Their Graphs

Section 2.1


1. $(-1, 3)$

$$\begin{aligned}
 2. \quad 3(-2)^2 - 5(-2) + \frac{1}{(-2)} &= 3(4) - 5(-2) - \frac{1}{2} \\
 &= 12 + 10 - \frac{1}{2} \\
 &= \frac{43}{2} \text{ or } 21\frac{1}{2} \text{ or } 21.5
 \end{aligned}$$

3. We must not allow the denominator to be 0.
 $x + 4 \neq 0 \Rightarrow x \neq -4$; Domain: $\{x \mid x \neq -4\}$.

4. $3 - 2x > 5$
 $-2x > 2$
 $x < -1$

Solution set: $\{x \mid x < -1\}$ or $(-\infty, -1)$




5. independent; dependent

6. range

7. $[0, 5]$

We need the intersection of the intervals $[0, 7]$ and $[-2, 5]$. That is, domain of $f \cap$ domain of g .



8. \neq ; f ; g

9. $(g - f)(x)$ or $g(x) - f(x)$

10. False; every function is a relation, but not every

11. True

12. True

13. False; if the domain is not specified, we assume it is the largest set of real numbers for which the value of f is a real number.

14. False; the domain of $f(x) = \frac{x^2 - 4}{x}$ is $\{x \mid x \neq 0\}$.

15. Function

Domain: {Elvis, Colleen, Kaleigh, Marissa}
 Range: {Jan. 8, Mar. 15, Sept. 17}

16. Not a function

17. Not a function

18. Function

Domain: {Less than 9th grade, 9th-12th grade,

High School Graduate, Some College, College Graduate}

Range: {\$18,120, \$23,251, \$36,055, \$45,810, \$67,165}

19. Not a function

20. Function

Domain: $\{-2, -1, 3, 4\}$
 Range: $\{3, 5, 7, 12\}$

21. Function

Range: $\{3\}$

22. Function

Domain: $\{0, 1, 2, 3\}$
 Range: $\{-2, 3, 7\}$

23. Not a function

24. Not a function

25. Function

Domain: $\{-2, -1, 0, 1\}$
 relation is a function. For example, the relation

$x^2 + y^2 = 1$ is not a function.


Range: $\{0, 1, 4\}$

26. Function


Domain: $\{-2, -1, 0, 1\}$

Range: $\{3, 4, 16\}$


27. Graph $y = x^2$. The graph passes the vertical line test. Thus, the equation represents a function.




28. Graph $y = x^3$. The graph passes the vertical line test. Thus, the equation represents a function.



29. Graph $y = \frac{1}{x}$. The graph passes the vertical line test. Thus, the equation represents a function.



30. Graph $y = |x|$. The graph passes the vertical line test. Thus, the equation represents a function.



31. $y^2 = 4 - x^2$

Solve for y : $y = \pm\sqrt{4 - x^2}$

For $x = 0$, $y = \pm 2$. Thus, $(0, 2)$ and $(0, -2)$ are on the graph. This is not a function, since a distinct x -value corresponds to two different y -values.


32. $y = \pm\sqrt{1 - 2x}$

34. $x + y^2 = 1$


Solve for y : $y = \pm\sqrt{1 - x}$

For $x = 0$, $y = \pm 1$. Thus, $(0, 1)$ and $(0, -1)$ are on the graph. This is not a function, since a distinct x -value corresponds to two different y -values.

35. Graph $y = 2x^2 - 3x + 4$. The graph passes the vertical line test. Thus, the equation represents a function.



36. Graph $y = \frac{3x - 1}{x + 2}$. The graph passes the vertical line test.



37. $2x^2 + 3y^2 = 1$

Solve for y : $2x^2 + 3y^2 = 1$

$$3y^2 = 1 - 2x^2$$

$$y^2 = \frac{1 - 2x^2}{3}$$

$$y = \pm\sqrt{\frac{1 - 2x^2}{3}}$$

For $x = 0$, $y = \pm\sqrt{\frac{1}{3}}$. Thus, $(0, \frac{1}{\sqrt{3}})$ and $(0, -\frac{1}{\sqrt{3}})$

are on the graph. This is not a function.

For $x = 0$, $y = \pm 1$. Thus, $(0, 1)$ and $(0, -1)$ are on the graph. This is not a function,

Chapter 2: Functions and Their Graphs

since a distinct x -value corresponds to two different y -values.

33. $x = y^2$

Solve for y : $y = \pm\sqrt{x}$

For $x = 1$, $y = \pm 1$. Thus, $(1, 1)$ and $(1, -1)$ are on the graph. This is not a function, since a distinct x -value corresponds to two different y -values.

Section 2.1: Functions

function, since a distinct x -value corresponds to two different y -values.

38. $x^2 - 4y^2 = 1$

Solve for y: $x^2 - 4y^2 = 1$

$$4y^2 = x^2 - 1$$

$$y^2 = \frac{x^2 - 1}{4}$$

$$y = \frac{\pm\sqrt{x^2 - 1}}{2}$$

For $x = \sqrt{2}$, $y = \pm \frac{1}{2}$. Thus, $(\sqrt{2}, \frac{1}{2})$ and $(\sqrt{2}, -\frac{1}{2})$ are on the graph. This is not a function, since a distinct x -value corresponds to two different y -values.

39. $f(x) = 3x^2 + 2x - 4$

a. $f(0) = 3(0)^2 + 2(0) - 4 = -4$

b. $f(1) = 3(1)^2 + 2(1) - 4 = 3 + 2 - 4 = 1$

c. $f(-1) = 3(-1)^2 + 2(-1) - 4 = 3 - 2 - 4 = -3$

d. $f(-x) = 3(-x)^2 + 2(-x) - 4 = 3x^2 - 2x - 4$

e. $-f(x) = -(3x^2 + 2x - 4) = -3x^2 - 2x + 4$

f. $f(x+1) = 3(x+1)^2 + 2(x+1) - 4$
 $= 3(x^2 + 2x + 1) + 2x + 2 - 4$

$$= 3x^2 + 6x + 3 + 2x + 2 - 4$$

$$= 3x^2 + 8x + 1$$

g. $f(2x) = 3(2x)^2 + 2(2x) - 4 = 12x^2 + 4x - 4$

e. $-f(x) = -(-2x^2 + x - 1) = 2x^2 - x + 1$

f. $f(x+1) = -2(x+1)^2 + (x+1) - 1$
 $= -2(x^2 + 2x + 1) + x + 1 - 1$
 $= -2x^2 - 4x - 2 + x$

$$= -2x^2 - 3x - 2$$

g. $f(2x) = -2(2x)^2 + (2x) - 1 = -8x^2 + 2x - 1$

h. $f(x+h) = -2(x+h)^2 + (x+h) - 1$
 $= -2(x^2 + 2xh + h^2) + x + h - 1$
 $= -2x^2 - 4xh - 2h^2 + x + h - 1$

41. $f(x) = \frac{x}{x^2 + 1}$

a. $f(0) = \frac{0}{0^2 + 1} = \frac{0}{1} = 0$

b. $f(1) = \frac{1}{1^2 + 1} = \frac{1}{2}$

c. $f(-1) = \frac{-1}{(-1)^2 + 1} = \frac{-1}{1 + 1} = -\frac{1}{2}$

d. $f(-x) = \frac{-x}{(-x)^2 + 1} = \frac{-x}{x^2 + 1}$

e. $-f(x) = -\left(\frac{x}{x^2 + 1}\right) = \frac{-x}{x^2 + 1}$

f. $f(x+1) = \frac{x+1}{(x+1)^2 + 1}$

Chapter 2: Functions and Their Graphs**Section 2.1: Functions**

$$\begin{aligned} \text{h. } f(x+h) &= 3(x+h)^2 + 2(x+h) - 4 \\ &= 3(x^2 + 2xh + h^2) + 2x + 2h - 4 \\ &= 3x^2 + 6xh + 3h^2 + 2x + 2h - 4 \end{aligned}$$

$$40. f(x) = -2x^2 + x - 1$$

$$\text{a. } f(0) = -2(0)^2 + 0 - 1 = -1$$

$$\text{b. } f(1) = -2(1)^2 + 1 - 1 = -2$$

$$\text{c. } f(-1) = -2(-1)^2 + (-1) - 1 = -4$$

$$\text{d. } f(-x) = -2(-x)^2 + (-x) - 1 = -2x^2 - x - 1$$

$$\begin{aligned} &= \frac{x+1}{x^2 + 2x + 1 + 1} \\ &= \frac{x+1}{x^2 + 2x + 2} \end{aligned}$$

$$\text{g. } f(2x) = \frac{2x}{(2x)^2 + 1} = \frac{2x}{4x^2 + 1}$$

$$\text{h. } f(x+h) = \frac{x+h}{(x+h)^2 + 1} = \frac{x+h}{x^2 + 2xh + h^2 + 1}$$

42. $f(x) = \frac{x^2 - 1}{x + 4}$

a. $f(0) = \frac{0^2 - 1}{0 + 4} = \frac{-1}{4} = -\frac{1}{4}$

b. $f(1) = \frac{1^2 - 1}{1 + 4} = \frac{0}{5} = 0$

c. $f(-1) = \frac{(-1)^2 - 1}{-1 + 4} = \frac{0}{3} = 0$

d. $f(-x) = \frac{(-x)^2 - 1}{-x + 4} = \frac{x^2 - 1}{-x + 4}$

e. $-f(x) = -\left(\frac{x^2 - 1}{x + 4}\right) = \frac{-x^2 + 1}{x + 4}$

f. $f(x+1) = \frac{(x+1)^2 - 1}{(x+1) + 4}$
 $= \frac{x^2 + 2x + 1 - 1}{x + 5} = \frac{x^2 + 2x}{x + 5}$

g. $f(2x) = \frac{(2x)^2 - 1}{2x + 4} = \frac{4x^2 - 1}{2x + 4}$

h. $f(x+h) = \frac{(x+h)^2 - 1}{(x+h) + 4} = \frac{x^2 + 2xh + h^2 - 1}{x + h + 4}$

43. $f(x) = |x| + 4$

a. $f(0) = |0| + 4 = 0 + 4 = 4$

b. $f(1) = |1| + 4 = 1 + 4 = 5$

c. $f(-1) = |-1| + 4 = 1 + 4 = 5$

44. $f(x) = \sqrt{x^2 + x}$

a. $f(0) = \sqrt{0^2 + 0} = \sqrt{0} = 0$

b. $f(1) = \sqrt{1^2 + 1} = \sqrt{2}$

c. $f(-1) = \sqrt{(-1)^2 + (-1)} = \sqrt{1 - 1} = \sqrt{0} = 0$

d. $f(-x) = \sqrt{(-x)^2 + (-x)} = \sqrt{x^2 - x}$

e. $-f(x) = -\left(\sqrt{x^2 + x}\right) = -\sqrt{x^2 + x}$

f. $f(x+1) = \sqrt{(x+1)^2 + (x+1)}$
 $= \sqrt{x^2 + 2x + 1 + x + 1}$

$= \sqrt{x^2 + 3x + 2}$

g. $f(2x) = \sqrt{(2x)^2 + 2x} = \sqrt{4x^2 + 2x}$

h. $f(x+h) = \sqrt{(x+h)^2 + (x+h)}$
 $= \sqrt{x^2 + 2xh + h^2 + x + h}$

45. $f(x) = \frac{2x+1}{3x-5}$

a. $f(0) = \frac{2(0)+1}{3(0)-5} = \frac{0+1}{0-5} = \frac{1}{-5} = -\frac{1}{5}$

b. $f(1) = \frac{2(1)+1}{3(1)-5} = \frac{2+1}{3-5} = \frac{3}{-2} = -\frac{3}{2}$

c. $f(-1) = \frac{2(-1)+1}{3(-1)-5} = \frac{-2+1}{-3-5} = \frac{-1}{-8} = \frac{1}{8}$

2 -x + 1

$$\text{d. } f(-x) = |-x| + 4 = |x| + 4$$

$$\text{e. } -f(x) = -(|x| + 4) = -|x| - 4$$

$$\text{f. } f(x+1) = |x+1| + 4$$

$$\text{g. } f(2x) = |2x| + 4 = 2|x| + 4$$

$$\text{h. } f(x+h) = |x+h| + 4$$

$$\text{d. } f(-x) = \frac{(-x)}{3(-x)-5} = \frac{-2x+1}{-3x-5} = \frac{2x-1}{3x+5}$$

$$\text{e. } -f(x) = -\left(\frac{2x+1}{3x-5}\right) = \frac{-2x-1}{3x-5}$$

$$\text{f. } f(x+1) = \frac{2(x+1)+1}{3(x+1)-5} = \frac{2x+2+1}{3x+3-5} = \frac{2x+3}{3x-2}$$

$$\text{g. } f(2x) = \frac{2(2x)+1}{3(2x)-5} = \frac{4x+1}{6x-5}$$

$$\text{h. } f(x+h) = \frac{2(x+h)+1}{3(x+h)-5} = \frac{2x+2h+1}{3x+3h-5}$$

46. $f(x) = 1 - \frac{1}{(x+2)^2}$

a. $f(0) = 1 - \frac{1}{(0+2)^2} = 1 - \frac{1}{4} = \frac{3}{4}$

b. $f(1) = 1 - \frac{1}{(1+2)^2} = 1 - \frac{1}{9} = \frac{8}{9}$

c. $f(-1) = 1 - \frac{1}{(-1+2)^2} = 1 - \frac{1}{1} = 0$

d. $f(-x) = 1 - \frac{1}{(-x+2)^2} = 1 - \frac{1}{(2-x)^2}$

e. $-f(x) = -\left(1 - \frac{1}{(x+2)^2}\right) = -1 + \frac{1}{(x+2)^2}$

f. $f(x+1) = 1 - \frac{1}{(x+1+2)^2} = 1 - \frac{1}{(x+3)^2}$

g. $f(2x) = 1 - \frac{1}{(2x+2)^2} = 1 - \frac{1}{4(x+1)^2}$

h. $f(x+h) = 1 - \frac{1}{(x+h+2)^2}$

47. $f(x) = -5x + 4$
 Domain: $\{x \mid x \text{ is any real number}\}$

48. $f(x) = x^2 + 2$
 Domain: $\{x \mid x \text{ is any real number}\}$

49.

52. $h(x) = \frac{2x}{x^2 - 4}$
 $x^2 - 4 \neq 0$
 $x^2 \neq 4 \Rightarrow x \neq \pm 2$
 Domain: $\{x \mid x \neq -2, x \neq 2\}$

53. $F(x) = \frac{x-2}{x^3 + x}$
 $x^3 + x \neq 0$
 $x(x^2 + 1) \neq 0$
 $x \neq 0, x^2 \neq -1$
 Domain: $\{x \mid x \neq 0\}$

54. $G(x) = \frac{x+4}{x^3 - 4x}$
 $x^3 - 4x \neq 0$
 $x(x^2 - 4) \neq 0$
 $x \neq 0, x^2 \neq 4$
 $x \neq 0, x \neq \pm 2$
 Domain: $\{x \mid x \neq -2, x \neq 0, x \neq 2\}$

55. $h(x) = 3x - 12$
 $3x - 12 \geq 0$
 $3x \geq 12$
 $x \geq 4$
 Domain: $\{x \mid x \geq 4\}$

56. $G(x) = \sqrt{1-x}$
 $1-x \geq 0$

5
0.

Chapter 2: Functions and Their Graphs

51. $f(x) = \frac{x}{x^2 + 1}$

Domain: $\{x \mid x \text{ is any real number}\}$

$$f(x) = \frac{x^2}{x^2 + 1}$$

Domain: $\{x \mid x \text{ is any real number}\}$

$$g(x) = \frac{x}{x^2 - 16}$$

$$x^2 - 16 \neq 0$$

$$x^2 \neq 16 \Rightarrow x \neq \pm 4$$

Domain: $\{x \mid x \neq -4, x \neq 4\}$

Section 2.1: Functions

$$-x \geq -1$$

$$x \leq 1$$

Domain: $\{x \mid x \leq 1\}$

57. $f(x) = \frac{4}{\sqrt{x-9}}$

$$x - 9 > 0$$

$$x > 9$$

Domain: $\{x \mid x > 9\}$

58. $f(x) = \frac{x}{\sqrt{x-4}}$
 $x-4 > 0$
 $x > 4$

Domain: $\{x \mid x > 4\}$

59. $p(x) = \sqrt{\frac{2}{x-1}} = \frac{\sqrt{2}}{\sqrt{x-1}}$

$x-1 > 0$

$x > 1$

Domain: $\{x \mid x > 1\}$

60. $q(x) = \sqrt{-x-2}$
 $-x-2 \geq 0$

$-x \geq 2$

$x \leq -2$

Domain: $\{x \mid x \leq -2\}$

61. $P(t) = \frac{\sqrt{t-4}}{3t-21}$

$t-4 \geq 0$

$t \geq 4$

Also $3t-21 \neq 0$

$3t-21 \neq 0$

$3t \neq 21$

$t \neq 7$

Domain: $\{t \mid t \geq 4, t \neq 7\}$

62. $h(z) = \frac{\sqrt{z+3}}{z-2}$

$z+3 \geq 0$

$z \geq -3$

Also $z-2 \neq 0$

$z \neq 2$

b. $(f-g)(x) = (3x+4) - (2x-3)$
 $= 3x+4-2x+3$
 $= x+7$

Domain: $\{x \mid x \text{ is any real number}\}$.

c. $(f \cdot g)(x) = (3x+4)(2x-3)$
 $= 6x^2 - 9x + 8x - 12$
 $= 6x^2 - x - 12$

Domain: $\{x \mid x \text{ is any real number}\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{3x+4}{2x-3}$

$2x-3 \neq 0 \Rightarrow 2x \neq 3 \Rightarrow x \neq \frac{3}{2}$

Domain: $\left\{x \mid x \neq \frac{3}{2}\right\}$.

e. $(f+g)(3) = 5(3)+1 = 15+1 = 16$

f. $(f-g)(4) = 4+7 = 11$

g. $(f \cdot g)(2) = 6(2)^2 - 2 - 12 = 24 - 2 - 12 = 10$

h. $\left(\frac{f}{g}\right)(1) = \frac{3(1)+4}{2(1)-3} = \frac{3+4}{2-3} = \frac{7}{-1} = -7$

64. $f(x) = 2x+1$ $g(x) = 3x-2$

a. $(f+g)(x) = 2x+1+3x-2 = 5x-1$

Domain: $\{x \mid x \text{ is any real number}\}$.

b. $(f-g)(x) = (2x+1) - (3x-2)$
 $= 2x+1-3x+2$
 $= -x+3$

Domain: $\{x \mid x \text{ is any real number}\}$.

c. $(f \cdot g)(x) = (2x+1)(3x-2)$
 $= 6x^2 - 4x + 3x - 2$
 $= 6x^2 - x - 2$

Domain: $\{x \mid x \text{ is any real number}\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{2x+1}{3x-2}$

Chapter 2: Functions and Their Graphs

Section 2.1: Functions

Domain: $\{z \mid z \geq -3, z \neq 2\}$

63. $f(x) = 3x + 4$ $g(x) = 2x - 3$

a. $(f + g)(x) = 3x + 4 + 2x - 3 = 5x + 1$

Domain: $\{x \mid x \text{ is any real number}\}$.

$$\left(\begin{array}{l} g \end{array} \right) \quad 3x - 2 \\ 3x - 2 \neq 0$$

$$3x \neq 2 \Rightarrow x \neq \frac{2}{3}$$

Domain: $\left\{ x \mid x \neq \frac{2}{3} \right\}$.

(3)

e. $(f + g)(3) = 5(3) - 1 = 15 - 1 = 14$

f. $(f - g)(4) = -4 + 3 = -1$

g. $(f \cdot g)(2) = 6(2)^2 - 2 - 2$
 $= 6(4) - 2 - 2$
 $= 24 - 2 - 2 = 20$

h. $\left(\frac{f}{g}\right)(1) = \frac{2(1)+1}{3(1)-2} = \frac{2+1}{3-2} = \frac{3}{1} = 3$

65. $f(x) = x - 1$ $g(x) = 2x^2$

a. $(f + g)(x) = x - 1 + 2x^2 = 2x^2 + x - 1$
 Domain: $\{x \mid x \text{ is any real number}\}$.

b. $(f - g)(x) = (x - 1) - (2x^2)$
 $= x - 1 - 2x^2$
 $= -2x^2 + x - 1$

Domain: $\{x \mid x \text{ is any real number}\}$.

c. $(f \cdot g)(x) = (x - 1)(2x^2) = 2x^3 - 2x^2$
 Domain: $\{x \mid x \text{ is any real number}\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{x-1}{2x^2}$

Domain: $\{x \mid x \neq 0\}$.

e. $(f + g)(3) = 2(3)^2 + 3 - 1$
 $= 2(9) + 3 - 1$
 $= 18 + 3 - 1 = 20$

f. $(f - g)(4) = -2(4)^2 + 4 - 1$
 $= -2(16) + 4 - 1$
 $= -32 + 4 - 1 = -29$

66. $f(x) = 2x^2 + 3$ $g(x) = 4x^3 + 1$

a. $(f + g)(x) = 2x^2 + 3 + 4x^3 + 1$
 $= 4x^3 + 2x^2 + 4$
 Domain: $\{x \mid x \text{ is any real number}\}$.

b. $(f - g)(x) = (2x^2 + 3) - (4x^3 + 1)$
 $= 2x^2 + 3 - 4x^3 - 1$
 $= -4x^3 + 2x^2 + 2$
 Domain: $\{x \mid x \text{ is any real number}\}$.

c. $(f \cdot g)(x) = (2x^2 + 3)(4x^3 + 1)$
 $= 8x^5 + 12x^3 + 2x^2 + 3$
 Domain: $\{x \mid x \text{ is any real number}\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{2x^2 + 3}{4x^3 + 1}$ $\sqrt{\quad}$
 $4x^3 + 1 \neq 0$
 $4x^3 \neq -1$

$$x^3 \neq -\frac{1}{4} \Rightarrow x \neq \sqrt[3]{-\frac{1}{4}} = -\frac{\sqrt[3]{2}}{2}$$

Domain: $\left\{x \mid x \neq -\frac{\sqrt[3]{2}}{2}\right\}$.

g. $(f \cdot g)(2) = 2(2)^3 - 2(2)^2$
 $= 2(8) - 2(4)$
 $= 16 - 8 = 8$

Chapter 2: Functions and Their Graphs

Section 2.1: Functions

e. $(f + g)(3) = 4(3)^3 + 2(3)^2 + 4$
 $= 4(27) + 2(9) + 4$
 $= 108 + 18 + 4 = 130$

f. $(f - g)(4) = -4(4)^3 + 2(4)^2 + 2$

h. $\left(\frac{f}{g}\right)(1) = \frac{1-1}{2(1)^2} = \frac{0}{2(1)} = \frac{0}{2} = 0$

$$= -4(64) + 2(16) + 2$$

$$= -256 + 32 + 2 = -222$$

g. $(f \cdot g)(2) = 8(2)^5 + 12(2)^3 + 2(2)^2 + 3$
 $= 8(32) + 12(8) + 2(4) + 3$
 $= 256 + 96 + 8 + 3 = 363$

h. $\left(\frac{f}{g}\right)(1) = \frac{2(1)^2 + 3}{4(1)^3 + 1} = \frac{2(1) + 3}{4(1) + 1} = \frac{2 + 3}{4 + 1} = \frac{5}{5} = 1$

67. $f(x) = \sqrt{x}$ $g(x) = 3x - 5$

a. $(f + g)(x) = \sqrt{x} + 3x - 5$

Domain: $\{x \mid x \geq 0\}$.

b. $(f - g)(x) = \sqrt{x} - (3x - 5) = \sqrt{x} - 3x + 5$

Domain: $\{x \mid x \geq 0\}$.

c. $(f \cdot g)(x) = \sqrt{x}(3x - 5) = 3x\sqrt{x} - 5\sqrt{x}$

Domain: $\{x \mid x \geq 0\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{\sqrt{x}}{3x - 5}$

$x \geq 0$ and $3x - 5 \neq 0$

$3x \neq 5 \Rightarrow x \neq \frac{5}{3}$

Domain: $\left\{x \mid x \geq 0 \text{ and } x \neq \frac{5}{3}\right\}$.

e. $(f + g)(3) = \sqrt{3} + 3(3) - 5$

$= \sqrt{3} + 9 - 5 = \sqrt{3} + 4$

f. $(f - g)(4) = \sqrt{4} - 3(4) + 5$

$= 2 - 12 + 5 = -5$

g. $(f \cdot g)(2) = 3(2)\sqrt{2} - 5\sqrt{2}$

$= 6\sqrt{2} - 5\sqrt{2} = \sqrt{2}$

h. $\left(\frac{f}{g}\right)(1) = \frac{\sqrt{1}}{3(1) - 5} = \frac{1}{3 - 5} = \frac{1}{-2} = -\frac{1}{2}$

68. $f(x) = |x| \quad g(x) = x$

a. $(f + g)(x) = |x| + x$

Domain: $\{x \mid x \text{ is any real number}\}$.

b. $(f - g)(x) = |x| - x$

h. $\left(\frac{f}{g}\right)(1) = \frac{1}{1} = \frac{1}{1} = 1$

69. $f(x) = 1 + \frac{1}{x} \quad g(x) = \frac{1}{x}$

a. $(f + g)(x) = 1 + \frac{1}{x} + \frac{1}{x} = 1 + \frac{2}{x}$

Domain: $\{x \mid x \neq 0\}$.

b. $(f - g)(x) = 1 + \frac{1}{x} - \frac{1}{x} = 1$

Domain: $\{x \mid x \neq 0\}$.

c. $(f \cdot g)(x) = \left(1 + \frac{1}{x}\right)\frac{1}{x} = \frac{1}{x} + \frac{1}{x^2}$

Domain: $\{x \mid x \neq 0\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{1 + \frac{1}{x}}{\frac{1}{x}} = \frac{x + 1}{1} = x + 1$

Domain: $\{x \mid x \neq 0\}$.

e. $(f + g)(3) = 1 + \frac{2}{3} = \frac{5}{3}$

f. $(f - g)(4) = 1$

g. $(f \cdot g)(2) = \frac{1}{2} + \frac{1}{(2)^2} = \frac{1}{2} + \frac{1}{4} = \frac{3}{4}$

h. $\left(\frac{f}{g}\right)(1) = 1 + 1 = 2$

70. $f(x) = \sqrt{x-1} \quad g(x) = \sqrt{4-x}$

Chapter 2: Functions and Their Graphs

Domain: $\{x \mid x \text{ is any real number}\}$.

c. $(f \cdot g)(x) = |x| \cdot x = x|x|$

Domain: $\{x \mid x \text{ is any real number}\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{|x|}{x}$

Domain: $\{x \mid x \neq 0\}$.

e. $(f + g)(3) = |3| + 3 = 3 + 3 = 6$

f. $(f - g)(4) = |4| - 4 = 4 - 4 = 0$

g. $(f \cdot g)(2) = 2|2| = 2 \cdot 2 = 4$

Section 2.1: Functions

a. $(f + g)(x) = \sqrt{x-1} + \sqrt{4-x}$

$$x - 1 \geq 0 \text{ and } 4 - x \geq 0$$

$$x \geq 1 \text{ and } -x \geq -4$$

$$x \leq 4$$

Domain: $\{x \mid 1 \leq x \leq 4\}$.

b. $(f - g)(x) = \sqrt{x-1} - \sqrt{4-x}$

$$x - 1 \geq 0 \text{ and } 4 - x \geq 0$$

$$x \geq 1 \text{ and } -x \geq -4$$

$$x \leq 4$$

Domain: $\{x \mid 1 \leq x \leq 4\}$.

c. $(f \cdot g)(x) = (\sqrt{x-1})(\sqrt{4-x})$
 $= \sqrt{-x^2 + 5x - 4}$
 $x-1 \geq 0$ and $4-x \geq 0$

$x \geq 1$ and $-x \geq -4$
 $x \leq 4$
 Domain: $\{x \mid 1 \leq x \leq 4\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{\sqrt{x-1}}{\sqrt{4-x}} = \sqrt{\frac{x-1}{4-x}}$
 $x-1 \geq 0$ and $4-x > 0$
 $x \geq 1$ and $-x > -4$
 $x < 4$

Domain: $\{x \mid 1 \leq x < 4\}$.

e. $(f + g)(3) = \sqrt{3-1} + 4-3$
 $= \sqrt{2} + \sqrt{1} = \sqrt{2} + 1$

f. $(f - g)(4) = \sqrt{4-1} - \sqrt{4-4}$
 $= \sqrt{3} - \sqrt{0} = \sqrt{3} - 0 = \sqrt{3}$

g. $(f \cdot g)(2) = \sqrt{-(2)^2 + 5(2) - 4}$
 $= \sqrt{-4 + 10 - 4} = \sqrt{2}$

h. $\left(\frac{f}{g}\right)(1) = \frac{\sqrt{1-1}}{4-1} = \frac{\sqrt{0}}{3} = \sqrt{0} = 0$

71. $f(x) = \frac{2x+3}{3x-2}$ $g(x) = \frac{4x}{3x-2}$

a. $(f + g)(x) = \frac{2x+3}{3x-2} + \frac{4x}{3x-2}$
 $3x-2 \neq 0$

b. $(f - g)(x) = \frac{2x+3}{3x-2} - \frac{4x}{3x-2}$
 $= \frac{2x+3-4x}{3x-2} = \frac{-2x+3}{3x-2}$

$3x-2 \neq 0$
 $3x \neq 2 \Rightarrow x \neq \frac{2}{3}$

Domain: $\left\{x \mid x \neq \frac{2}{3}\right\}$.

c. $(f \cdot g)(x) = \left(\frac{2x+3}{3x-2}\right)\left(\frac{4x}{3x-2}\right) = \frac{8x+12x}{(3x-2)^2}$
 $3x-2 \neq 0$

$3x \neq 2 \Rightarrow x \neq \frac{2}{3}$

Domain: $\left\{x \mid x \neq \frac{2}{3}\right\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{\frac{2x+3}{3x-2}}{\frac{4x}{3x-2}} = \frac{2x+3}{4x} \cdot \frac{3x-2}{4x} = \frac{2x+3}{4x}$

$3x-2 \neq 0$ and $x \neq 0$

$3x \neq 2$
 $x \neq \frac{2}{3}$

Domain: $\left\{x \mid x \neq \frac{2}{3} \text{ and } x \neq 0\right\}$.

e. $(f + g)(3) = \frac{6(3)+3}{3(3)-2} = \frac{18+3}{9-2} = \frac{21}{7} = 3$
 $= \frac{2x+3+4x}{3x-2} = \frac{6x+3}{3x-2}$

Chapter 2: Functions and Their Graphs

Section 2.1: Functions

$$\frac{3x-2}{2} \quad 3x-$$

$$\mathbf{f.} \quad (f-g)(4) = \frac{2(4)+3}{2} = \frac{-8+3}{2} = \frac{5}{2} = -\frac{1}{2}$$

$$\frac{3(4)}{2} = \frac{12}{2} = 6$$

$$3x \neq 2 \Rightarrow x \neq \frac{2}{3}$$

$$\text{Domain: } \left\{ x \mid x \neq \frac{2}{3} \right\}$$

$$\mathbf{g.} \quad (f \cdot g)(2) = \frac{8(2) + 12(2)}{(3(2) - 2)^2}$$

$$= \frac{8(4) + 24}{(6-2)^2} = \frac{32+24}{(4)^2} = \frac{56}{16} = \frac{7}{2}$$

$$\mathbf{h.} \quad \left(\frac{f}{g} \right)(1) = \frac{2(1)+3}{4(1)} = \frac{2+3}{4} = \frac{5}{4}$$

72. $f(x) = \sqrt{x+1}$ $g(x) = \frac{2}{x}$

a. $(f + g)(x) = \sqrt{x+1} + \frac{2}{x}$

$x+1 \geq 0$ and $x \neq 0$
 $x \geq -1$

Domain: $\{x | x \geq -1, \text{ and } x \neq 0\}$.

b. $(f - g)(x) = \sqrt{x+1} - \frac{2}{x}$

$x+1 \geq 0$ and $x \neq 0$
 $x \geq -1$

Domain: $\{x | x \geq -1, \text{ and } x \neq 0\}$.

c. $(f \cdot g)(x) = \sqrt{x+1} \cdot \frac{2}{x} = \frac{2\sqrt{x+1}}{x}$

$x+1 \geq 0$ and $x \neq 0$
 $x \geq -1$

Domain: $\{x | x \geq -1, \text{ and } x \neq 0\}$.

d. $\left(\frac{f}{g}\right)(x) = \frac{\sqrt{x+1}}{\frac{2}{x}} = \frac{x\sqrt{x+1}}{2}$

$x+1 \geq 0$ and $x \neq 0$
 $x \geq -1$

Domain: $\{x | x \geq -1, \text{ and } x \neq 0\}$.

e. $(f + g)(3) = \sqrt{3+1} + \frac{2}{3} = \sqrt{4} + \frac{2}{3} = 2 + \frac{2}{3} = \frac{8}{3}$

f. $(f - g)(4) = \sqrt{4+1} - \frac{2}{4} = \sqrt{5} - \frac{1}{2}$

74. $f(x) = \frac{1}{x}$ $\left(\frac{f}{g}\right)(x) = \frac{x+1}{x^2 - x}$

$\frac{x+1}{x^2 - x} = \frac{x+1}{x(x-1)}$

$g(x) = \frac{1}{x} = \frac{1}{x} \cdot \frac{x+1}{x+1}$

$\frac{x+1}{x(x-1)}$

$= \frac{1}{x} \cdot \frac{x(x-1)}{x+1} = \frac{x-1}{x+1}$

75. $f(x) = 4x+3$

$\frac{f(x+h) - f(x)}{h} = \frac{4(x+h)+3 - (4x+3)}{h}$

$= \frac{4x+4h+3-4x-3}{h}$

$= \frac{4h}{h} = 4$

h

76. $f(x) = -3x+1$

$\frac{f(x+h) - f(x)}{h} = \frac{-3(x+h)+1 - (-3x+1)}{h}$

$= \frac{-3x-3h+1+3x-1}{h}$

$= \frac{-3h}{h} = -3$

77. $f(x) = x^2 - x + 4$

$\frac{f(x+h) - f(x)}{h}$

$\frac{(x+h)^2 - (x+h) + 4 - (x^2 - x + 4)}{h}$

Chapter 2: Functions and Their Graphs

g. $(f \cdot g)(2) = \frac{2\sqrt{2+1}}{2} = \frac{2\sqrt{3}}{2} = \sqrt{3}$

h. $\begin{pmatrix} f \\ g \end{pmatrix}(1) = \frac{1\sqrt{1+1}}{2} = \frac{\sqrt{2}}{2}$

73. $f(x) = 3x + 1 \quad (f + g)(x) = 6 - \frac{1}{2}x$

$$6 - \frac{1}{2}x = 3x + 1 + g(x)$$

$$5 - \frac{7}{2}x = g(x)$$

$$g(x) = 5 - \frac{7}{2}x$$

Section 2.1: Functions

$$\begin{aligned} &= \frac{h}{h} \\ &= \frac{x^2 + 2xh + h^2 - x - h + 4 - x^2 + x - 4}{h} \\ &= \frac{2xh + h^2 - h}{h} \\ &= 2x + h - 1 \end{aligned}$$

$$\begin{aligned}
 78. \quad f(x) &= 3x^2 - 2x + 6 \\
 \frac{f(x+h) - f(x)}{h} &= \frac{[3(x+h)^2 - 2(x+h) + 6] - [3x^2 - 2x + 6]}{h} \\
 &= \frac{3(x^2 + 2xh + h^2) - 2x - 2h + 6 - 3x^2 + 2x - 6}{h} \\
 &= \frac{3x^2 + 6xh + 3h^2 - 2h - 3x^2}{h} = \frac{6xh + 3h^2 - 2h}{h} \\
 &= 6x + 3h - 2
 \end{aligned}$$

$$\begin{aligned}
 79. \quad f(x) &= \frac{1}{x^2} \\
 \frac{f(x+h) - f(x)}{h} &= \frac{\frac{1}{(x+h)^2} - \frac{1}{x^2}}{h} \\
 &= \frac{\frac{x^2 - (x+h)^2}{x^2(x+h)^2}}{h} \\
 &= \frac{x - (x^2 + 2xh + h^2)}{x^2(x+h)^2} \\
 &= \frac{-2xh - h^2}{x^2(x+h)^2} \\
 &= \frac{h(-2x - h)}{x^2(x+h)^2} \\
 &= \frac{-2x - h}{x^2(x+h)^2} = -\frac{2x+h}{x^2(x+h)^2}
 \end{aligned}$$

$$\begin{aligned}
 80. \quad f(x) &= \frac{1}{x+3} \\
 \frac{f(x+h) - f(x)}{h} &= \frac{\frac{1}{x+h+3} - \frac{1}{x+3}}{h} \\
 &= \frac{(x+3) - (x+h+3)}{h(x+3)(x+h+3)} \\
 &= \frac{-h}{h(x+3)(x+h+3)} \\
 &= \frac{-1}{(x+h+3)(x+3)}
 \end{aligned}$$

$$\begin{aligned}
 81. \quad f(x) &= \sqrt{x} \\
 \frac{f(x+h) - f(x)}{h} &= \frac{\sqrt{x+h} - \sqrt{x}}{h} \\
 &= \frac{\sqrt{x+h} - \sqrt{x}}{h} \cdot \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\
 &= \frac{\sqrt{x+h} - x\sqrt{x}}{h(x+h+x)} = \frac{h}{h(x+h+\sqrt{x})} \\
 &= \frac{1}{\sqrt{x+h} + \sqrt{x}}
 \end{aligned}$$

$$\begin{aligned}
 82. \quad f(x) &= \sqrt{x+1} \\
 \frac{f(x+h) - f(x)}{h} &= \frac{\sqrt{x+h+1} - \sqrt{x+1}}{h}
 \end{aligned}$$

Chapter 2: Functions and Their Graphs

Section 2.1: Functions

$$= \frac{x+h+1 - x+1}{h}$$

$$= \frac{\sqrt{x+h+1} - \sqrt{x+1}}{h} \cdot \frac{\sqrt{x+h+1} + \sqrt{x+1}}{\sqrt{x+h+1} + \sqrt{x+1}}$$

$$= \frac{x+h+1 - (x+1)}{h(\sqrt{x+h+1} + \sqrt{x+1})} = \frac{h}{h(\sqrt{x+h+1} + \sqrt{x+1})}$$

$$= \frac{1}{\sqrt{x+h+1} + \sqrt{x+1}}$$

83. $f(x) = 2x^3 + Ax^2 + 4x - 5$ and $f(2) = 5$

$$f(2) = 2(2)^3 + A(2)^2 + 4(2) - 5$$

$$5 = 16 + 4A + 8 - 5$$

$$5 = 4A + 19$$

$$-14 = 4A$$

$$A = \frac{-14}{4} = -\frac{7}{2}$$

84. $f(x) = 3x^2 - Bx + 4$ and $f(-1) = 12$:

$$f(-1) = 3(-1)^2 - B(-1) + 4$$

$$12 = 3 + B + 4$$

$$B = 5$$

85. $f(x) = \frac{3x+8}{2x-A}$ and $f(0) = 2$

$$f(0) = \frac{3(0)+8}{2(0)-A}$$

$$2 = \frac{8}{-A}$$

$$-2A = 8$$

$$A = -4$$

86. $f(x) = \frac{2x-B}{3x+4}$ and $f(2) = \frac{1}{2}$

$$f(2) = \frac{2(2)-B}{3(2)+4} = \frac{1}{2}$$

$$\frac{1}{2} = \frac{4-B}{10}$$

$$5 = 4 - B$$

$$B = -1$$

87. $f(x) = \frac{2x-A}{x-3}$ and $f(4) = 0$

$$f(4) = \frac{2(4)-A}{4-3} = 0$$

$$0 = \frac{8-A}{1}$$

$$0 = 8 - A$$

88. $f(x) = \frac{x-B}{x-A}$, $f(2) = 0$ and $f(1)$ is undefined

$$1 - A = 0 \Rightarrow A = 1$$

$$f(2) = \frac{2-B}{2-1} = 0$$

$$0 = \frac{2-B}{1}$$

$$0 = 2 - B$$

$$B = 2$$

89. Let x represent the length of the rectangle.

Then, $\frac{x}{2}$ represents the width of the rectangle

since the length is twice the width. The function

for the area is: $A(x) = x \cdot \frac{x}{2} = \frac{x^2}{2} = \frac{1}{2}x^2$

90. Let x represent the length of one of the two equal sides. The function for the area is:

$$A(x) = \frac{1}{2} \cdot x \cdot x = \frac{1}{2}x^2$$

91. Let x represent the number of hours worked. The function for the gross salary is:

$$G(x) = 10x$$

92. Let x represent the number of items sold. The function for the gross salary is:

$$G(x) = 10x + 100$$

93. a. P is the dependent variable; a is the independent variable

$$\begin{aligned} \text{b. } P(20) &= 0.015(20)^2 - 4.962(20) + 290.580 \\ &= 6 - 99.24 + 290.580 \\ &= 197.34 \end{aligned}$$

In 2005 there are 197.34 million people who are 20 years of age or older.

$$\begin{aligned} \text{c. } P(0) &= 0.015(0)^2 - 4.962(0) + 290.580 \\ &= 290.580 \end{aligned}$$

In 2005 there are 290.580 million people.

Chapter 2: Functions and Their Graphs

$A = 8$
 f is undefined when $x = 3$.

Section 2.1: Functions

94. a. N is the dependent variable; r is the independent variable

b.
$$\begin{aligned} N(3) &= -1.44(3)^2 + 14.52(3) - 14.96 \\ &= -12.96 + 43.56 - 14.96 \\ &= 15.64 \end{aligned}$$

In 2005, there are 15.64 million housing units with 3 rooms.

95. a. $H(1) = 20 - 4.9(1)^2$
 $= 20 - 4.9 = 15.1$ meters
 $H(1.1) = 20 - 4.9(1.1)^2$
 $= 20 - 4.9(1.21)$
 $= 20 - 5.929 = 14.071$ meters
 $H(1.2) = 20 - 4.9(1.2)^2$
 $= 20 - 4.9(1.44)$
 $= 20 - 7.056 = 12.944$ meters
 $H(1.3) = 20 - 4.9(1.3)^2$
 $= 20 - 4.9(1.69)$
 $= 20 - 8.281 = 11.719$ meters

b. $H(x) = 15$:
 $15 = 20 - 4.9x^2$
 $-5 = -4.9x^2$
 $x^2 \approx 1.0204$
 $x \approx 1.01$ seconds

$H(x) = 10$:
 $10 = 20 - 4.9x^2$
 $-10 = -4.9x^2$
 $x^2 \approx 2.0408$
 $x \approx 1.43$ seconds

$H(x) = 5$:
 $5 = 20 - 4.9x^2$
 $-15 = -4.9x^2$
 $x^2 \approx 3.0612$
 $x \approx 1.75$ seconds

c. $H(x) = 0$
 $0 = 20 - 4.9x^2$

b. $H(x) = 15$
 $15 = 20 - 13x^2$
 $-5 = -13x^2$
 $x^2 \approx 0.3846$
 $x \approx 0.62$ seconds

$H(x) = 10$
 $10 = 20 - 13x^2$
 $-10 = -13x^2$
 $x^2 \approx 0.7692$
 $x \approx 0.88$ seconds

$H(x) = 5$
 $5 = 20 - 13x^2$
 $-15 = -13x^2$
 $x^2 \approx 1.1538$
 $x \approx 1.07$ seconds

c. $H(x) = 0$
 $0 = 20 - 13x^2$
 $-20 = -13x^2$
 $x^2 \approx 1.5385$
 $x \approx 1.24$ seconds

97. $C(x) = 100 + \frac{x}{10} + \frac{36,000}{x}$

a. $C(500) = 100 + \frac{500}{10} + \frac{36,000}{500}$
 $= 100 + 50 + 72$
 $= \$222$

$\frac{450}{10} + \frac{36,000}{450}$

b. $C(450) = 100 + \frac{450}{10} + \frac{36,000}{450}$

Chapter 2: Functions and Their Graphs**Section 2.1: Functions**

96. a. $-20 = -4.9x^2$

$x^2 \approx 4.0816$

$x \approx 2.02$ seconds

$H(1) = 20 - 13(1)^2 = 20 - 13 = 7$ meters

$H(1.1) = 20 - 13(1.1)^2 = 20 - 13(1.21)$

$= 20 - 15.73 = 4.27$ meters

$H(1.2) = 20 - 13(1.2)^2 = 20 - 13(1.44)$

$= 20 - 18.72 = 1.28$ meters

$= 100 + 45 + 80$

$= \$225$

c. $C(600) = 100 + \frac{600}{10} + \frac{36,000}{600}$

$= 100 + 60 + 60$

$= \$220$

d. $C(400) = 100 + \frac{400}{10} + \frac{36,000}{400}$

$= 100 + 40 + 90$

$= \$230$

98. $A(x) = 4x\sqrt{1-x^2}$

a. $A\left(\frac{1}{3}\right) = 4 \cdot \frac{1}{3} \sqrt{1 - \left(\frac{1}{3}\right)^2} = \frac{4}{3} \sqrt{\frac{8}{9}} = \frac{4}{3} \cdot \frac{2\sqrt{2}}{3}$
 $= \frac{8\sqrt{2}}{9} \approx 1.26 \text{ ft}^2$

b. $A\left(\frac{1}{2}\right) = 4 \cdot \frac{1}{2} \sqrt{1 - \left(\frac{1}{2}\right)^2} = 2\sqrt{\frac{3}{4}} = 2 \cdot \frac{\sqrt{3}}{2}$
 $= \sqrt{3} \approx 1.73 \text{ ft}^2$

c. $A\left(\frac{2}{3}\right) = 4 \cdot \frac{2}{3} \sqrt{1 - \left(\frac{2}{3}\right)^2} = \frac{8}{3} \sqrt{\frac{5}{9}} = \frac{8}{3} \cdot \frac{\sqrt{5}}{3}$
 $= \frac{8\sqrt{5}}{9} \approx 1.99 \text{ ft}^2$

99. $R(x) = \left(\frac{L}{P}\right)(x) = \frac{L(x)}{P(x)}$

100. $T(x) = (V + P)(x) = V(x) + P(x)$

101. $H(x) = (P \cdot I)(x) = P(x) \cdot I(x)$

102. $N(x) = (I - T)(x) = I(x) - T(x)$

103. a. $P(x) = R(x) - C(x)$

$$= (-1.2x^2 + 220x) - (0.05x^3 - 2x^2 + 65x + 500)$$

$$= -1.2x^2 + 220x - 0.05x^3 + 2x^2 - 65x - 500$$

$$= -0.05x^3 + 0.8x^2 + 155x - 500$$

b. $P(15) = -0.05(15)^3 + 0.8(15)^2 + 155(15) - 500$
 $= -168.75 + 180 + 2325 - 500$
 $= \$1836.25$

c. When 30 clocks are sold, the profit is \$200.

105. a. $h(x) = 2x$

$$h(a+b) = 2(a+b) = 2a + 2b$$

$$= h(a) + h(b)$$

$h(x) = 2x$ has the property.

b. $g(x) = x^2$

$$g(a+b) = (a+b)^2 = a^2 + 2ab + b^2$$

Since

$$a^2 + 2ab + b^2 \neq a^2 + b^2 = g(a) + g(b),$$

$g(x) = x^2$ does not have the property.

c. $F(x) = 5x - 2$

$$F(a+b) = 5(a+b) - 2 = 5a + 5b - 2$$

Since

$$5a + 5b - 2 \neq 5a - 2 + 5b - 2 = F(a) + F(b),$$

$F(x) = 5x - 2$ does not have the property.

d. $G(x) = \frac{1}{x}$

$$G(a+b) = \frac{1}{a+b} \neq \frac{1}{a} + \frac{1}{b} = G(a) + G(b)$$

$G(x) = \frac{1}{x}$ does not have the property.

106. No. The domain of f is $\{x \mid x \text{ is any real number}\}$, but the domain of g is $\{x \mid x \neq -1\}$.

107. Answers will vary.

$$\frac{3x - x^3}{c}$$

When 15 hundred cell phones are sold, the profit is \$1836.25.

108.

(
you
r
age
)

104. a. $P(x) = R(x) - C(x)$
 $= 30x - (0.1x^2 + 7x + 400)$
 $= 30x - 0.1x^2 - 7x - 400$
 $= -0.1x^2 + 23x - 400$
- b. $P(30) = -0.1(30)^2 + 23(30) - 400$
 $= -90 + 690 - 400$
 $= \$200$

Section 2.2

1. $x^2 + 4y^2 = 16$

x-intercepts:

$$x^2 + 4(0)^2 = 16$$

$$x^2 = 16$$

$$x = \pm 4 \Rightarrow (-4, 0), (4, 0)$$

y-intercepts:

$$(0)^2 + 4y^2 = 16$$

$$4y^2 = 16$$

$$y^2 = 4$$

$$y = \pm 2 \Rightarrow (0, -2), (0, 2)$$

2. False; $x = 2y - 2$

$$-2 = 2y - 2$$

$$0 = 2y$$

$$0 = y$$

The point $(-2, 0)$ is on the graph.

3. vertical

4. $f(5) = -3$

5. $f(x) = ax^2 + 4$

$$a(-1)^2 + 4 = 2 \Rightarrow a = -2$$

6. False; it would fail the vertical line test.

7. False; e.g. $y = \frac{1}{x}$.

8. True

9. a. $f(0) = 3$ since $(0, 3)$ is on the graph.

$$f(-6) = -3 \text{ since } (-6, -3) \text{ is on the graph.}$$

g. The domain of f is $\{x \mid -6 \leq x \leq 11\}$ or $[-6, 11]$.

h. The range of f is $\{y \mid -3 \leq y \leq 4\}$ or $[-3, 4]$.

i. The *x*-intercepts are $-3, 6,$ and 10 .

j. The *y*-intercept is 3 .

k. The line $y = \frac{1}{2}$ intersects the graph 3 times.

l. The line $x = 5$ intersects the graph 1 time.

m. $f(x) = 3$ when $x = 0$ and $x = 4$.

n. $f(x) = -2$ when $x = -5$ and $x = 8$.

10. a. $f(0) = 0$ since $(0, 0)$ is on the graph.
 $f(6) = 0$ since $(6, 0)$ is on the graph.

b. $f(2) = -2$ since $(2, -2)$ is on the graph.
 $f(-2) = 1$ since $(-2, 1)$ is on the graph.

c. $f(3)$ is negative since $f(3) \approx -1$.

d. $f(-1)$ is positive since $f(-1) \approx 1.0$.

e. $f(x) = 0$ when $x = 0, x = 4,$ and $x = 6$.

f. $f(x) < 0$ when $0 < x < 4$.

g. The domain of f is $\{x \mid -4 \leq x \leq 6\}$ or $[-4, 6]$.

h. The range of f is $\{y \mid -2 \leq y \leq 3\}$ or $[-2, 3]$.

i. The *x*-intercepts are $0, 4,$ and 6 .

j. The *y*-intercept is 0 .

k. The line $y = -1$ intersects the graph 2 times.

l. The line $x = 1$ intersects the graph 1 time.

Chapter 2: Functions and Their Graphs

- b.** $f(6) = 0$ since $(6, 0)$ is on the graph.
 $f(11) = 1$ since $(11, 1)$ is on the graph.
- c.** $f(3)$ is positive since $f(3) \approx 3.7$.
- d.** $f(-4)$ is negative since $f(-4) \approx -1$.
- e.** $f(x) = 0$ when $x = -3$, $x = 6$, and $x = 10$.
- f.** $f(x) > 0$ when $-3 < x < 6$, and $10 < x \leq 11$.

Section 2.2: The Graph of a Function

- m.** $f(x) = 3$ when $x = 5$.
- n.** $f(x) = -2$ when $x = 2$.
- 11.** Not a function since vertical lines will intersect the graph in more than one point.
- 12.** Function

a. Domain: $\{x \mid x \text{ is any real number}\}$;

Range: $\{y \mid y > 0\}$

b. Intercepts: (0,1)

c. None

13. Function

a. Domain: $\{x \mid -\pi \leq x \leq \pi\}$;

Range: $\{y \mid -1 \leq y \leq 1\}$

b. Intercepts: $\left(-\frac{\pi}{2}, 0\right), \left(\frac{\pi}{2}, 0\right), (0,1)$

c. Symmetry about y-axis.

14. Function

a. Domain: $\{x \mid -\pi \leq x \leq \pi\}$;

Range: $\{y \mid -1 \leq y \leq 1\}$

b. Intercepts: $(-\pi, 0), (\pi, 0), (0, 0)$

c. Symmetry about the origin.

15. Not a function since vertical lines will intersect the graph in more than one point.

16. Not a function since vertical lines will intersect the graph in more than one point.

17. Function

a. Domain: $\{x \mid 0 < x < 3\}$;

Range: $\{y \mid y < 2\}$

b. Intercepts: (1, 0)

c. None

18. Function

a. Domain: $\{x \mid 0 \leq x < 4\}$;

Range: $\{y \mid 0 \leq y < 3\}$

c. Symmetry about y-axis.

20. Function

a. Domain: $\{x \mid x \geq -3\}$;

Range: $\{y \mid y \geq 0\}$

b. Intercepts: $(-3, 0), (2,0), (0,2)$

c. None

21. Function

a. Domain: $\{x \mid x \text{ is any real number}\}$;

Range: $\{y \mid y \geq -3\}$

b. Intercepts: $(1, 0), (3,0), (0,9)$

c. None

22. Function

a. Domain: $\{x \mid x \text{ is any real number}\}$;

Range: $\{y \mid y \leq 5\}$

b. Intercepts: $(-1, 0), (2,0), (0,4)$

c. None

23. $f(x) = 2x^2 - x - 1$

a. $f(-1) = 2(-1)^2 - (-1) - 1 = 2$

The point $(-1, 2)$ is on the graph of f .

b. $f(-2) = 2(-2)^2 - (-2) - 1 = 9$

The point $(-2, 9)$ is on the graph of f .

c. Solve for x :

$$-1 = 2x^2 - x - 1$$

$$0 = 2x^2 - x$$

$$0 = x(2x - 1) \Rightarrow x = 0, x = \frac{1}{2}$$

$(0, -1)$ and $\left(\frac{1}{2}, -1\right)$ are on the graph of f .

Chapter 2: Functions and Their Graphs

- b. Intercepts: (0, 0)
- c. None

19. Function

- a. Domain: $\{x \mid x \text{ is any real number}\}$;
Range: $\{y \mid y \leq 2\}$
- b. Intercepts: $(-3, 0), (3, 0), (0, 2)$

Section 2.2: The Graph of a Function

- d. The domain of f is $\{x \mid x \text{ is any real number}\}$.

- e. x -intercepts:

$$f(x)=0 \Rightarrow 2x^2 - x - 1 = 0$$

$\frac{1}{2}$

$$(2x+1)(x-1) = 0 \Rightarrow x = -\frac{1}{2}, x = 1$$

$$\left(-\frac{1}{2}, 0\right) \text{ and } (1, 0)$$

$$\left(\frac{1}{2} \right)$$

f. y-intercept:

$$f(0) = 2(0)^2 - 0 - 1 = -1 \Rightarrow (0, -1)$$

24. $f(x) = -3x^2 + 5x$

a. $f(-1) = -3(-1)^2 + 5(-1) = -8 \neq 2$

The point $(-1, 2)$ is not on the graph of f .

b. $f(-2) = -3(-2)^2 + 5(-2) = -22$

The point $(-2, -22)$ is on the graph of f .

c. Solve for x :

$$-2 = -3x^2 + 5x \Rightarrow 3x^2 - 5x - 2 = 0$$

$$(3x+1)(x-2) = 0 \Rightarrow x = -\frac{1}{3}, x = 2$$

$(2, -2)$ and $(-\frac{1}{3}, -2)$ on the graph of f .

d. The domain of f is $\{x \mid x \text{ is any real number}\}$.

e. x-intercepts:

$$f(x) = 0 \Rightarrow -3x^2 + 5x = 0$$

$$x(-3x+5) = 0 \Rightarrow x = 0, x = \frac{5}{3}$$

$$(0, 0) \text{ and } (\frac{5}{3}, 0)$$

f. y-intercept:

$$f(0) = -3(0)^2 + 5(0) = 0 \Rightarrow (0, 0)$$

25. $f(x) = \frac{x+2}{x-6}$

a. $f(3) = \frac{3+2}{3-6} = -\frac{5}{3} \neq 14$

The point $(3, 14)$ is not on the graph of f .

b. $f(4) = \frac{4+2}{4-6} = \frac{6}{-2} = -3$

The point $(4, -3)$ is on the graph of f .

c. Solve for x :

e. x-intercepts:

$$f(x) = 0 \Rightarrow \frac{x+2}{x-6} = 0$$

$$x+2 = 0 \Rightarrow x = -2 \Rightarrow (-2, 0)$$

f. y-intercept: $f(0) = \frac{0+2}{0-6} = -\frac{1}{3} \Rightarrow (0, -\frac{1}{3})$

26. $f(x) = \frac{x^2+2}{x+4}$

a. $f(1) = \frac{1+2}{1+4} = \frac{3}{5}$

The point $(1, \frac{3}{5})$ is on the graph of f .

b. $f(0) = \frac{0^2+2}{0+4} = \frac{2}{4} = \frac{1}{2}$

The point $(0, \frac{1}{2})$ is on the graph of f .

c. Solve for x :

$$\frac{1}{2} = \frac{x^2+2}{x+4} \Rightarrow x+4 = 2x^2+4$$

$$2 = x^2 - x$$

$$0 = 2x^2 - x$$

$$x(2x-1) = 0 \Rightarrow x = 0 \text{ or } x = \frac{1}{2}$$

$(0, \frac{1}{2})$ and $(\frac{1}{2}, \frac{1}{2})$ are on the graph of f .

(\quad, \quad) (\quad, \quad)

d. The domain of f is $\{x \mid x \neq -4\}$.

e. x-intercepts:

$$f(x) = 0 \Rightarrow \frac{x^2+2}{x+4} = 0 \Rightarrow x^2+2 = 0$$

This is impossible, so there are no x-intercepts.

f. y-intercept:

$$\frac{0^2+2}{0+4} = \frac{2}{4} = \frac{1}{2} \Rightarrow (0, \frac{1}{2})$$

Chapter 2: Functions and Their Graphs

$$2 = \frac{x+2}{x-6}$$

$$2x - 12 = x + 2$$

$$x = 14$$

(14, 2) is a point on the graph of f .

d. The domain of f is $\{x \mid x \neq 6\}$.

Section 2.2: The Graph of a Function

$$f(0) = 0 + 4 = 4 = 2 \Rightarrow (0, 2)$$

27. $f(x) = \frac{2x^2}{x^4 + 1}$

$$x^4 + 1$$

$$\frac{2(-1)^2}{2} = 1$$

a. $f(-1) = \frac{2(-1)^2}{(-1)^4 + 1} = \frac{2}{2} = 1$

The point $(-1, 1)$ is on the graph of f .

b. $f(2) = \frac{2(2)^2}{(2)^4 + 1} = \frac{8}{17}$

The point $\left(2, \frac{8}{17}\right)$ is on the graph of f .

c. Solve for x :

$$1 = \frac{2x^2}{x^4 + 1}$$

$$x^4 + 1 = 2x^2$$

$$x^4 - 2x^2 + 1 = 0$$

$$(x^2 - 1)^2 = 0$$

$$x^2 - 1 = 0 \Rightarrow x = \pm 1$$

$(1, 1)$ and $(-1, 1)$ are on the graph of f .

d. The domain of f is $\{x \mid x \text{ is any real number}\}$.

e. x -intercept:

$$f(x) = 0 \Rightarrow \frac{2x^2}{x + 1} = 0$$

$$2x^2 = 0 \Rightarrow x = 0 \Rightarrow (0, 0)$$

f. y -intercept:

$$f(0) = \frac{2(0)^2}{0^4 + 1} = \frac{0}{0 + 1} = 0 \Rightarrow (0, 0)$$

28. $f(x) = \frac{2x}{x - 2}$

$$2\left(\frac{1}{2}\right)$$

a. $f\left(\frac{1}{2}\right) = \frac{2\left(\frac{1}{2}\right)}{\left(\frac{1}{2}\right) - 2} = \frac{1}{-\frac{3}{2}} = -\frac{2}{3}$

The point $\left(\frac{1}{2}, -\frac{2}{3}\right)$ is on the graph of f .

b. $f(4) = \frac{2(4)}{4 - 2} = \frac{8}{2} = 4$

The point $(4, 4)$ is on the graph of f .

e. x -intercept:

$$f(x) = 0 \Rightarrow \frac{2x}{x - 2} = 0 \Rightarrow 2x = 0$$

$$\Rightarrow x = 0 \Rightarrow (0, 0)$$

f. y -intercept: $f(0) = \frac{0}{0 - 2} = 0 \Rightarrow (0, 0)$

29. $h(x) = -\frac{44x^2}{v^2} + x + 6$

a. $h(8) = -\frac{44(8)^2}{28^2} + (8) + 6$

$$= -\frac{2816}{784} + 14$$

$$\approx 10.4 \text{ feet}$$

b. $h(12) = -\frac{44(12)^2}{28^2} + (12) + 6$

$$= -\frac{6336}{784} + 18$$

$$\approx 9.9 \text{ feet}$$

$h(12) \approx 9.9$ represents the height of the ball, in feet, after it has traveled 12 feet in front of the foul line.

c. From part (a) we know the point $(8, 10.4)$ is on the graph and from part (b) we know the point $(12, 9.9)$ is on the graph. We could

evaluate the function at several more values of x (e.g. $x = 0$, $x = 15$, and $x = 20$) to obtain additional points.

$$h(0) = -\frac{44(0)^2}{28^2} + (0) + 6 = 6$$

$$\frac{44(15)^2}{28^2}$$

Chapter 2: Functions and Their Graphs

c. Solve for x :

$$1 = \frac{2x}{x-2} \Rightarrow x-2 = 2x \Rightarrow -2 = x$$

$(-2,1)$ is a point on the graph of f .

d. The domain of f is $\{x \mid x \neq 2\}$.


Section 2.2: The Graph of a Function

$$h(15) = -\frac{28^2}{44(15)^2} + (15) + 6 \approx 8.4$$

$$h(20) = -\frac{28^2}{44(20)^2} + (20) + 6 \approx 3.6$$

Some additional points are $(0,6)$, $(15,8.4)$
and $(20,3.6)$. The complete graph is given

below.



d.
$$h(15) = -\frac{44(15)^2}{28^2} + (15) + 6 \approx 8.4 \text{ feet}$$

No; when the ball is 15 feet in front of the foul line, it will be below the hoop. Therefore it cannot go through the hoop.

In order for the ball to pass through the hoop, we need to have $h(15) = 10$.

$$10 = -\frac{44(15)^2}{v^2} + (15) + 6$$

$$-11 = -\frac{44(15)^2}{v^2}$$

$$v^2 = 4(225)$$

$$v^2 = 900$$

$$v = 30 \text{ ft/sec}$$

The ball must be shot with an initial velocity of 30 feet per second in order to go through the hoop.

30.
$$h(x) = -\frac{136x^2}{v^2} + 2.7x + 3.5$$

a. We want $h(15) = 10$.

$$-\frac{136(15)^2}{v^2} + 2.7(15) + 3.5 = 10$$

$$-\frac{30,600}{v^2} = -34$$

b.
$$h(x) = -\frac{126x^2}{30^2} + 2.7x + 3.5$$

which simplifies to

$$h(x) = -\frac{34}{225}x^2 + 2.7x + 3.5$$

c. Using the velocity from part (b),

$$h(9) = -\frac{34}{225}(9)^2 + 2.7(9) + 3.5 = 15.56 \text{ ft}$$

The ball will be 15.56 feet above the floor when it has traveled 9 feet in front of the foul line.

d. Select several values for x and use these to find the corresponding values for h . Use the results to form ordered pairs (x, h) . Plot the points and connect with a smooth curve.

$$h(0) = -\frac{34}{225}(0)^2 + 2.7(0) + 3.5 = 3.5 \text{ ft}$$

$$h(5) = -\frac{34}{225}(5)^2 + 2.7(5) + 3.5 \approx 13.2 \text{ ft}$$

$$(0, 3.5), (5, 13.2), (15, 10)$$


$$h(15) = -\frac{34}{225}(15)^2 + 2.7(15) + 3.5 \approx 10 \text{ ft}$$

$$(0, 3.5),$$

Thus, some points on the graph are

$(5, 13.2)$, and $(15, 10)$. The complete graph

is given below.



31.
$$h(x) = \frac{-32x^2}{2} + x$$

Chapter 2: Functions and Their Graphs

$$v^2$$

$$v^2 = 900$$

$$v = 30 \text{ ft/sec}$$

The ball needs to be thrown with an initial velocity of 30 feet per second.

Section 2.2: The Graph of a Function

130

$$\begin{aligned} \text{a. } h(100) &= \frac{-32(100)^2}{130^2} + 100 \\ &= \frac{-320,000}{16,900} + 100 \approx 81.07 \text{ feet} \end{aligned}$$

$$\begin{aligned} \text{b. } h(300) &= \frac{-32(300)^2}{130^2} + 300 \\ &= \frac{-2,880,000}{16,900} + 300 \approx 129.59 \text{ feet} \end{aligned}$$

c.
$$h(500) = \frac{-32(500)^2}{130^2} + 500$$

$$= \frac{-8,000,000}{16,900} + 500 \approx 26.63 \text{ feet}$$

$h(500) \approx 26.63$ feet represents the height of the golf ball, in feet, after it has traveled a horizontal distance of 500 feet.

d. Solving $h(x) = \frac{-32x^2}{130^2} + x = 0$

$$\frac{-32x^2}{130^2} + x = 0$$

$$x \left(\frac{-32x}{130^2} + 1 \right) = 0$$

$$x = 0 \text{ or } \frac{-32x}{130^2} + 1 = 0$$

$$1 = \frac{32x}{130^2}$$


$$130^2 = 32x$$

$$x = \frac{130^2}{32} = 528.13 \text{ feet}$$

Therefore, the golf ball travels 528.13 feet.

e. $y_1 = \frac{-32x^2}{130^2} + x$

150



f. Use INTERSECT on the graphs of

$$\frac{-32x^2}{130^2}$$

$$y_1 = \frac{-32x^2}{130^2} + x \text{ and } y_2 = 90.$$

150

The ball reaches a height of 90 feet twice. The first time is when the ball has traveled approximately 115.07 feet, and the second time is when the ball has traveled about 413.05 feet.

g. The ball travels approximately 275 feet before it reaches its maximum height of approximately 131.8 feet.

X	Y1
200	124.26
225	129.14
250	131.66
275	131.8
300	129.59
325	125
350	118.05

X=275

h. The ball travels approximately 264 feet before it reaches its maximum height of approximately 132.03 feet.

X	Y1	Y2	Y1
260	132	132	132
261	132.01	132.01	132.01
262	132.02	132.02	132.02
263	132.03	132.03	132.03
264	132.03	132.03	132.03
265	132.02	132.02	132.02
266	132.01	132.01	132.01

Y1=132.029112426 Y1=132.031242604

X	Y1
260	132
261	132.01
262	132.02
263	132.03
264	132.03
265	132.02
266	132.01

Y1=132.029585799

32. $A(x) = 4x \sqrt{1-x^2}$

a. Domain of $A(x) = 4x\sqrt{1-x^2}$; we know

that x must be greater than or equal to zero, since x represents a length. We also need

$1-x^2 \geq 0$, since this expression occurs under a square root. In fact, to avoid Area = 0, we require

$$x > 0 \text{ and } 1-x^2 > 0.$$


$$\text{Solve: } 1-x^2 > 0$$

$$\left(\frac{\quad}{1+x} \right) \left(\frac{\quad}{1-x} \right) > 0$$


$$\text{Case 1: } 1+x > 0 \text{ and } 1-x > 0$$

$$x > -1 \text{ and } x < 1$$

$$\text{(i.e. } -1 < x < 1)$$



150



Section 2.2: The Graph of a Function


Case2: $1 + x < 0$ and $1 - x < 0$

$$x < -1 \text{ and } x > 1$$

(which is impossible)

Therefore the domain of A is $\{x \mid 0 < x < 1\}$.

- b. Graphing $A(x) = 4x\sqrt{1-x^2}$




- c. When $x = 0.7$ feet, the cross-sectional area is maximized at approximately 1.9996 square feet. Therefore, the length of the base of the beam should be 1.4 feet in order to maximize the cross-sectional area.

X	Y1
.3	1.1447
.4	1.4664
.5	1.7321
.6	1.82
.7	1.9996
.8	1.82
.9	1.5692

X=.7

33. $C(x) = 100 + \frac{x}{10} + \frac{36000}{x}$

- a. Graphing:



- b. TblStart = 0; ΔTbl = 50

X	Y1
0	ERROR
50	825
100	470
150	355
200	300
250	269
300	250

Y1=100+X/10+36000...

- c. The cost per passenger is minimized to about \$220 when the ground speed is roughly 600 miles per hour.


X	Y1
450	225
500	225
550	220.45
600	220
650	220.38
700	221.43
750	223

X=600

34. $W(h) = m\left(\frac{4000}{4000+h}\right)^2$

On Pike's Peak, Amy will weigh about 119.84 pounds.

- b. Graphing:



- c. Create a TABLE:

X	Y1
0	120
.5	119.97
1	119.94
1.5	119.91
2	119.88
2.5	119.85
3	119.82

X=0

X	Y1
2	119.88
2.5	119.85
3	119.82
3.5	119.79
4	119.76
4.5	119.73
5	119.7

X=5

The weight W will vary from 120 pounds to about 119.7 pounds.

- d. By refining the table, Amy will weigh 119.95 lbs at a height of about 0.83 miles

X	Y1
.5	119.97
.6	119.96
.7	119.96
.8	119.95
.9	119.95
1	119.94
1.1	119.93

X=.8

X	Y1
.8	119.95
.81	119.95
.82	119.95
.83	119.95
.84	119.95
.85	119.95
.86	119.95

Y1=119.950215496

- e. Yes, 4382 feet is reasonable.

35. a. $(f + g)(2) = f(2) + g(2) = 2 + 1 = 3$

b. $(f + g)(4) = f(4) + g(4) = 1 + (-3) = -2$

c. $(f - g)(6) = f(6) - g(6) = 0 - 1 = -1$

d. $(g - f)(6) = g(6) - f(6) = 1 - 0 = 1$

e. $(f \cdot g)(2) = f(2) \cdot g(2) = 2(1) = 2$

f. $\left(\frac{f}{g}\right)(4) = \frac{f(4)}{g(4)} = \frac{1}{-3} = -\frac{1}{3}$

36. a. $C(0) = 5000$

This represents the fixed overhead costs. That is, the company will incur costs of \$5000 per day even if no computers are manufactured.

b. $C(10) = 19,000$

Chapter 2: Functions and Their Graphs

a. $h = 14110$ feet ≈ 2.67 miles ;

$$W(2.67) = 120 \left(\frac{4000}{4000 + 2.67} \right)^2 \approx 119.84$$


Section 2.2: The Graph of a Function


It costs the company \$19,000 to produce 10 computers in a day.

- c. $C(50) = 51,000$
It costs the company \$51,000 to produce 50 computers in a day.
 - d. The domain is $\{q \mid 0 \leq q \leq 100\}$. This indicates that production capacity is limited to 100 computers in a day.
 - e. The graph is curved down and rises slowly at first. As production increases, the graph becomes rises more quickly and changes to being curved up.
 - f. The inflection point is where the graph changes from being curved down to being curved up.
37. a. $C(0) = 80$
This represents the monthly fee. The plan costs \$80 per month even if no minutes are used.
- b. $C(1000) = 80$
The monthly charge is \$80 if 1000 minutes are used. Since this is the same as the cost for 0 minutes, all these minutes are included in the base plan.
- c. $C(2000) = 210$
The monthly charge is \$210 if 2000 minutes are used.
- d. The domain is $\{m \mid 0 \leq m \leq 14,400\}$. The domain implies that there are at most 14,400 anytime minutes in a month.
- e. The graph starts off flat (horizontal line), then increases at a constant rate (straight line with positive slope) after $m = 1000$.

38. Answers will vary. From a graph, the domain can be found by visually locating the x-values for which the graph is defined. The range can be found in a similar fashion by visually locating the y-values for which the function is defined. If an equation is given, the domain can be found by locating any restricted values and removing them from the set of real numbers. The range can be found by using known properties of the graph of the equation, or estimated by means of a table of values.
39. The graph of a function can have any number of x-intercepts. The graph of a function can have at most one y-intercept (otherwise the graph would fail the vertical line test).
40. Yes, the graph of a single point is the graph of a function since it would pass the vertical line test. The equation of such a function would be something like the following: $f(x) = 2$, where $x = 7$.

41. (a) III; (b) IV; (c) I; (d) V; (e) II
42. (a) II; (b) V; (c) IV; (d) III; (e) I





45. a. 2 hours elapsed; Kevin was between 0 and 3 miles from home.
 b. 0.5 hours elapsed; Kevin was 3 miles from home.
 c. 0.3 hours elapsed; Kevin was between 0 and 3 miles from home.
 d. 0.2 hours elapsed; Kevin was at home.
 e. 0.9 hours elapsed; Kevin was between 0 and 2.8 miles from home.
 f. 0.3 hours elapsed; Kevin was 2.8 miles from home.
 g. 1.1 hours elapsed; Kevin was between 0 and 2.8 miles from home.
 h. The farthest distance Kevin is from home is 3 miles.
 i. Kevin returned home 2 times.
46. a. Michael travels fastest between 7 and 7.4 minutes. That is, $(7, 7.4)$.
 b. Michael's speed is zero between 4.2 and 6 minutes. That is, $(4.2, 6)$.
 c. Between 0 and 2 minutes, Michael's speed increased from 0 to 30 miles/hour.
 d. Between 4.2 and 6 minutes, Michael was stopped (i.e., his speed was 0 miles/hour).
 e. Between 7 and 7.4 minutes, Michael was traveling at a steady rate of 50 miles/hour.
 f. Michael's speed is constant between 2 and 4 minutes, between 4.2 and 6 minutes,

47. Answers (graphs) will vary. Points of the form $(5, y)$ and of the form $(x, 0)$ cannot be on the graph of the function.
48. The only such function is $f(x) = 0$ because it is the only function for which $f(x) = -f(x)$. Any other such graph would fail the vertical line test.

Section 2.3

- $2 < x < 5$
- slope = $\frac{\Delta y}{\Delta x} = \frac{8-3}{3-(-2)} = \frac{5}{5} = 1$
- x-axis: $y \rightarrow -y$
 $(-y) = 5x^2 - 1$
 $-y = 5x^2 - 1$
 $y = -5x^2 + 1$ different
 y-axis: $x \rightarrow -x$
 $y = 5(-x)^2 - 1$
 $y = 5x^2 - 1$ same
 origin: $x \rightarrow -x$ and $y \rightarrow -y$
 $(-y) = 5(-x)^2 - 1$
 $-y = 5x^2 - 1$
 $y = -5x^2 + 1$ different
 The equation has symmetry with respect to the y-axis only.
- $y - y_1 = m(x - x_1)$
 $y - (-2) = 5(x - 3)$
 $y + 2 = 5(x - 3)$
- $y = x^2 - 9$
 x-intercepts:
 $0 = x^2 - 9$
 $x^2 = 9 \rightarrow x = \pm 3$
 y-intercept:
 2

Chapter 2: Functions and Their Graphs

between 7 and 7.4 minutes, and between 7.6

and 8 minutes. That is, on the intervals
(2, 4), (4.2, 6), (7, 7.4), and (7.6, 8).

Section 2.3: Properties of Functions

$$y = (0) - 9 = -9$$

The intercepts are $(-3, 0)$, $(3, 0)$, and $(0, -9)$.

6. increasing

7. even; odd
8. True
9. True
10. False; odd functions are symmetric with respect to the origin. Even functions are symmetric with respect to the y -axis.
11. Yes
12. No, it is increasing.
13. No, it only increases on $(5, 10)$.
14. Yes
15. f is increasing on the intervals $(-8, -2)$, $(0, 2)$, $(5, \infty)$.
16. f is decreasing on the intervals: $(-\infty, -8)$, $(-2, 0)$, $(2, 5)$.
17. Yes. The local maximum at $x = 2$ is 10.
18. No. There is a local minimum at $x = 5$; the local minimum is 0.
19. f has local maxima at $x = -2$ and $x = 2$. The local maxima are 6 and 10, respectively.
20. f has local minima at $x = -8$, $x = 0$ and $x = 5$. The local minima are -4 , 0, and 0, respectively.
21. a. Intercepts: $(-2, 0)$, $(2, 0)$, and $(0, 3)$.
- b. Domain: $\{x \mid -4 \leq x \leq 4\}$ or $[-4, 4]$;
Range: $\{y \mid 0 \leq y \leq 3\}$ or $[0, 3]$.
- c. Increasing: $(-2, 0)$ and $(2, 4)$;
Decreasing: $(-4, -2)$ and $(0, 2)$.
23. a. Intercepts: $(0, 1)$.
- b. Domain: $\{x \mid x \text{ is any real number}\}$;
Range: $\{y \mid y > 0\}$ or $(0, \infty)$.
- c. Increasing: $(-\infty, \infty)$; Decreasing: never.
- d. Since the graph is not symmetric with respect to the y -axis or the origin, the function is neither even nor odd.
24. a. Intercepts: $(1, 0)$.
- b. Domain: $\{x \mid x > 0\}$ or $(0, \infty)$;
Range: $\{y \mid y \text{ is any real number}\}$.
- c. Increasing: $(0, \infty)$; Decreasing: never.
- d. Since the graph is not symmetric with respect to the y -axis or the origin, the function is neither even nor odd.
25. a. Intercepts: $(-\pi, 0)$, $(\pi, 0)$, and $(0, 0)$.
- b. Domain: $\{x \mid -\pi \leq x \leq \pi\}$ or $[-\pi, \pi]$;
Range: $\{y \mid -1 \leq y \leq 1\}$ or $[-1, 1]$.
- c. Increasing: $(-\frac{\pi}{2}, \frac{\pi}{2})$;
Decreasing: $(-\pi, -\frac{\pi}{2})$ and $(\frac{\pi}{2}, \pi)$.
- d. Since the graph is symmetric with respect to the origin, the function is odd.
26. a. Intercepts: $(-\frac{\pi}{2}, 0)$, $(\frac{\pi}{2}, 0)$, and $(0, 1)$.
- b. Domain: $\{x \mid -\pi \leq x \leq \pi\}$ or $[-\pi, \pi]$;
Range: $\{y \mid -1 \leq y \leq 1\}$ or $[-1, 1]$.

Chapter 2: Functions and Their Graphs

- d. Since the graph is symmetric with respect to the y -axis, the function is even.
22. a. Intercepts: $(-1, 0)$, $(1, 0)$, and $(0, 2)$.
- b. Domain: $\{x \mid -3 \leq x \leq 3\}$ or $[-3, 3]$;
Range: $\{y \mid 0 \leq y \leq 3\}$ or $[0, 3]$.
- c. Increasing: $(-1, 0)$ and $(1, 3)$;
Decreasing: $(-3, -1)$ and $(0, 1)$.
- d. Since the graph is symmetric with respect to the y -axis, the function is even.

Section 2.3: Properties of Functions

- c. Increasing: $(-\pi, 0)$; Decreasing: $(0, \pi)$.
- d. Since the graph is symmetric with respect to the y -axis, the function is even.
- (1) (5) (1)
27. a. Intercepts: $(3, 0)$, $(2, 0)$, and $(0, 2)$.
- b. Domain: $\{x \mid -3 \leq x \leq 3\}$ or $[-3, 3]$;
Range: $\{y \mid -1 \leq y \leq 2\}$ or $[-1, 2]$.

- c. Increasing: $(2, 3)$; Decreasing: $(-1, 1)$;
Constant: $(-3, -1)$ and $(1, 2)$
- d. Since the graph is not symmetric with respect to the y -axis or the origin, the function is neither even nor odd.
28. a. Intercepts: $(-2.3, 0)$, $(3, 0)$, and $(0, 1)$.
- b. Domain: $\{x \mid -3 \leq x \leq 3\}$ or $[-3, 3]$;
Range: $\{y \mid -2 \leq y \leq 2\}$ or $[-2, 2]$.
- c. Increasing: $(-3, -2)$ and $(0, 2)$;
Decreasing: $(2, 3)$; Constant: $(-2, 0)$.
- d. Since the graph is not symmetric with respect to the y -axis or the origin, the function is neither even nor odd.
29. a. f has a local maximum of 3 at $x = 0$.
- b. f has a local minimum of 0 at both $x = -2$ and $x = 2$.
30. a. f has a local maximum of 2 at $x = 0$.
- b. f has a local minimum of 0 at both $x = -1$ and $x = 1$.
31. a. f has a local maximum of 1 at $x = \frac{\pi}{2}$.
- b. f has a local minimum of -1 at $x = -\frac{\pi}{2}$.
32. a. f has a local maximum of 1 at $x = 0$.
- b. f has a local minimum of -1 both at $x = -\pi$ and $x = \pi$.
33. $f(x) = 4x^3$
 $f(-x) = 4(-x)^3 = -4x^3 = -f(x)$

36. $h(x) = 3x^3 + 5$
 $h(-x) = 3(-x)^3 + 5 = -3x^3 + 5$
 h is neither even nor odd.
37. $F(x) = \sqrt[3]{x}$
 $F(-x) = \sqrt[3]{-x} = -\sqrt[3]{x} = -F(x)$
Therefore, F is odd.
38. $G(x) = \sqrt{x}$
 $G(-x) = \sqrt{-x}$
 G is neither even nor odd.
39. $f(x) = x + |x|$
 $f(-x) = -x + |-x|$
 f is neither even nor odd.
40. $f(x) = \sqrt[3]{2x^2 + 1}$
 $f(-x) = \sqrt[3]{2(-x)^2 + 1} = \sqrt[3]{2x^2 + 1} = f(x)$
Therefore, f is even.
41. $g(x) = \frac{x^2 + 3}{x^2 - 1}$
 $g(-x) = \frac{(-x)^2 + 3}{(-x)^2 - 1} = \frac{x^2 + 3}{x^2 - 1} = g(x)$
Therefore, g is even.
42. $h(x) = \frac{x}{x^2 - 1}$
 $h(-x) = \frac{-x}{(-x)^2 - 1} = \frac{-x}{x^2 - 1} = -h(x)$
Therefore, h is odd.
43. $h(x) = \frac{-x^3}{3x^2 - 9}$

Chapter 2: Functions and Their Graphs

Therefore, x^3 is odd.

34. $f(x) = 2x^4 - x^2$
 $f(-x) = 2(-x)^4 - (-x)^2 = 2x^4 - x^2 = f(x)$

Therefore, f is even.

35. $g(x) = -3x^2 - 5$
 $g(-x) = -3(-x)^2 - 5 = -3x^2 - 5 = g(x)$
 Therefore, g is even.

Section 2.3: Properties of Functions

$$h(-x) = \frac{2(-x)}{3(-x)^2 - 9} = \frac{-2x}{3x^2 - 9} = -h(x)$$

Therefore, h is odd.

44. $F(x) = \frac{2x}{|x|}$



$$F(-x) = \frac{2(-x)}{|-x|} = \frac{-2x}{|x|} = -F(x)$$

Therefore, F is odd.

- 45. f has an absolute maximum of 4 at $x = 1$.
 f has an absolute minimum of 1 at $x = 5$.
- 46. f has an absolute maximum of 4 at $x = 4$.
 f has an absolute minimum of 0 at $x = 5$.
- 47. f has an absolute minimum of 1 at $x = 1$.
 f has an absolute maximum of 4 at $x = 3$.
- 48. f has an absolute minimum of 1 at $x = 0$.
 f has no absolute maximum.
- 49. f has an absolute minimum of 0 at $x = 0$.
 f has no absolute maximum.
- 50. f has an absolute maximum of 4 at $x = 2$.
 f has no absolute minimum.
- 51. f has no absolute maximum or minimum.
- 52. f has no absolute maximum or minimum.
- 53. $f(x) = x^3 - 3x + 2$ on the interval $(-2, 2)$



Use MAXIMUM and MINIMUM on the graph

of $y_1 = x^3 - 3x + 2$.





local maximum at: $(-1, 4)$;
 local minimum at: $(1, 0)$
 f is increasing on: $(-2, -1)$ and $(1, 2)$;
 f is decreasing on: $(-1, 1)$

- 54. $f(x) = x^3 - 3x^2 + 5$ on the interval $(-1, 3)$
Use MAXIMUM and MINIMUM on the graph of



local maximum at: $(0, 5)$;
 local minimum at: $(2, 1)$
 f is increasing on: $(-1, 0)$ and $(2, 3)$;
 f is decreasing on: $(0, 2)$


- 55. $f(x) = x^5 - x^3$ on the interval $(-2, 2)$
Use MAXIMUM and MINIMUM on the graph of $y_1 = x^5 - x^3$.



local maximum at: $(-0.77, 0.19)$;
 local minimum at: $(0.77, -0.19)$;
 f is increasing on: $(-2, -0.77)$ and $(0.77, 2)$;
 f is decreasing on: $(-0.77, 0.77)$

- 56. $f(x) = x^4 - x^2$ on the interval $(-2, 2)$
Use MAXIMUM and MINIMUM on the graph of $y_1 = x^4 - x^2$.

$$y_1 = x^2 - 3x + 5.$$




local maximum at: $(0, 0)$;
 local minimum at: $(-0.71, -0.25)$, $(0.71, -0.25)$
 f is increasing on: $(-0.71, 0)$ and $(0.71, 2)$;
 f is decreasing on: $(-2, -0.71)$ and $(0, 0.71)$

57. $f(x) = -0.2x^3 - 0.6x^2 + 4x - 6$ on the interval $(-6, 4)$

Use MAXIMUM and MINIMUM on the graph of $y_1 = -0.2x^3 - 0.6x^2 + 4x - 6$.

58. $f(x) = -0.4x^3 + 0.6x^2 + 3x - 2$ on the interval $(-4, 5)$


Use MAXIMUM and MINIMUM on the graph of $y_1 = -0.4x^3 + 0.6x^2 + 3x - 2$.



local maximum at: $(2.16, 3.25)$;
 local minimum at: $(-1.16, -4.05)$
 f is increasing on: $(-1.16, 2.16)$;
 f is decreasing on: $(-4, -1.16)$ and $(2.16, 5)$

59. $f(x) = 0.25x^4 + 0.3x^3 - 0.9x^2 + 3$ on the interval $(-3, 2)$


Use MAXIMUM and MINIMUM on the graph of $y_1 = 0.25x^4 + 0.3x^3 - 0.9x^2 + 3$.




Chapter 2

Graphs

- local maximum at: $(1.77, -1.91)$;
- local minimum at: $(-3.77, -18.89)$
- f is increasing on: $(-3.77, 1.77)$;
- f is decreasing on: $(-6, -3.77)$ and $(1.77, 4)$




Properties of Functions



local maximum at: $(0, 3)$;

local minimum at: $(-1.87, 0.95)$, $(0.97, 2.65)$
 f is increasing on: $(-1.87, 0)$ and $(0.97, 2)$;
 f is decreasing on: $(-3, -1.87)$ and $(0, 0.97)$

60. $f(x) = -0.4x^4 - 0.5x^3 + 0.8x^2 - 2$ on the interval $(-3, 2)$
 Use MAXIMUM and MINIMUM on the graph of $y_1 = -0.4x^4 - 0.5x^3 + 0.8x^2 - 2$.



local maxima at: $(-1.57, -0.52)$, $(0.64, -1.87)$;
 local minimum at: $(0, -2)$

f is increasing on: $(-3, -1.57)$ and $(0, 0.64)$;
 f is decreasing on: $(-1.57, 0)$ and $(0.64, 2)$

61. $f(x) = -2x^2 + 4$
 a. Average rate of change of f from $x = 0$ to $x = 2$

$$\frac{f(2) - f(0)}{2 - 0} = \frac{-2(2)^2 + 4 - (-2(0)^2 + 4)}{2 - 0}$$

- c. Average rate of change of f from $x = 1$ to $x = 4$:

$$\begin{aligned} \frac{f(4) - f(1)}{4 - 1} &= \frac{-2(4)^2 + 4 - (-2(1)^2 + 4)}{3} \\ &= \frac{-28 - 2}{3} = \frac{-30}{3} = -10 \end{aligned}$$

62. $f(x) = -x^3 + 1$

- a. Average rate of change of f from $x = 0$ to $x = 2$:

$$\begin{aligned} \frac{f(2) - f(0)}{2 - 0} &= \frac{-2^3 + 1 - (-0^3 + 1)}{2} \\ &= \frac{-7 - 1}{2} = \frac{-8}{2} = -4 \end{aligned}$$

- b. Average rate of change of f from $x = 1$ to $x = 3$:

$$\begin{aligned} \frac{f(3) - f(1)}{3 - 1} &= \frac{-3^3 + 1 - (-1^3 + 1)}{2} \\ &= \frac{-26 - 0}{2} = \frac{-26}{2} = -13 \end{aligned}$$

- c. Average rate of change of f from $x = -1$ to $x = 1$:

$$\begin{aligned} \frac{f(1) - f(-1)}{1 - (-1)} &= \frac{-1^3 + 1 - (-(-1)^3 + 1)}{2} \\ &= \frac{0 - 2}{2} = \frac{-2}{2} = -1 \end{aligned}$$

63. $g(x) = x^3 - 2x + 1$

- a. Average rate of change of g from $x = -3$ to $x = -2$:

$$\begin{aligned} \frac{g(-2) - g(-3)}{-2 - (-3)} &= \frac{(-2)^3 - 2(-2) + 1 - [(-3)^3 - 2(-3) + 1]}{-2 - (-3)} \\ &= \frac{[-2^3 - 2(-2) + 1] - [-3^3 - 2(-3) + 1]}{1} \\ &= \frac{(-8) - (-20)}{1} = 12 \end{aligned}$$

Chapter 2: Functions and Their Graphs

$$2 - 0 = \frac{(-4) - (-4)}{2} = \frac{-8}{2} = -4$$

- b. Average rate of change of f from $x = 1$ to $x = 3$:

$$\begin{aligned} \frac{f(3) - f(1)}{3 - 1} &= \frac{(-2(3)^2 + 4) - (-2(1)^2 + 4)}{2} \\ &= \frac{(-14) - (2)}{2} = \frac{-16}{2} = -8 \end{aligned}$$

Section 2.3: Properties of Functions

$$= \frac{17 - 1}{1} = 16$$

- b. Average rate of change of g from $x = -1$ to $x = 1$:

$$\frac{g(1) - g(-1)}{1 - (-1)}$$

$$= \frac{[(1)^3 - 2(1) + 1] - [(-1)^3 - 2(-1) + 1]}{2}$$

$$= \frac{(0) - (2)}{2} = \frac{-2}{2} = -1$$

- c. Average rate of change of g from $x = 1$ to $x = 3$:

$$\frac{g(3) - g(1)}{3 - 1}$$

$$= \frac{[(3)^3 - 2(3) + 1] - [(1)^3 - 2(1) + 1]}{2}$$

$$= \frac{(22) - (0)}{2} = \frac{22}{2} = 11$$

64. $h(x) = x^2 - 2x + 3$

- a. Average rate of change of h from $x = -1$ to $x = 1$:

$$\frac{h(1) - h(-1)}{1 - (-1)}$$

$$= \frac{[(1)^2 - 2(1) + 3] - [(-1)^2 - 2(-1) + 3]}{2}$$

$$= \frac{(2) - (6)}{2} = \frac{-4}{2} = -2$$

- b. Average rate of change of h from $x = 0$ to $x = 2$:

$$\frac{h(2) - h(0)}{2 - 0}$$

$$= \frac{[(2)^2 - 2(2) + 3] - [(0)^2 - 2(0) + 3]}{2}$$

$$= \frac{(3) - (3)}{2} = \frac{0}{2} = 0$$

- c. Average rate of change of h from $x = 2$ to $x = 5$:

$$\frac{h(5) - h(2)}{5 - 2}$$

$$= \frac{[(5)^2 - 2(5) + 3] - [(2)^2 - 2(2) + 3]}{3}$$

$$= \frac{(18) - (3)}{3} = \frac{15}{3} = 5$$

65. $f(x) = 5x - 2$

- a. Average rate of change of f from 1 to 3:

$$\frac{\Delta y}{\Delta x} = \frac{f(3) - f(1)}{3 - 1} = \frac{13 - 3}{2} = 5$$

Thus, the average rate of change of f from 1 to 3 is 5.

- b. From (a), the slope of the secant line joining

$(1, f(1))$ and $(3, f(3))$ is 5. We use the

point-slope form to find the equation of the secant line:

$$y - y_1 = m_{\text{sec}}(x - x_1)$$

$$y - 3 = 5(x - 1)$$

$$y - 3 = 5x - 5$$

$$y = 5x - 2$$

66. $f(x) = -4x + 1$

- a. Average rate of change of f from 2 to 5:

$$\frac{\Delta y}{\Delta x} = \frac{f(5) - f(2)}{5 - 2} = \frac{-19 - (-7)}{3}$$

$$= \frac{-12}{3} = -4$$

67.

Chapter 2: Functions and Their Graphs

Therefore, the average rate of change of f from 2 to 5 is -4 .

- b. From (a), the slope of the secant line joining $(2, f(2))$ and $(5, f(5))$ is -4 . We use the point-slope form to find the equation of the secant line:

$$y - y_1 = m_{\text{sec}}(x - x_1)$$

Section 2.3: Properties of Functions

$$\begin{aligned} y &= -4(x - 2) \\ &= y + 7 = -4x + 8 \\ &= y = -4x + 1 \end{aligned}$$

$$g(x) = x^2 - 2$$

- a. Average rate of change of g from -2 to 1 :

$$\frac{\Delta y}{\Delta x} = \frac{g(1) - g(-2)}{1 - (-2)} = \frac{-1 - 2}{1 - (-2)} = \frac{-3}{3} = -1$$

Therefore, the average rate of change of g from -2 to 1 is -1 .

- b. From (a), the slope of the secant line joining $(-2, g(-2))$ and $(1, g(1))$ is -1 . We use the point-slope form to find the equation of the secant line:

$$y - y_1 = m_{\text{sec}}(x - x_1)$$

$$y - 2 = -1(x - (-2))$$

$$y - 2 = -x - 2$$

$$y = -x$$

68. $g(x) = x^2 + 1$

- a. Average rate of change of g from -1 to 2 :

$$\frac{\Delta y}{\Delta x} = \frac{g(2) - g(-1)}{2 - (-1)} = \frac{5 - 2}{2 - (-1)} = \frac{3}{3} = 1$$

Therefore, the average rate of change of g from -1 to 2 is 1 .

- b. From (a), the slope of the secant line joining

$$(-1, g(-1)) \text{ and } (2, g(2)) \text{ is } 1. \text{ We use the}$$

point-slope form to find the equation of the secant line:

$$y - y_1 = m_{\text{sec}}(x - x_1)$$

$$y - 2 = 1(x - (-1))$$

$$y - 2 = x + 1$$

$$y = x + 3$$

69. $h(x) = x^2 - 2x$

- a. Average rate of change of h from 2 to 4 :

$$\frac{\Delta y}{\Delta x} = \frac{h(4) - h(2)}{4 - 2} = \frac{8 - 0}{4 - 2} = \frac{8}{2} = 4$$

Therefore, the average rate of change of h from 2 to 4 is 4 .

- b. From (a), the slope of the secant line joining $(2, h(2))$ and $(4, h(4))$ is 4 . We use the point-slope form to find the equation of the secant line:

Therefore, the average rate of change of h from 0 to 3 is -5 .

- b. From (a), the slope of the secant line joining $(0, h(0))$ and $(3, h(3))$ is -5 . We use the point-slope form to find the equation of the secant line:

$$y - y_1 = m_{\text{sec}}(x - x_1)$$

$$y - 0 = -5(x - 0)$$

$$y = -5x$$

71. a. $g(x) = x^3 - 27x$

$$g(-x) = (-x)^3 - 27(-x)$$

$$= -x^3 + 27x$$

$$= -(x^3 - 27x)$$

$$= -g(x)$$

Since $g(-x) = -g(x)$, the function is odd.

- b. Since $g(x)$ is odd then it is symmetric

about the origin so there exist a local maximum at $x = -3$.

$$g(-3) = (-3)^3 - 27(-3) = -27 + 81 = 54$$

So there is a local maximum of 54 at $x = -3$.

72. $f(x) = -x^3 + 12x$

a. $f(-x) = -(-x)^3 + 12(-x)$

$$= x^3 - 12x$$

$$= -(-x^3 + 12x)$$

$$= -f(x)$$

Since $f(-x) = -f(x)$, the function is odd.

- b. Since $f(x)$ is odd then it is symmetric

Chapter 2: Functions and Their Graphs

$$y - y_1 = m_{\text{sec}}(x - x_1)$$

$$y - 0 = 4(x - 2)$$

$$y = 4x - 8$$

70. $h(x) = -2x^2 + x$

a. Average rate of change from 0 to 3:

$$\frac{\Delta y}{\Delta x} = \frac{h(3) - h(0)}{3 - 0} = \frac{-15 - 0}{3 - 0}$$

$$= \frac{-15}{3} = -5$$

Section 2.3: Properties of Functions

about the origin so there exist a local maximum at $x = -3$.

$$f(-2) = -(-2)^3 + 12(-2) = 8 - 24 = -16$$

So there is a local maximum of -16 at $x = -2$.

73. $F(x) = -x^4 + 8x^2 + 8$

a. $F(-x) = -(-x)^4 + 8(-x)^2 + 8$
 $= -x^4 + 8x^2 + 8$
 $= F(x)$

Since $F(-x) = F(x)$, the function is even.

b. Since the function is even, its graph has y-axis symmetry. The second local maximum is in quadrant II and is 24 and occurs at $x = -2$.

c. Because the graph has y-axis symmetry, the area under the graph between $x = 0$ and

$x = 3$ bounded below by the x-axis is the same as the area under the graph between $x = -3$ and $x = 0$ bounded below the x-axis. Thus, the area is 47.4 square units.

74. $G(x) = -x^4 + 32x^2 + 144$

a. $G(-x) = -(-x)^4 + 32(-x)^2 + 144$
 $= -x^4 + 32x^2 + 144$
 $= G(x)$

Since $G(-x) = G(x)$, the function is even.


b. Since the function is even, its graph has y-axis symmetry. The second local maximum is in quadrant II and is 400 and occurs at $x = -4$.

c. Because the graph has y-axis symmetry, the area under the graph between $x = 0$ and $x = 6$ bounded below by the x-axis is the same as the area under the graph between


$x = -6$ and $x = 0$ bounded below the x-axis. Thus, the area is 1612.8 square units.

75. $\bar{C}(x) = 0.3x^2 + 21x - 251 + \frac{2500}{x}$

a. $y_1 = 0.3x^2 + 21x - 251 + \frac{2500}{x}$



b. Use MINIMUM. Rounding to the nearest whole number, the average cost is minimized when approximately 10 lawnmowers are produced per hour.




c. The minimum average cost is approximately \$239 per mower.

76. a. $C(t) = -.002t^4 + .039t^3 - .285t^2 + .766t + .085$

Graph the function on a graphing utility and use the Maximum option from the CALC menu.


1



The concentration will be highest after about 2.16 hours.


b. Enter the function in Y1 and 0.5 in Y2. Graph the two equations in the same window and use the Intersect option from the CALC menu.

1



0

1



0

0


Chapter 2: Functions and Their Graphs

0 30
-300

Section 2.3: Properties of Functions

After taking the medication, the woman can feed her child within the first 0.71 hours (about 42 minutes) or after 4.47 hours (about 4 hours 28 minutes) have elapsed.

77. a.



b. The slope represents the average rate of change of the debt from 2000 to 2002.


$$\begin{aligned} \text{c. avg. rate of change} &= \frac{P(2002) - P(2000)}{2002 - 2000} \\ &= \frac{6228 - 5674}{2} \\ &= \frac{554}{2} \\ &= \$ 277 \text{ billion} \end{aligned}$$

$$\begin{aligned} \text{d. avg. rate of change} &= \frac{P(2006) - P(2004)}{2006 - 2004} \\ &= \frac{8507 - 7379}{2} \\ &= \frac{1128}{2} \\ &= \$ 564 \text{ billion} \end{aligned}$$

$$\begin{aligned} \text{e. avg. rate of change} &= \frac{P(2010) - P(2008)}{2010 - 2008} \\ &= \frac{13562 - 10025}{2} \\ &= \frac{3537}{2} \\ &= \$ 1768.5 \text{ billion} \end{aligned}$$

c. The average rate of change is increasing as time passes.

78. a.



b. The slope represents the average rate of change of the population from 0 to 2.5 hours.

$$\begin{aligned} \text{c. avg. rate of change} &= \frac{P(2.5) - P(0)}{2.5 - 0} \\ &= \frac{0.18 - 0.09}{2.5 - 0} \\ &= \frac{0.09}{2.5} \\ &= 0.036 \text{ gram per hour} \end{aligned}$$

$$\begin{aligned} \text{c. avg. rate of change} &= \frac{P(6) - P(4.5)}{6 - 4.5} \\ &= \frac{0.50 - 0.35}{6 - 4.5} \\ &= \frac{0.15}{1.5} \\ &= 0.1 \text{ gram per hour} \end{aligned}$$

d. The average rate of change is increasing as time passes.

79. $f(x) = x^2$

a. Average rate of change of f from $x = 0$ to $x = 1$:

$$\frac{f(1) - f(0)}{1 - 0} = \frac{1^2 - 0^2}{1} = \frac{1}{1} = 1$$

b. Average rate of change of f from $x = 0$ to $x = 0.5$:

$$\frac{f(0.5) - f(0)}{0.5 - 0} = \frac{(0.5)^2 - 0^2}{0.5} = \frac{0.25}{0.5} = 0.5$$

c. Average rate of change of f from $x = 0$ to $x = 0.1$:

$$\frac{f(0.1) - f(0)}{0.1 - 0} = \frac{(0.1)^2 - 0^2}{0.1} = \frac{0.01}{0.1} = 0.1$$

$$0.1 - 0 = 0.1 = 0.1 = 0.1$$






- d. Average rate of change of f from $x = 0$ to $x = 0.01$:

$$\frac{f(0.01) - f(0)}{0.01 - 0} = \frac{(0.01)^2 - 0^2}{0.01} = \frac{0.0001}{0.01} = 0.01$$

- e. Average rate of change of f from $x = 0$ to $x = 0.001$:

$$\frac{f(0.001) - f(0)}{0.001 - 0} = \frac{(0.001)^2 - 0^2}{0.001} = \frac{0.000001}{0.001} = 0.001$$

- f. Graphing the secant lines:



- g. The secant lines are beginning to look more and more like the tangent line to the graph of f at the point where $x = 0$.
- h. The slopes of the secant lines are getting smaller and smaller. They seem to be approaching the number zero.

80. $f(x) = x^2$

- a. Average rate of change of f from $x = 1$ to $x = 2$:

$$\frac{f(2) - f(1)}{2 - 1} = \frac{2^2 - 1^2}{1} = \frac{3}{1} = 3$$

- b. Average rate of change of f from $x = 1$ to $x = 1.5$:

$$\frac{f(1.5) - f(1)}{1.5 - 1} = \frac{(1.5)^2 - 1^2}{0.5} = \frac{1.25}{0.5} = 2.5$$

- c. Average rate of change of f from $x = 1$ to $x = 1.1$:

$$\frac{f(1.1) - f(1)}{1.1 - 1} = \frac{(1.1)^2 - 1^2}{0.1} = \frac{0.21}{0.1} = 2.1$$






- d. Average rate of change of f from $x = 1$ to $x = 1.01$:

$$\frac{f(1.01) - f(1)}{1.01 - 1} = \frac{(1.01)^2 - 1^2}{0.01} = \frac{0.0201}{0.01} = 2.01$$

- e. Average rate of change of f from $x = 1$ to $x = 1.001$:

$$\begin{aligned}\frac{f(1.001) - f(1)}{1.001 - 1} &= \frac{(1.001)^2 - 1^2}{0.001} \\ &= \frac{0.002001}{0.001} = 2.001\end{aligned}$$

f. Graphing the secant lines:



g. The secant lines are beginning to look more and more like the tangent line to the graph of f at the point where $x = 1$.

h. The slopes of the secant lines are getting smaller and smaller. They seem to be approaching the number 2.

81. $f(x) = 2x + 5$

a. m_{sec}

b. When $x = 1$:

$h = 0.5 \Rightarrow m_{\text{sec}} = 2$

$h = 0.1 \Rightarrow m_{\text{sec}} = 2$

$h = 0.01 \Rightarrow m_{\text{sec}} = 2$

as $h \rightarrow 0, m_{\text{sec}} \rightarrow 2$

c. Using the point $(1, f(1)) = (1, 7)$ and slope,


$m = 2$, we get the secant line:

$y - 7 = 2(x - 1)$

$y - 7 = 2x - 2$

$y = 2x + 5$

d. Graphing:



The graph and the secant line coincide.

82. $f(x) = -3x + 2$

a.
$$m_{\text{sec}} = \frac{f(x+h) - f(x)}{h}$$

$$= \frac{-3(x+h) + 2 - (-3x + 2)}{h} = \frac{-3h}{h} = -3$$

b. When $x = 1$,

$h = 0.5 \Rightarrow m_{\text{sec}} = -3$

$h = 0.1 \Rightarrow m_{\text{sec}} = -3$

$h = 0.01 \Rightarrow m_{\text{sec}} = -3$

as $h \rightarrow 0, m_{\text{sec}} \rightarrow -3$

c. Using point $(1, f(1)) = (1, -1)$ and slope = -3 , we get the secant line:

$y - (-1) = -3(x - 1)$

$y + 1 = -3x + 3$

$y = -3x + 2$

d. Graphing:


Chapter 2: Functions and Their Graphs

$$= \frac{f(x+h) - f(x)}{h}$$

$$= \frac{2(x+h) + 5 - 2}{h}$$

$$\frac{x-5}{h} = \frac{2h}{h} = 2$$

The graph and the secant line coincide.



Properties of Functions

83. $f(x) = x^2 + 2x$

a.
$$m_{\text{sec}} = \frac{f(x+h) - f(x)}{h}$$

$$= \frac{(x+h)^2 + 2(x+h) - (x^2 + 2x)}{h}$$

$$= \frac{x^2 + 2xh + h^2 + 2x + 2h - x^2 - 2x}{h}$$


$$= \frac{2xh + h^2 + 2h}{h}$$

$$= 2x + h + 2$$

b. When $x = 1$,
 $h = 0.5 \Rightarrow m_{\text{sec}} = 2 \cdot 1 + 0.5 + 2 = 4.5$
 $h = 0.1 \Rightarrow m_{\text{sec}} = 2 \cdot 1 + 0.1 + 2 = 4.1$
 $h = 0.01 \Rightarrow m_{\text{sec}} = 2 \cdot 1 + 0.01 + 2 = 4.01$
 as $h \rightarrow 0$, $m_{\text{sec}} \rightarrow 2 \cdot 1 + 0 + 2 = 4$

c. Using point $(1, f(1)) = (1, 3)$ and slope = 4.01, we get the secant line:
 $y - 3 = 4.01(x - 1)$
 $y - 3 = 4.01x - 4.01$
 $y = 4.01x - 1.01$

d. Graphing:



84. $f(x) = 2x^2 + x$

a.
$$m_{\text{sec}} = \frac{f(x+h) - f(x)}{h}$$

$$= \frac{2(x+h)^2 + (x+h) - (2x^2 + x)}{h}$$

$$= \frac{2(x^2 + 2xh + h^2) + x + h - 2x^2 - x}{h}$$

$$= 4x + 2h + 1$$


$h = 0.1 \Rightarrow m_{\text{sec}} = 4 \cdot 1 + 2(0.1) + 1 = 5.2$

$h = 0.01 \Rightarrow m_{\text{sec}} = 4 \cdot 1 + 2(0.01) + 1 = 5.02$

as $h \rightarrow 0$, $m_{\text{sec}} \rightarrow 4 \cdot 1 + 2 \cdot 0 + 1 = 5$

c. Using point $(1, f(1)) = (1, 3)$ and slope = 5.02, we get the secant line:
 $y - 3 = 5.02(x - 1)$
 $y - 3 = 5.02x - 5.02$
 $y = 5.02x - 2.02$

d. Graphing:



85. $f(x) = 2x^2 - 3x + 1$

a.
$$m_{\text{sec}} = \frac{f(x+h) - f(x)}{h}$$

$$= \frac{2(x+h)^2 - 3(x+h) + 1 - (2x^2 - 3x + 1)}{h}$$

$$= \frac{2(x^2 + 2xh + h^2) - 3x - 3h + 1 - 2x^2 + 3x - 1}{h}$$

$$= \frac{2x^2 + 4xh + 2h^2 - 3x - 3h + 1 - 2x^2 + 3x - 1}{h}$$

$$= \frac{4xh + 2h^2 - 3h}{h}$$

$$= 4x + 2h - 3$$

b. When $x = 1$,
 $h = 0.5 \Rightarrow m_{\text{sec}} = 4 \cdot 1 + 2(0.5) - 3 = 2$
 $h = 0.1 \Rightarrow m_{\text{sec}} = 4 \cdot 1 + 2(0.1) - 3 = 1.2$
 $h = 0.01 \Rightarrow m_{\text{sec}} = 4 \cdot 1 + 2(0.01) - 3 = 1.02$
 as $h \rightarrow 0$, $m_{\text{sec}} \rightarrow 4 \cdot 1 + 2(0) - 3 = 1$

Chapter 2: Functions and Their Graphs

$$= \frac{2x^2 + 4xh + 2h^2 + x + h - 2x^2 - x}{h}$$

$$= \frac{4xh + 2h^2 + h}{h}$$

$$= 4x + 2h + 1$$


b. When $x = 1$,

$$h = 0.5 \Rightarrow m_{\text{sec}} = 4 \cdot 1 + 2(0.5) + 1 = 6$$

Section 2.3: Properties of Functions

- c. Using point $(1, f(1)) = (1, 0)$ and slope = 1.02, we get the secant line:
 $y - 0 = 1.02(x - 1)$
 $y = 1.02x - 1.02$

d. Graphing:



86. $f(x) = -x^2 + 3x - 2$

a.
$$m_{\text{sec}} = \frac{f(x+h) - f(x)}{h}$$

$$= \frac{-(x+h)^2 + 3(x+h) - 2 - (-x^2 + 3x - 2)}{h}$$

$$= \frac{-(x^2 + 2xh + h^2) + 3x + 3h - 2 + x^2 - 3x + 2}{h}$$

$$= \frac{-x^2 - 2xh - h^2 + 3x + 3h - 2 + x^2 - 3x + 2}{h}$$

$$= \frac{-2xh - h^2 + 3h}{h}$$

$= -2x - h + 3$

b. When $x = 1$,
 $h = 0.5 \Rightarrow m_{\text{sec}} = -2 \cdot 1 - 0.5 + 3 = 0.5$


$h = 0.1 \Rightarrow m_{\text{sec}} = -2 \cdot 1 - 0.1 + 3 = 0.9$

$h = 0.01 \Rightarrow m_{\text{sec}} = -2 \cdot 1 - 0.01 + 3 = 0.99$

as $h \rightarrow 0$, $m_{\text{sec}} \rightarrow -2 \cdot 1 - 0 + 3 = 1$

c. Using point $(1, f(1)) = (1, 0)$ and slope = 0.99, we get the secant line:
 $y - 0 = 0.99(x - 1)$
 $y = 0.99x - 0.99$

d. Graphing:



a.
$$m_{\text{sec}} = \frac{f(x+h) - f(x)}{h}$$

$$= \frac{\left(\frac{1}{x+h} - \frac{1}{x}\right) - \left(\frac{x - (x+h)}{(x+h)x}\right)}{h}$$

$$= \frac{\left(\frac{x - (x+h)}{(x+h)x}\right) - \left(\frac{x - (x+h)}{(x+h)x}\right)}{h}$$

$$= \frac{\left(\frac{x - x - h}{(x+h)x}\right) - \left(\frac{-h}{(x+h)x}\right)}{h}$$

$$= -\frac{1}{(x+h)x}$$


b. When $x = 1$,
 $h = 0.5 \Rightarrow m_{\text{sec}} = -\frac{1}{(1+0.5)(1)} = -\frac{1}{1.5} = -\frac{2}{3} \approx -0.667$
 $h = 0.1 \Rightarrow m_{\text{sec}} = -\frac{1}{(1+0.1)(1)} = -\frac{1}{1.1} = -\frac{10}{11} \approx -0.909$
 $h = 0.01 \Rightarrow m_{\text{sec}} = -\frac{1}{(1+0.01)(1)} = -\frac{1}{1.01} = -\frac{100}{101} \approx -0.990$

as $h \rightarrow 0$,
 $m_{\text{sec}} \rightarrow -\frac{1}{(1+0)(1)} = -\frac{1}{1} = -1$

c. Using point $(1, f(1)) = (1, 1)$ and slope = $-\frac{100}{101}$, we get the secant line:
 $y - 1 = -\frac{100}{101}(x - 1)$
 $y - 1 = -\frac{100}{101}x + \frac{100}{101}$

$y = -\frac{100}{101}x + \frac{201}{101}$

d. Graphing:



87. $f(x) = \frac{1}{x}$

88. $f(x) = \frac{1}{x^2}$

a.
$$m_{\text{sec}} = \frac{f(x+h) - f(x)}{h}$$

$$= \frac{\left(\frac{1}{(x+h)^2} - \frac{1}{x^2} \right)}{h}$$

$$= \frac{(x^2 - (x+h)^2)}{h(x+h)^2 x^2}$$

$$= \frac{(x^2 - (x^2 + 2xh + h^2)) \left(\frac{1}{h} \right)}{(x+h)^2 x^2}$$

$$= \frac{(-2xh - h^2) \left(\frac{1}{h} \right)}{(x+h)^2 x^2}$$

$$= \frac{-2x - h}{(x+h)^2 x^2} = \frac{-2x - h}{(x^2 + 2xh + h^2) x^2}$$

b. When $x = 1$,

$$h = 0.5 \Rightarrow m_{\text{sec}} = \frac{-2 \cdot 1 - 0.5}{(1 + 0.5)^2 1^2} = -\frac{10}{9} \approx -1.1111$$

$$h = 0.1 \Rightarrow m_{\text{sec}} = \frac{-2 \cdot 1 - 0.1}{(1 + 0.1)^2 1^2} = -\frac{210}{121} \approx -1.7355$$


$$h = 0.01 \Rightarrow m_{\text{sec}} = \frac{-2 \cdot 1 - 0.01}{(1 + 0.01)^2 1^2}$$

$$= -\frac{20,100}{10,201} \approx -1.9704$$


$$\text{as } h \rightarrow 0, m_{\text{sec}} \rightarrow \frac{-2 \cdot 1 - 0}{(1 + 0)^2 1^2} = -2$$

c. Using point $(1, f(1)) = (1, 1)$ and slope = -1.9704 , we get the secant line:

d. Graphing:




89. Answers will vary. One possibility follows:



90. Answers will vary. See solution to Problem 89 for one possibility.

91. A function that is increasing on an interval can have at most one x -intercept on the interval. The graph of f could not "turn" and cross it again or it would start to decrease.

92. An increasing function is a function whose graph goes up as you read from left to right.




$$y = -1.9704x + 2.9704$$

-3

3



-5

A decreasing function is a function whose graph goes down as you read from left to right.



93. To be an even function we need $f(-x) = f(x)$
 and to be an odd function we need $f(-x) = -f(x)$. In order for a function be both even and odd, we would need $f(x) = -f(x)$.
 This is only possible if $f(x) = 0$.

94. The graph of $y = 5$ is a horizontal line.




The local maximum is $y = 5$ and it occurs at each x -value in the interval.

95. Not necessarily. It just means $f(5) > f(2)$.
 The function could have both increasing and decreasing intervals.

96.
$$\frac{f(x_2) - f(x_1)}{x_2 - x_1} = \frac{b - b}{x_2 - x_1} = 0$$


Section 2.4

1. $y = \sqrt{x}$



1

2. $y = -\frac{1}{x}$



3. $y = x^3 - 8$

y-intercept:

Let $x = 0$, then $y = (0)^3 - 8 = -8$.

x-intercept:

Let $y = 0$, then $0 = x^3 - 8$

$x^3 = 8$

$x = 2$

The intercepts are $(0, -8)$ and $(2, 0)$.

4. $(-\infty, 0)$

5. piecewise-defined

Chapter 2: Functions and Their Graphs

$$\frac{f(2) - f(-2)}{2 - (-2)} = \frac{0 - 0}{4} = 0$$

$$2 - (-2) \quad 4$$


97 – 98. Interactive Exercises

Section 2.3: Properties of Functions


6. True
7. False; the cube root function is odd and increasing on the interval $(-\infty, \infty)$.
8. False; the domain and range of the reciprocal function are both the set of real numbers except for 0.

- 9. C
- 10. A
- 11. E
- 12. G
- 13. B
- 14. D
- 15. F
- 16. H

17. $f(x) = x$




18. $f(x) = x^2$




19. $f(x) = x^3$


20. $f(x) = \sqrt{x}$




21. $f(x) = \frac{1}{x}$




22. $f(x) = |x|$




23. $f(x) = \sqrt[3]{x}$





24. $f(x) = 3$



25. a. $f(-2) = (-2)^2 = 4$

b. $f(0) = 2$

c. $f(2) = 2(2) + 1 = 5$

26. a. $f(-2) = -3(-2) = 6$

b. $f(-1) = 0$

c. $f(0) = 2(0)^2 + 1 = 1$

27. a. $f(0) = 2(0) - 4 = -4$

b. $f(1) = 2(1) - 4 = -2$

c. $f(2) = 2(2) - 4 = 0$

d. $f(3) = (3)^3 - 2 = 25$

28. a. $f(-1) = (-1)^3 = -1$

b. $f(0) = (0)^3 = 0$


c. $f(1) = 3(1) + 2 = 5$

d. $f(3) = 3(3) + 2 = 11$

29. $f(x) = \begin{cases} 2x & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$

The only intercept is $(0, 1)$.

c. Graph:



d. Range: $\{y \mid y \neq 0\}; (-\infty, 0) \cup (0, \infty)$

e. The graph is not continuous. There is a jump at $x = 0$.

30. $f(x) = \begin{cases} 3x & \text{if } x \neq 0 \\ 4 & \text{if } x = 0 \end{cases}$


a. Domain: $\{x \mid x \text{ is any real number}\}$

b. x -intercept: none

y -intercept: $f(0) = 4$

The only intercept is $(0, 4)$.

c. Graph:



d. Range: $\{y \mid y \neq 0\}; (-\infty, 0) \cup (0, \infty)$

a. Domain: $\{x \mid x \text{ is any real number}\}$

Chapter 2: Functions and Their Graphs

- b.** x -intercept: none
 y -intercept:
 $f(0) = 1$

Section 2.4: Library of Functions; Piecewise-defined Functions

- e.** The graph is not continuous. There is a jump at $x = 0$.

31. $f(x) = \begin{cases} -2x + 3 & \text{if } x < 1 \\ 3x - 2 & \text{if } x \geq 1 \end{cases}$


- a.** Domain: $\{x \mid x \text{ is any real number}\}$

b. x -intercept: none

y -intercept: $f(0) = -2(0) + 3 = 3$

The only intercept is $(0, 3)$.

c. Graph:



d. Range: $\{y \mid y \geq 1\}$; $[1, \infty)$

e. The graph is continuous. There are no holes or gaps.

32. $f(x) = \begin{cases} x + 3 & \text{if } x < -2 \\ -2x - 3 & \text{if } x \geq -2 \end{cases}$

a. Domain: $\{x \mid x \text{ is any real number}\}$

b. $x + 3 = 0$ $-2x - 3 = 0$
 $x = -3$ $-2x = 3$

$$x = -\frac{3}{2}$$


x -intercepts: $-3, -\frac{3}{2}$

2

y -intercept: $f(0) = -2(0) - 3 = -3$

The intercepts are $(-3, 0)$, $(-\frac{3}{2}, 0)$, and $(0, -3)$.

c. Graph:



e. The graph is continuous. There are no holes or gaps.

33. $f(x) = \begin{cases} x + 3 & \text{if } -2 \leq x < 1 \\ 5 & \text{if } x = 1 \\ -x + 2 & \text{if } x > 1 \end{cases}$


a. Domain: $\{x \mid x \geq -2\}$; $[-2, \infty)$

b. $x + 3 = 0$ $-x + 2 = 0$
 $x = -3$ $-x = -2$
(not in domain) $x = 2$
 x -intercept: 2

y -intercept: $f(0) = 0 + 3 = 3$

The intercepts are $(2, 0)$ and $(0, 3)$.

c. Graph:



d. Range: $\{y \mid y < 4, y = 5\}$; $(-\infty, 4) \cup \{5\}$

e. The graph is not continuous. There is a jump at $x = 1$.

34. $f(x) = \begin{cases} 2x + 5 & \text{if } -3 \leq x < 0 \\ -3 & \text{if } x = 0 \\ -5x & \text{if } x > 0 \end{cases}$

a. Domain: $\{x \mid x \geq -3\}$; $[-3, \infty)$

b. $2x + 5 = 0$ $x = -\frac{5}{2}$
 $2x = -5$

Chapter 2: Functions and Their Graphs

$$-5x = 0$$

$$x = 0$$

Section 2.4: Library of Functions; Piecewise-defined Functions

(not in domain of piece)


$$x\text{-intercept: } -\frac{5}{2}$$

$$y\text{-intercept: } f(0) = -3$$

The intercepts are $\left(-\frac{5}{2}, 0\right)$ and $(0, -3)$.

d. Range: $\{y \mid y \leq 1\}; (-\infty, 1]$

c. Graph:



d. Range: $\{y \mid y < 5\}; (-\infty, 5)$

e. The graph is not continuous. There is a jump at $x = 0$.

35. $f(x) = \begin{cases} 1+x & \text{if } x < 0 \\ x^2 & \text{if } x \geq 0 \end{cases}$

a. Domain: $\{x \mid x \text{ is any real number}\}$


b. $1+x=0 \quad x^2=0$
 $x=-1 \quad x=0$

x -intercepts: $-1, 0$

y -intercept: $f(0) = 0^2 = 0$

The intercepts are $(-1, 0)$ and $(0, 0)$.

c. Graph:



36.

b. $\frac{1}{x} = 0 \quad \sqrt[3]{x} = 0$


(no solution) $x = 0$

x -intercept: 0

y -intercept: $f(0) = \sqrt[3]{0} = 0$

The only intercept is $(0, 0)$.

c. Graph:



d. Range: $\{y \mid y \text{ is any real number}\}$

e. The graph is not continuous. There is a break at $x = 0$.

37. $f(x) = \begin{cases} |x| & \text{if } -2 \leq x < 0 \\ x^3 & \text{if } x > 0 \end{cases}$

a. Domain: $\{x \mid -2 \leq x < 0 \text{ and } x > 0\}$ or

$\{x \mid x \geq -2, x \neq 0\}; [-2, 0) \cup (0, \infty)$.

b. x -intercept: none


There are no x -intercepts since there are no values for x such that $f(x) = 0$.

y -intercept:

d. Range: $\{y \mid y \text{ is any real number}\}$

e. There is no y-intercept since $x = 0$ is not in the domain.

c. Graph:



The graph is not continuous. There is a jump at $x = 0$.

$$f(x) = \begin{cases} x & \text{if } x < 0 \\ \sqrt[3]{x} & \text{if } x \geq 0 \end{cases}$$

if $x < 0$

$$\sqrt[3]{x} \quad \text{if } x \geq 0$$

a. Domain: $\{x \mid x \text{ is any real number}\}$

d. Range: $\{y \mid y > 0\}; (0, \infty)$

- e. The graph is not continuous. There is a hole at $x = 0$.

38. $f(x) = \begin{cases} 2 - x & \text{if } -3 \leq x < 1 \\ \sqrt{x} & \text{if } x > 1 \end{cases}$

- a. Domain: $\{x \mid -3 \leq x < 1 \text{ and } x > 1\}$ or

$\{x \mid x \geq -3, x \neq 1\}; [-3, 1) \cup (1, \infty)$.


- b. $2 - x = 0 \quad \sqrt{x} = 0$
 $x = 2 \quad x = 0$
 (not in domain of piece)

no x -intercepts

y -intercept: $f(0) = 2 - 0 = 2$

The intercept is $(0, 2)$.

- c. Graph:




- d. Range: $\{y \mid y > 1\}; (1, \infty)$
 e. The graph is not continuous. There is a hole at $x = 1$.

39. $f(x) = 2 \text{int}(x)$

- a. Domain: $\{x \mid x \text{ is any real number}\}$
 b. x -intercepts:
 All values for x such that $0 \leq x < 1$.

- c. Graph:



- d. Range: $\{y \mid y \text{ is an even integer}\}$
 e. The graph is not continuous. There is a jump at each integer value of x .

40. $f(x) = \text{int}(2x)$


- a. Domain: $\{x \mid x \text{ is any real number}\}$

- b. x -intercepts:
 All values for x such that $0 \leq x < \frac{1}{2}$.

y -intercept: $f(0) = \text{int}(2(0)) = \text{int}(0) = 0$

The intercepts are all ordered pairs $(x, 0)$ when $0 \leq x < \frac{1}{2}$.

- c. Graph:



- d. Range: $\{y \mid y \text{ is an integer}\}$
 e. The graph is not continuous. There is a jump $\frac{k}{2}$

Chapter 2: Functions and Their Graphs**Section 2.4: Library of Functions; Piecewise-defined Functions**

y-intercept: $f(0) = 2 \operatorname{int}(0) = 0$

at each $x = \frac{k}{2}$, where k is an integer.

The intercepts are all ordered pairs $(x, 0)$

when $0 \leq x < 1$.

41. Answers may vary. One possibility follows:

$$f(x) = \begin{cases} -x & \text{if } -1 \leq x \leq 0 \\ 1 - x & \text{if } 0 < x \leq 2 \end{cases}$$

42. Answers may vary. One possibility follows:

$$f(x) = \begin{cases} x & \text{if } -1 \leq x \leq 0 \\ 1 - x & \text{if } 0 < x \leq 2 \end{cases}$$

43. Answers may vary. One possibility follows:

$$f(x) = \begin{cases} -x & \text{if } x \leq 0 \\ -x + 2 & \text{if } 0 < x \leq 2 \end{cases}$$

44. Answers may vary. One possibility follows:

$$f(x) = \begin{cases} 2x + 2 & \text{if } -1 \leq x \leq 0 \\ x & \text{if } x > 0 \end{cases}$$

45. a. $f(1.2) = \text{int}(2(1.2)) = \text{int}(2.4) = 2$

b. $f(1.6) = \text{int}(2(1.6)) = \text{int}(3.2) = 3$

c. $f(-1.8) = \text{int}(2(-1.8)) = \text{int}(-3.6) = -4$

46. a. $f(1.2) = \text{int}\left(\frac{1.2}{2}\right) = \text{int}(0.6) = 0$

b. $f(1.6) = \text{int}\left(\frac{1.6}{2}\right) = \text{int}(0.8) = 0$

c. $f(-1.8) = \text{int}\left(\frac{-1.8}{2}\right) = \text{int}(-0.9) = -1$

47. $C = \begin{cases} 39.99 & \text{if } 0 < x \leq 450 \\ 0.45x - 162.51 & \text{if } x > 450 \end{cases}$

a. $C(200) = \$39.99$

b. $C(465) = 0.45(465) - 162.51 = \46.74

c. $C(451) = 0.45(451) - 162.51 = \40.44

48. $F(x) = \begin{cases} 2 & \text{if } 0 < x \leq 1 \\ 4 & \text{if } 1 < x \leq 3 \\ 10 & \text{if } 3 < x \leq 4 \\ 5 \text{int}(x+1) + 2 & \text{if } 4 < x < 9 \\ 51 & \text{if } 9 \leq x \leq 24 \end{cases}$

a. $F(2) = 4$

Parking for 2 hours costs \$4.

b. $F(7) = 5 \text{int}(7+1) + 2 = 42$

Parking for 7 hours costs \$42.

d. $24 \text{ min} \cdot \frac{1 \text{ hr}}{60 \text{ min}} = 0.4 \text{ hr}$

$$F(8.4) = 5 \text{int}(8.4 + 1) + 2 = 5(9) + 2 = 47$$

Parking for 8 hours and 24 minutes costs \$47.

49. a. Charge for 50 therms:

$$C = 18.95 + 0.5038(50) + 0.33372(50) = \$60.83$$

b. Charge for 500 therms:

$$C = 18.95 + 0.5038(500) + 0.12360(450) + 0.33372(50) = \$343.16$$

c. For $0 \leq x \leq 50$:

$$C = 18.95 + 0.33372x + 0.5038x = 0.83752x + 18.95$$


For $x > 50$:

$$C = 18.95 + 0.33372(50) + 0.12360(x - 50) + 0.5038x = 18.95 + 16.686 + 0.12360x - 6.18 + 0.5038x = 0.6274x + 29.456$$

The monthly charge function:

$$C = \begin{cases} 0.83752x + 18.95 & \text{for } 0 \leq x \leq 50 \\ 0.6274x + 29.456 & \text{for } x > 50 \end{cases}$$

d. Graph:



c. $F(15) = 39.99$
Parking for 15 hours costs \$39.99

50. a. Charge for
40 therms:

$$C = 13.55 + 0.1473(20) + \\ 0.0579(20) \\ + 0.51(40)$$

=

\$

3

8

.

0

5

b. Charge for 100 therms:

$$\begin{aligned}
 C &= 13.55 + 0.1473(20) + 0.0579(30) \\
 &\quad + 0.0519(100) + 0.51(150) \\
 &= \$99.92
 \end{aligned}$$

c. For $0 \leq x \leq 20$:

$$\begin{aligned}
 C &= 13.55 + 0.1473x + 0.51x \\
 &= 0.6573x + 13.55
 \end{aligned}$$

For $20 < x \leq 50$:

$$\begin{aligned}
 C &= 13.55 + 0.1473(20) + 0.0579(x - 20) \\
 &\quad + 0.51x \\
 &= 13.55 + 2.946 + 0.0579x - 1.158 \\
 &\quad + 0.51x \\
 &= 0.5679x + 15.388
 \end{aligned}$$


For $x > 50$:

$$\begin{aligned}
 C &= 13.55 + 0.1473(20) + 0.0579(30) \\
 &\quad + 0.0519(x - 50) + 0.51x \\
 &= 13.55 + 2.946 + 1.737 + 0.0519x - 2.595 \\
 &\quad + 0.51x \\
 &= 0.5619x + 15.638
 \end{aligned}$$

The monthly charge function:

$$C(x) = \begin{cases} 0.6573x + 13.55 & \text{if } 0 \leq x \leq 20 \\ 0.5679x + 15.388 & \text{if } 20 < x \leq 50 \\ 0.5619x + 15.638 & \text{if } x > 50 \end{cases}$$

d. Graph:



51. For schedule X:

$$f(x) = \begin{cases} 0.10x & \text{if } 0 < x \leq 8500 \\ 850.00 + 0.15(x - 8500) & \text{if } 8500 < x \leq 34,500 \\ 4750.00 + 0.25(x - 34,500) & \text{if } 34,500 < x \leq 83,600 \\ 17,025.00 + 0.28(x - 83,600) & \text{if } 83,600 < x \leq 174,400 \\ 42,449.00 + 0.33(x - 174,400) & \text{if } 174,400 < x \leq 379,150 \\ 110,016.50 + 0.35(x - 379,150) & \text{if } x > 379,150 \end{cases}$$


52. For Schedule Y - 1:

$$f(x) = \begin{cases} 0.10x & \text{if } 0 < x \leq 17,000 \\ 1700.00 + 0.15(x - 17,000) & \text{if } 17,000 < x \leq 69,000 \\ 9500.00 + 0.25(x - 69,000) & \text{if } 69,000 < x \leq 139,350 \\ 27,087.50 + 0.28(x - 139,350) & \text{if } 139,350 < x \leq 212,300 \\ 47,513.50 + 0.33(x - 212,300) & \text{if } 212,300 < x \leq 379,150 \\ 102,574.00 + 0.35(x - 379,150) & \text{if } x > 379,150 \end{cases}$$

53. a. Let x represent the number of miles and C be the cost of transportation.

$$C(x) = \begin{cases} 0.50x & \text{if } 0 \leq x \leq 100 \\ 0.50(100) + 0.40(x - 100) & \text{if } 100 < x \leq 400 \\ 0.50(100) + 0.40(300) + 0.25(x - 400) & \text{if } 400 < x \leq 800 \\ 0.50(100) + 0.40(300) + 0.25(400) + 0(x - 800) & \text{if } 800 < x \leq 960 \end{cases}$$

$$C(x) = \begin{cases} 0.50x & \text{if } 0 \leq x \leq 100 \\ 10 + 0.40x & \text{if } 100 < x \leq 400 \\ 70 + 0.25x & \text{if } 400 < x \leq 800 \\ 270 & \text{if } 800 < x \leq 960 \end{cases}$$




b. For hauls between 100 and 400 miles the cost is: $C(x) = 10 + 0.40x$.

c. For hauls between 400 and 800 miles the cost is: $C(x) = 70 + 0.25x$.

54. Let x = number of days car is used. The cost of renting is given by

$$C(x) = \begin{cases} 95 & \text{if } x = 7 \\ 119 & \text{if } 7 < x \leq 8 \\ 143 & \text{if } 8 < x \leq 9 \\ 167 & \text{if } 9 < x \leq 10 \\ 190 & \text{if } 10 < x \leq 14 \end{cases}$$




55. a. Let s = the credit score of an individual who wishes to borrow \$300,000 with an 80% LTV ratio. The adverse market delivery charge is given by

$$C(s) = \begin{cases} 9750 & \text{if } s \leq 659 \\ 8250 & \text{if } 660 \leq s \leq 679 \\ 4500 & \text{if } 680 \leq s \leq 699 \\ 3000 & \text{if } 700 \leq s \leq 719 \\ 1500 & \text{if } 720 \leq s \leq 739 \\ 750 & \text{if } s \geq 740 \end{cases}$$

- b. 725 is between 720 and 739 so the charge would be \$1500.
 c. 670 is between 660 and 679 so the charge would be \$8250.
 56. Let x = the amount of the bill in dollars. The minimum payment due is given by

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < 10 \\ 10 & \text{if } 10 \leq x < 500 \\ 30 & \text{if } 500 \leq x < 1000 \\ 50 & \text{if } 1000 \leq x < 1500 \\ 70 & \text{if } x \geq 1500 \end{cases}$$



57. a.

b. $\sqrt{\quad}$

- c.

- d.

b. $W = 33 - \frac{(10.45 + 10\sqrt{5} - 5)(33 - (-10))}{22.04}$

$\approx -21^\circ C$

c. $W = 33 - \frac{(10.45 + 10\sqrt{15 - 15})(33 - (-10))}{22.04}$

$\approx -34^\circ C$

d. $W = 33 - 1.5958(33 - (-10)) = -36^\circ C$

59. Let x = the number of ounces and $C(x)$ = the postage due.

For $0 < x \leq 1$: $C(x) = \$0.88$


For $1 < x \leq 2$: $C(x) = 0.88 + 0.17 = \$1.05$

For $2 < x \leq 3$: $C(x) = 0.88 + 2(0.17) = \1.22

For $3 < x \leq 4$: $C(x) = 0.88 + 3(0.17) = \1.39

⋮

For $12 < x \leq 13$: $C(x) = 0.88 + 12(0.17) = \2.92



$W = 10^\circ C$

$\frac{0}{4}$

$W = 33 - \frac{(10.45 + 10\sqrt{5} - 5)(33 - (-10))}{22.04} \approx 4^\circ C$

$W = 33 - \frac{(10.45 + 10\sqrt{15 - 15})(33 - (-10))}{22.04} \approx -3^\circ C$

$\frac{2}{2}$

$\frac{2}{2}$


60. Each graph is that of vertically.

$y = x^2$, but shifted

$W = 33$
 -1.5958
 $(33$
 $-10) =$
 $-4^\circ C$

- e. When $0 \leq v < 1.79$, the wind speed is so small that there is no effect on the temperature.
- f. When the wind speed exceeds 20, the wind chill depends only on the air temperature.


58. a. $W = -10^\circ C$



If $y = x^2 + k$, $k > 0$, the shift is up k units; if $y = x^2 - k$, $k > 0$, the shift is down k units. The graph of $y = x^2 - 4$ is the same as the graph of


$y = x^2$, but shifted down 4 units. The graph of $y = x^2 + 5$ is the graph of $y = x^2$, but shifted up 5 units.

61. Each graph is that of $y = x^2$, but shifted horizontally.




If $y = (x - k)^2$, $k > 0$, the shift is to the right k units; if $y = (x + k)^2$, $k > 0$, the shift is to the left k units. The graph of $y = (x + 4)^2$ is the same as the graph of $y = x^2$, but shifted to the left 4 units. The graph of $y = (x - 5)^2$ is the graph of $y = x^2$, but shifted to the right 5 units.


62. Each graph is that of $y = |x|$, but either compressed or stretched vertically.



If $y = k|x|$ and $k > 1$, the graph is stretched vertically; if $y = k|x|$ and $0 < k < 1$, the graph is compressed vertically. The graph of $y = \frac{1}{4}|x|$ is the same as the graph of $y = |x|$, but compressed




The graph of $y = -|x|$ is the reflection of the graph of $y = |x|$ about the x -axis.




Multiplying a function by -1 causes the graph to be a reflection about the x -axis of the original function's graph.

64. The graph of $y = \sqrt{-x}$ is the reflection about the y -axis of the graph of $y = \sqrt{x}$.



The same type of reflection occurs when graphing $y = 2x + 1$ and $y = 2(-x) + 1$.




vertically. The graph of $y = 5|x|$ is the same as the graph of $y = |x|$, but stretched vertically.

- 63.** The graph of $y = -x^2$ is the reflection of the graph of $y = x^2$ about the x -axis.

The graph of $y = f(-x)$ is the reflection about the y -axis of the graph of $y = f(x)$.

- 65.** The graph of $y = (x - 1)^3 + 2$ is a shifting of the graph of $y = x^3$ one unit to the right and two


units up. Yes, the result could be predicted.



66. The graphs of $y = x^n$, n a positive even integer,

are all U-shaped and open upward. All go through the points $(-1, 1)$, $(0, 0)$, and $(1, 1)$. As


n increases, the graph of the function is narrower for $|x| > 1$ and flatter for $|x| < 1$.



67. The graphs of $y = x^n$, n a positive odd integer,

all have the same general shape. All go through the points $(-1, -1)$, $(0, 0)$, and $(1, 1)$. As n increases, the graph of the function increases at a greater rate for $|x| > 1$ and is flatter around 0 for

$|x| < 1$.



68. $f(x) = \begin{cases} 1 & \text{if } x \text{ is rational} \\ 0 & \text{if } x \text{ is irrational} \end{cases}$

Yes, it is a function.

namely, there is an x -intercept at each irrational value of x .

$$f(-x) = 1 = f(x) \text{ when } x \text{ is rational;}$$

$$f(-x) = 0 = f(x) \text{ when } x \text{ is irrational.}$$

Thus, f is even.

The graph of f consists of 2 infinite clusters of distinct points, extending horizontally in both directions. One cluster is located 1 unit above the x -axis, and the other is located along the x -axis.

69. For $0 < x < 1$, the graph of $y = x^r$, r rational and

$r > 0$, flattens down toward the x -axis as r gets

bigger. For $x > 1$, the graph of $y = x^r$ increases at a greater rate as r gets bigger.

Section 2.5

1. horizontal; right

2. y

3. vertical; up

4. True; the graph of $y = -f(x)$ is the reflection about the x -axis of the graph of $y = f(x)$.

5. False; to obtain the graph of $y = f(x+2) - 3$ you shift the graph of $y = f(x)$ to the left 2 units and down 3 units.

6. True

7. B

8. E

Domain = $\{x \mid x \text{ is any real number}\}$ or $(-\infty, \infty)$

Range = $\{0, 1\}$ or $\{y \mid y = 0 \text{ or } y = 1\}$

Chapter 2: Functions and Their Graphs

9 1
· 0
·

H D

y-intercept: $x = 0 \Rightarrow x$ is rational $\Rightarrow y = 1$

So the y-intercept is $y = 1$.

x-intercept: $y = 0 \Rightarrow x$ is irrational

So the graph has infinitely many x-intercepts,

Section 2.5: Graphing Techniques: Transformations

11. I

12. A

13. L

14. C

15. F

16. J

17. G

18. K

19. $y = (x - 4)^3$

20. $y = (x + 4)^3$

21. $y = x^3 + 4$

22. $y = x^3 - 4$

23. $y = (-x)^3 = -x^3$

24. $y = -x^3$

25. $y = 4x^3$

26. $y = \left(\frac{1}{4}x\right)^3 = \frac{1}{64}x^3$

27. (1) $y = \sqrt{x} + 2$

(2) $y = -(\sqrt{x+2})$

(3) $y = -(-x+2) = -x-2$

28. (1) $y = -\sqrt{x}$

(2) $y = -\sqrt{x-3}$

(3) $y = -\sqrt{x-3} - 2$

29. (1) $y = -\sqrt{x}$

(2) $y = -\sqrt{x} + 2$

(3) $y = -\sqrt{x+3} + 2$

30. (1) $y = \sqrt{x} + 2$

(2) $y = \sqrt{x}$

about the y -axis. This means we change the sign of the x -coordinate for each point on the graph of $y = f(x)$. Thus, the point $(3, 6)$ would become $(-3, 6)$.

33. (c); To go from $y = f(x)$ to $y = 2f(x)$, we

stretch vertically by a factor of 2. Multiply the y -coordinate of each point on the graph of

$y = f(x)$ by 2. Thus, the point $(1, 3)$ would become $(1, 6)$.

34. (c); To go from $y = f(x)$ to $y = f(2x)$, we

compress horizontally by a factor of 2. Divide the x -coordinate of each point on the graph of $y = f(x)$ by 2. Thus, the point $(4, 2)$ would become $(2, 2)$.

35. a. The graph of $y = f(x+2)$ is the same as the graph of $y = f(x)$, but shifted 2 units to

the left. Therefore, the x -intercepts are -7 and 1 .

b. The graph of $y = f(x-2)$ is the same as the graph of $y = f(x)$, but shifted 2 units to the right. Therefore, the x -intercepts are -3 and 5 .

c. The graph of $y = 4f(x)$ is the same as the graph of $y = f(x)$, but stretched vertically by a factor of 4. Therefore, the x -intercepts are still -5 and 3 since the y -coordinate of each is 0.

d. The graph of $y = f(-x)$ is the same as the

$y = -x + 2$

Chapter 2: Functions and Their Graphs

graph of $y = f(x)$,
but
reflected
about the

$$(3) \quad y = \sqrt{-(x+3)} + 2 = \sqrt{-x-3} + 2$$

- 31.** (c); To go from $y = f(x)$ to $y = -f(x)$ we reflect about the x -axis. This means we change the sign of the y -coordinate for each point on the graph of $y = f(x)$. Thus, the point $(3, 6)$ would become $(3, -6)$.
- 32.** (d); To go from $y = f(x)$ to $y = f(-x)$, we reflect each point on the graph of $y = f(x)$

Section 2.5: Graphing Techniques: Transformations

y -axis. Therefore, the x -intercepts are 5 and -3.


- 36. a.** The graph of $y = f(x+4)$ is the same as the graph of $y = f(x)$, but shifted 4 units to the left. Therefore, the x -intercepts are -12 and -3.
- b.** The graph of $y = f(x-3)$ is the same as the graph of $y = f(x)$, but shifted 3 units to the right. Therefore, the x -intercepts are -5 and 4.

- c. The graph of $y = 2f(x)$ is the same as the graph of $y = f(x)$, but stretched vertically by a factor of 2. Therefore, the x -intercepts are still -8 and 1 since the y -coordinate of each is 0.
 - d. The graph of $y = f(-x)$ is the same as the graph of $y = f(x)$, but reflected about the y -axis. Therefore, the x -intercepts are 8 and -1 .
37. a. The graph of $y = f(x + 2)$ is the same as the graph of $y = f(x)$, but shifted 2 units to the left. Therefore, the graph of $f(x + 2)$ is increasing on the interval $(-3, 3)$.
- b. The graph of $y = f(x - 5)$ is the same as the graph of $y = f(x)$, but shifted 5 units to the right. Therefore, the graph of $f(x - 5)$ is increasing on the interval $(4, 10)$.
 - c. The graph of $y = -f(x)$ is the same as the graph of $y = f(x)$, but reflected about the x -axis. Therefore, we can say that the graph of $y = -f(x)$ must be *decreasing* on the interval $(-1, 5)$.
 - d. The graph of $y = f(-x)$ is the same as the graph of $y = f(x)$, but reflected about the y -axis. Therefore, we can say that the graph

- c. The graph of $y = -f(x)$ is the same as the graph of $y = f(x)$, but reflected about the x -axis. Therefore, we can say that the graph of $y = -f(x)$ must be *increasing* on the interval $(-2, 7)$.
- d. The graph of $y = f(-x)$ is the same as the graph of $y = f(x)$, but reflected about the y -axis. Therefore, we can say that the graph of $y = f(-x)$ must be *increasing* on the interval $(-7, 2)$.

39. $f(x) = x^2 - 1$

Using the graph of $y = x^2$, vertically shift downward 1 unit.



The domain is $(-\infty, \infty)$ and the range is $[-1, \infty)$.

2

40. $f(x) = x + 4$

Using the graph of $y = x^2$, vertically shift

Chapter 2: Functions and Their Graphs


of $y = f(-x)$ must be *decreasing* on the

interval $(-5, 1)$.

- 38. a.** The graph of $y = f(x + 2)$ is the same as the graph of $y = f(x)$, but shifted 2 units to the left. Therefore, the graph of $f(x + 2)$ is decreasing on the interval $(-4, 5)$.
- b.** The graph of $y = f(x - 5)$ is the same as the graph of $y = f(x)$, but shifted 5 units to the right. Therefore, the graph of $f(x - 5)$ is decreasing on the interval $(3, 12)$.

Section 2.5: Graphing Techniques: Transformations


upward 4 units.



The domain is $(-\infty, \infty)$ and the range is $[4, \infty)$.

41. $g(x) = x^3 + 1$


Using the graph of $y = x^3$, vertically shift upward 1 unit.



The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

42. $g(x) = x^3 - 1$

Using the graph of $y = x^3$, vertically shift downward 1 unit.




The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

43. $h(x) = \sqrt{x-2}$

Using the graph of $y = \sqrt{x}$, horizontally shift to the right 2 units.

44. $h(x) = \sqrt{x+1}$

Using the graph of $y = \sqrt{x}$, horizontally shift to the left 1 unit.




The domain is $[-1, \infty)$ and the range is $[0, \infty)$.

45. $f(x) = (x-1)^3 + 2$

Using the graph of $y = x^3$, horizontally shift to the right 1 unit [$y = (x-1)^3$], then vertically


shift up 2 units [$y = (x-1)^3 + 2$].



The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

46. $f(x) = (x+2)^3 - 3$


Using the graph of $y = x^3$, horizontally shift to the left 2 units [$y = (x+2)^3$], then vertically



The domain is $[2, \infty)$ and the range is $[0, \infty)$.

Section 2.5: Graphing Techniques: Transformations


shift down 3 units $\lceil y = (x + 2)^3 - 3 \rceil$.



The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

47. $g(x) = 4\sqrt{x}$


Using the graph of $y = \sqrt{x}$, vertically stretch by a factor of 4.



The domain is $[0, \infty)$ and the range is $[0, \infty)$.


48. $g(x) = \frac{1}{2}\sqrt{x}$

Using the graph of $y = \sqrt{x}$, vertically compress by a factor of $\frac{1}{2}$.



49.


by a factor of $\frac{1}{2}$.



The domain is $(-\infty, 0) \cup (0, \infty)$ and the range is $(-\infty, 0) \cup (0, \infty)$.

50. $h(x) = \sqrt[3]{2x}$

Using the graph of $y = \sqrt[3]{x}$, horizontally compress by a factor of 2.



The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

51. $f(x) = -\sqrt[3]{x}$


Using the graph of $y = \sqrt[3]{x}$, reflect the graph about the x -axis.

The domain is $[0, \infty)$ and the range is $[0, \infty)$.

Chapter 2: Functions and Their Graphs

Section

5: Transformations




$f(x)$
 $\sqrt{x-1}$
 The domain is $(-\infty, \infty)$ and the range is

$$2x \quad (2)(x)$$

$$(-\infty, \infty)$$


Using the graph of $y = \frac{1}{x}$, vertically compress

52. $f(x) = -\sqrt{x}$
 Using the graph of $y = \sqrt{x}$, reflect the graph about the x -axis.




The domain is $[0, \infty)$ and the range is $(-\infty, 0]$.

53. $g(x) = \sqrt[3]{-x}$
 Using the graph of $y = \sqrt[3]{x}$, reflect the graph about the y -axis.




The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

54. $g(x) = \frac{1}{-x}$
 Using the graph of $y = \frac{1}{x}$, reflect the graph about the y -axis.




55. $h(x) = -x^3 + 2$
 Using the graph of the x -axis $[y = -x^3]$, then shift vertically upward 2 units $[y = -x^3 + 2]$.




The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

56. $h(x) = \frac{1}{-x} + 2$
 Using the graph of $y = \frac{1}{x}$, reflect the graph about the y -axis $[y = \frac{1}{-x}]$, then shift vertically upward 2 units $[y = \frac{1}{-x} + 2]$.



The domain is $(-\infty, 0) \cup (0, \infty)$ and the range is $(-\infty, 2) \cup (2, \infty)$.


57. $f(x) = 2(x+1)^2 - 3$
 Using the graph of $y = x^2$, horizontally shift to the left 1 unit $[y = (x+1)^2]$, vertically stretch



The domain is $(-\infty, \infty)$ and the range is $[-3, \infty)$.

58. $f(x) = 3(x-2)^2 + 1$


Using the graph of $y = x^2$, horizontally shift to the right 2 units $[y = (x-2)^2]$, vertically stretch by a factor of 3 $[y = 3(x-2)^2]$, and then vertically shift upward 1 unit $[y = 3(x-2)^2 + 1]$.



The domain is $(-\infty, \infty)$ and the range is $[1, \infty)$.

59. $g(x) = 2\sqrt{x-2} + 1$


Using the graph of $y = \sqrt{x}$, horizontally shift to the right 2 units $[y = \sqrt{x-2}]$, vertically stretch by a factor of 2 $[y = 2\sqrt{x-2}]$, and vertically shift upward 1 unit $[y = 2\sqrt{x-2} + 1]$.



The domain is $[2, \infty)$ and the range is $[1, \infty)$.

60. $g(x) = 3|x+1| - 3$

Using the graph of $y = |x|$, horizontally shift to the left 1 unit $[y = |x+1|]$, vertically stretch by a factor of 3 $[y = 3|x+1|]$, and vertically shift downward 3 units $[y = 3|x+1| - 3]$.



The domain is $(-\infty, \infty)$ and the range is $[-3, \infty)$.


61. $h(x) = \sqrt{-x} - 2$

Using the graph of $y = \sqrt{x}$, reflect the graph about the y-axis $[y = \sqrt{-x}]$ and vertically shift downward 2 units $[y = \sqrt{-x} - 2]$.

Chapter 2: Functions and Their Graphs

Section 2.5: Graphing Techniques: Transformations

shift upward 1 unit $\left[y = 2\sqrt{x-2} + 1 \right]$.




The domain is $(-\infty, 0]$ and the range is $[-2, \infty)$.

62.
$$h(x) = \frac{4}{x} + 2 = 4\left(\frac{1}{x}\right) + 2$$

Stretch the graph of $y = \frac{1}{x}$ vertically by a factor of 4


of 4 $\left[y = 4 \cdot \frac{1}{x} = \frac{4}{x} \right]$ and vertically shift upward 2 units $\left[y = \frac{4}{x} + 2 \right]$.



The domain is $(-\infty, 0) \cup (0, \infty)$ and the range is $(-\infty, 2) \cup (2, \infty)$.

63.
$$f(x) = -(x+1)^3 - 1$$


Using the graph of $y = x^3$, horizontally shift to the left 1 unit $\left[y = (x+1)^3 \right]$, reflect the graph



The domain is $(-\infty, \infty)$ and the range is $(-\infty, \infty)$.

64.
$$f(x) = -4\sqrt{x-1}$$

Using the graph of $y = \sqrt{x}$, horizontally shift to the right 1 unit $\left[y = \sqrt{x-1} \right]$, reflect the graph about the x -axis $\left[y = -\sqrt{x-1} \right]$, and stretch vertically by a factor of 4 $\left[y = -4\sqrt{x-1} \right]$.



The domain is $[1, \infty)$ and the range is $(-\infty, 0]$.

65.
$$g(x) = 2|1-x| = 2|-(1+x)| = 2|x-1|$$


Using the graph of $y = |x|$, horizontally shift to the right 1 unit $\left[y = |x-1| \right]$, and vertically stretch by a factor of 2 $\left[y = 2|x-1| \right]$.

Chapter 2: Functions and Their Graphs

Section 2.5: Graphing Techniques: Transformations

about the x -axis $\lceil y = -(x+1)^3 \rceil$, and vertically


shift downward 1 unit $\lceil y = -(x+1)^3 - 1 \rceil$.



The domain is $(-\infty, \infty)$ and the range is $[0, \infty)$.

66. $g(x) = 4 \sqrt{2-x} = 4 \sqrt{-(x-2)}$


Using the graph of $y = \sqrt{x}$, reflect the graph about the y -axis [$y = \sqrt{-x}$], horizontally shift to the right 2 units [$y = \sqrt{-(x-2)}$], and vertically stretch by a factor of 4 [$y = 4\sqrt{-(x-2)}$].



The domain is $(-\infty, 2]$ and the range is $[0, \infty)$.


67. $h(x) = 2 \text{int}(x-1)$

Using the graph of $y = \text{int}(x)$, horizontally shift to the right 1 unit [$y = \text{int}(x-1)$], and vertically stretch by a factor of 2 [$y = 2 \text{int}(x-1)$].



68. $h(x) = \text{int}(-x)$


Reflect the graph of $y = \text{int}(x)$ about the y -axis.



The domain is $(-\infty, \infty)$ and the range is $\{y \mid y \text{ is an integer}\}$.


69. a. $F(x) = f(x) + 3$

Shift up 3 units.




b. $G(x) = f(x+2)$

Shift left 2 units.




The domain is $(-\infty, \infty)$ and the range is $\{y \mid y \text{ is an even integer}\}$.

- c. $P(x) = -f(x)$
Reflect about the x -axis.




d. $H(x) = f(x+1) - 2$

Shift left 1 unit and shift down 2 units.




e. $Q(x) = \frac{1}{2}f(x)$

Compress vertically by a factor of $\frac{1}{2}$.




f. $g(x) = f(-x)$

Reflect about the y-axis.




g. $h(x) = f(2x)$

Compress horizontally by a factor of $\frac{1}{2}$.




70. a. $F(x) = f(x) + 3$

Shift up 3 units.




b. $G(x) = f(x+2)$

Shift left 2 units.




c. $P(x) = -f(x)$

Reflect about the x-axis.




d. $H(x) = f(x+1) - 2$

Shift left 1 unit and shift down 2 units.




e. $Q(x) = \frac{1}{2} f(x)$

Compress vertically by a factor of $\frac{1}{2}$.




f. $g(x) = f(-x)$

Reflect about the y-axis.




g. $h(x) = f(2x)$

Compress horizontally by a factor of $\frac{1}{2}$.




71. a. $F(x) = f(x) + 3$
Shift up 3 units.


b. $G(x) = f(x + 2)$
Shift left 2 units.




c. $P(x) = -f(x)$
Reflect about the x-axis.




d. $H(x) = f(x + 1) - 2$
Shift left 1 unit and shift down 2 units.




e. $Q(x) = \frac{1}{2} f(x)$
Compress vertically by a factor of $\frac{1}{2}$.







- f. $g(x) = f(-x)$
Reflect about the y-axis.




- c. $P(x) = -f(x)$
Reflect about the x-axis.




- g. $h(x) = f(2x)$
Compress horizontally by a factor of $\frac{1}{2}$.




- d. $H(x) = f(x+1) - 2$
Shift left 1 unit and shift down 2 units.




72. a. $F(x) = f(x) + 3$
Shift up 3 units.




- e. $Q(x) = \frac{1}{2}f(x)$
Compress vertically by a factor of $\frac{1}{2}$.



- b. $G(x) = f(x+2)$
Shift left 2 units.




- f. $g(x) = f(-x)$
Reflect about the y-axis.




g. $h(x) = f(2x)$

Compress horizontally by a factor of $\frac{1}{2}$.



73. a. $f(x) = x^3 - 9x, -4 < x < 4$



b. $0 = x^3 - 9x$


$0 = x(x^2 - 9)$

$0 = x(x - 3)(x + 3)$

$x = 0, x = 3, x = -3$


The x-intercepts are $-3, 0,$ and $3.$

- c. The local minimum can be found by using the Minimum feature from the CALC menu on a TI-84 Plus graphing calculator.



The local minimum is approximately -10.39 when $x \approx 1.73.$

The local maximum can be found by using the Maximum feature from the CALC menu on a TI-84 Plus graphing calculator.



- d. From the graph above, we see that f is initially increasing, decreasing between the two extrema, and increasing again at the end. Thus, f is increasing on the interval $(-4, -1.73)$ and on the interval $(1.73, 4).$ It is decreasing on the interval $(-1.73, 1.73).$

- e. $y = f(x + 2)$ involves a shift to the left 2 units so we subtract 2 from each x-value. Therefore, we have the following:
x-intercepts: $-5, -2,$ and 1

local minimum: -10.39 when $x \approx -0.27$

local maximum: 10.39 when $x \approx -3.73$

increasing on $(-6, -3.73)$ and $(-0.27, 2);$

decreasing on $(-3.73, -0.27).$

- f. $y = 2f(x)$ involves a vertical stretch by a factor of 2 so we multiply each y-value by 2. Therefore, we have the following:

x-intercepts: $-3, 0,$ and 3

local minimum: -20.78 when $x \approx 1.73$

local maximum: 20.78 when $x \approx -1.73$

increasing on $(-4, -1.73)$ and $(1.73, 4);$

decreasing on $(-1.73, 1.73).$

- g. $y = f(-x)$ involves a reflection about the y-axis. Therefore, we have the following:

x-intercepts: $-3, 0,$ and 3

local minimum: -10.39 when $x \approx -1.73$

local maximum: 10.39 when $x \approx 1.73$

increasing on $(-1.73, 1.73);$


decreasing on $(-4, -1.73)$ and $(1.73, 4).$

74. a. $f(x) = x^3 - 4x, -3 < x < 3$

15

Chapter 2: Functions and Their Graphs

Section 2.5: Graphical Information



The local maximum is approximately 10.39 when $x \approx -1.73$.

b. $0 = x^3 - 4x$


$$0 = x(x^2 - 4)$$

$$0 = x(x-2)(x+2)$$

$$x = 0, x = 2, x = -2$$

The x-intercepts are -2, 0, and 2.

- c. The local minimum can be found by using the Minimum feature from the CALC menu on a TI-84 Plus graphing calculator.




The local minimum is approximately -3.08

when $x \approx 1.15$.

The local maximum can be found by using the Maximum feature from the CALC menu

on a TI-84 Plus graphing calculator.



The local maximum is approximately 3.08 when $x \approx -1.15$.

- d. From the graph above, we see that f is initially increasing, decreasing between the two extrema, and increasing again at the end. Thus, f is increasing on the interval $(-3, -1.15)$ and on the interval $(1.15, 3)$. It is decreasing on the interval $(-1.15, 1.15)$.
- e. $y = f(x - 4)$ involves a shift to the right 4

units so we add 4 to each x-value. Therefore, we have the following:

local minimum: -3.08 when $x \approx 0.575$

local maximum: 3.08 when $x \approx -0.575$

increasing on $(-\frac{3}{2}, -0.575)$ and $(0.575, \frac{3}{2})$;

decreasing on $(-0.575, 0.575)$.

- g. $y = -f(x)$ involves a reflection about the x-axis. Therefore, we have the following:

x-intercepts: -2, 0, and 2

local minimum: -3.08 when $x \approx -1.15$


local maximum: 3.08 when $x \approx 1.15$

increasing on $(-1.15, 1.15)$;

decreasing on $(-3, -1.15)$ and $(1.15, 3)$.


75. $f(x) = x^2 + 2x$
 $f(x) = (x^2 + 2x + 1) - 1$
 $f(x) = (x + 1)^2 - 1$

Using $f(x) = x^2$, shift left 1 unit and shift down 1 unit.



76. $f(x) = x^2 - 6x$
 $f(x) = (x^2 - 6x + 9) - 9$
 $f(x) = (x - 3)^2 - 9$

Using $f(x) = x^2$, shift right 3 units and shift



Chapter 2: Functions and Their Graphs

Section 2.5: Graphing Techniques: Transformations

down 9 units.

local minimum: -3.08 when $x \approx 5.15$

local maximum: 3.08 when $x \approx 2.85$

increasing on $(1, 2.85)$ and $(5.15, 7)$;

decreasing on $(2.85, 5.15)$.

- f. $y = f(2x)$ involves a horizontal compression by a factor of 2 so we divide each x-value by 2. Therefore, we have the following:
x-intercepts: -1 , 0 , and 1


77. $f(x) = x^2 - 8x + 1$

$$f(x) = (x^2 - 8x + 16) + 1 - 16$$

$$f(x) = (x - 4)^2 - 15$$

Using $f(x) = x^2$, shift right 4 units and shift

down 15 units.




78. $f(x) = x^2 + 4x + 2$

$$f(x) = (x^2 + 4x + 4) + 2 - 4$$

$$f(x) = (x + 2)^2 - 2$$

Using $f(x) = x^2$, shift left 2 units and shift

down 2 units.



79. $f(x) = 2x^2 - 12x + 19$


$$= 2(x^2 - 6x) + 19$$

$$= 2(x^2 - 6x + 9) + 19 - 18$$

$$= 2(x - 3)^2 + 1$$

Using $f(x) = x^2$, shift right 3 units, vertically


stretch by a factor of 2, and then shift up 1 unit.



80. $f(x) = 3x^2 + 6x + 1$
 $= 3(x^2 + 2x) + 1$
 $= 3(x^2 + 2x + 1) + 1 - 3$
 $= 3(x + 1)^2 - 2$

Using $f(x) = x^2$, shift left 1 unit, vertically


stretch by a factor of 3, and shift down 2 units.




81. $f(x) = -3x^2 - 12x - 17$
 $= -3(x^2 + 4x) - 17$
 $= -3(x^2 + 4x + 4) - 17 + 12$
 $= -3(x + 2)^2 - 5$

Using $f(x) = x^2$, shift left 2 units, stretch

vertically by a factor of 3, reflect about the x-axis, and shift down 5 units.





86. a. $R(0) = 170.7(0)^2 + 1373(0) + 1080 = 1080$
 The estimated worldwide music revenue for 2005 is \$1080 million.

$$R(3) = 170.7(3)^2 + 1373(3) + 1080 = 6735.3$$

The estimated worldwide music revenue for 2008 is \$6735.3 million.

$$R(5) = 170.7(5)^2 + 1373(5) + 1080 = 12,212.5$$

The estimated worldwide music revenue for 2010 is \$12,212.5 million.

b. $r(x) = R(x - 5)$

$$= 170.7(x - 5)^2 + 1373(x - 5) + 1080$$

$$= 170.7(x^2 - 10x + 25) + 1373(x - 5) + 1080$$

$$= 170.7x^2 - 1707x + 4267.5 + 1373x - 6865 + 1080$$

$$= 170.7x^2 - 334x - 1517.5$$

c. The graph of $r(x)$ is the graph of $R(x)$ shifted 5 units to the left. Thus, $r(x)$ represents the estimated worldwide music revenue, x years after 2000.

$$r(5) = 170.7(5)^2 - 334(5) - 1517.5 = 1080$$

The estimated worldwide music revenue for 2005 is \$1080 million.

$$r(8) = 170.7(8)^2 - 334(8) - 1517.5 = 6735.3$$


The estimated worldwide music revenue for 2008 is \$6735.3 million.

$$r(10) = 170.7(10)^2 - 334(10) - 1517.5 = 12,212.5$$

The estimated worldwide music revenue for 2010 is \$12,212.5 million.


- d. In $r(x)$, x represents the number of years after 2000 (see the previous part).
- e. Answers will vary. One advantage might be that it is easier to determine what value should be substituted for x when using $r(x)$ instead of $R(x)$ to estimate worldwide music revenue.

87. $F = \frac{9}{5}C + 32$




$$F = \frac{9}{5}(K - 273) + 32$$

Shift the graph 273 units to the right.




88. a. $T = 2\pi \sqrt{\frac{l}{g}}$




b. $T_1 = 2\pi \sqrt{\frac{l+1}{g}}$; $T_2 = 2\pi \sqrt{\frac{l+2}{g}}$;

$T_3 = 2\pi \sqrt{\frac{l+3}{g}}$




c. As the length of the pendulum increases, the period increases.

d. $T_1 = 2\pi \sqrt{\frac{2l}{g}}$; $T_2 = 2\pi \sqrt{\frac{3l}{g}}$; $T_3 = 2\pi \sqrt{\frac{4l}{g}}$



e. If the length of the pendulum is multiplied by k , the period is multiplied by \sqrt{k} .

89. a. $p(x) = -0.05x^2 + 100x - 2000$




b. Select the 10% tax since the profits are higher.

c. The graph of Y1 is obtained by shifting the graph of $p(x)$ vertically down 10,000 units.


0.9. Thus, Y2 is the graph of $p(x)$

vertically compressed by a factor of 0.9.


d. Select the 10% tax since the graph of $Y1 = 0.9p(x) \geq Y2 = -0.05x^2 + 100x - 6800$ for all x in the domain.



90. a. $y = |f(x)|$




b. $y = f(|x|)$



91. a. To graph $y = |f(x)|$, the part of the graph

for f that lies in quadrants III or IV is reflected about the x -axis.




Chapter 2: Functions and Their Graphs

Section 2.5: Graphing Techniques: Transformations

The graph of Y_2 is obtained by multiplying the y -coordinate of the graph of $p(x)$ by

- b. To graph $y = f(|x|)$, the part of the graph for f that lies in quadrants II or III is replaced by the reflection of the part in quadrants I and IV reflected about the y -axis.



92. a. The graph of $y = f(x+3) - 5$ is the graph of $y = f(x)$ but shifted left 3 units and down 5 units. Thus, the point $(1, 3)$ becomes the point $(-2, -2)$.
- b. The graph of $y = -2f(x-2) + 1$ is the graph of $y = f(x)$ but shifted right 2 units, stretched vertically by a factor of 2, reflected about the x -axis, and shifted up 1 unit. Thus, the point $(1, 3)$ becomes the point $(3, -5)$.
- c. The graph of $y = f(2x+3)$ is the graph of $y = f(x)$ but shifted left 3 units and horizontally compressed by a factor of 2. Thus, the point $(1, 3)$ becomes the point $(-1, 3)$.
93. a. The graph of $y = g(x+1) - 3$ is the graph

vertically by a factor of 3, reflected about the x -axis, and shifted up 3 units. Thus, the point $(-3, 5)$ becomes the point $(1, -12)$.

- c. The graph of $y = g(3x+9)$ is the graph of $y = f(x)$ but shifted left 9 units and horizontally compressed by a factor of 3. Thus, the point $(-3, 5)$ becomes the point $(-4, 5)$.

94. The graph of $y = 4f(x)$ is a vertical stretch of the graph of f by a factor of 4, while the graph of $y = f(4x)$ is a horizontal compression of the graph of f by a factor of $\frac{1}{4}$.
95. The graph of $y = f(x) - 2$ will shift the graph of $y = f(x)$ down by 2 units. The graph of $y = f(x-2)$ will shift the graph of $y = f(x)$ to the right by 2 units.
96. The graph of $y = \sqrt{-x}$ is the graph of $y = \sqrt{x}$ but reflected about the y -axis. Therefore, our region is simply rotated about the y -axis and does not change shape. Instead of the region being bounded on the right by $x = 4$, it is bounded on the left by $x = -4$. Thus, the area of the second region would also be $\frac{16}{3}$ square units.
- 97 – 98. Answers will vary.
- 99 – 104. Interactive Exercises.

Section 2.6

Chapter 2: Functions and Their Graphs

of $y = g(x)$ but shifted left 1 unit and down 3 units. Thus, the point $(-3, 5)$

becomes the point $(-4, 2)$.

b. The graph of $y = -3g(x-4) + 3$ is the graph

of $y = g(x)$ but shifted right 4 units, stretched

c.
$$d(1) = \sqrt{(1)^4 - 15(1)^2 + 64}$$

$$= \sqrt{1 - 15 + 64} = \sqrt{50} = 5\sqrt{2} \approx 7.07$$

Section 2.6: Mathematical Models: Building Functions


1. a. The distance d from P to the origin is

$$d = \sqrt{x^2 + y^2}. \text{ Since } P \text{ is a point on the}$$


graph of $y = \sqrt{x^2 - 8}$, we have: _____

$$d(x) = x^2 + (x^2 - 8)^2 = x^4 - 15x^2 + 64$$

b. $d(0) = \sqrt{0^4 - 15(0)^2 + 64} = \sqrt{64} = 8$



e. d is smallest when $x \approx -2.74$ or when $x \approx 2.74$.




2. a. The distance d from P to $(0, -1)$ is $d = \sqrt{x^2 + (y+1)^2}$. Since P is a point on the graph of $y = x^2 - 8$, we have:

$$d(x) = \sqrt{x^2 + (x^2 - 8 + 1)^2}$$


$$= \sqrt{x^2 + (x^2 - 7)^2} = \sqrt{x^4 - 13x^2 + 49}$$

b. $d(0) = \sqrt{0^4 - 13(0)^2 + 49} = \sqrt{49} = 7$

c. $d(-1) = \sqrt{(-1)^4 - 13(-1)^2 + 49} = \sqrt{37} \approx 6.08$




e. d is smallest when $x \approx -2.55$ or when $x \approx 2.55$.




3. a. The distance d from P to the point $(1, 0)$ is

$$d = \sqrt{(x-1)^2 + y^2}$$
. Since P is a point on




c. d is smallest when $x = \frac{1}{2}$.




4. a. The distance d from P to the origin is $d = \sqrt{x^2 + y^2}$. Since P is a point on the graph of $y = \frac{1}{x}$, we have:

$$d(x) = \sqrt{x^2 + \left(\frac{1}{x}\right)^2} = \sqrt{x^2 + \frac{1}{x^2}} = \sqrt{\frac{x^4 + 1}{x^2}}$$

$$= \frac{\sqrt{x^4 + 1}}{x}$$



c. d is smallest when $x = -1$ or $x = 1$.



5. By definition, a triangle has area $A = \frac{1}{2}bh$, b = base, h = height. From the figure,

we know that $b = x$ and $h = y$. Expressing the

Chapter 2: Functions and Their Graphs

the graph of $y = \sqrt{x}$, we have:

$$d(x) = \sqrt{(x-1)^2 + (\sqrt{x})^2} = \sqrt{x^2 - x + 1}$$

where $x \geq 0$.


Section 2.6: Mathematical Models: Building Functions

area of the triangle as a function of x , we have:

$$A(x) = \frac{1}{2}xy = \frac{1}{2}x(x^3) = \frac{1}{2}x^4.$$

6. By definition, a triangle has area


$A = \frac{1}{2}bh$, b =base, h = height. Because one vertex of the triangle is at the origin and the



other is on the x -axis, we know that $b = x$ and $h = y$. Expressing the area of the

triangle as a function of x , we have:

$$A(x) = \frac{1}{2}xy = \frac{1}{2}x(9 - x^2) = \frac{9}{2}x - \frac{1}{2}x^3.$$





The perimeter is largest when $x \approx 1.79$.

7. a. $A(x) = xy = x(16 - x^2) = -x^3 + 16x$

b. Domain: $\{x \mid 0 < x < 4\}$


c. The area is largest when $x \approx 2.31$.
30



8. a. $A(x) = 2xy = 2x\sqrt{4 - x^2}$

b. $p(x) = 2(2x) + 2(y) = 4x + 2\sqrt{4 - x^2}$

c. Graphing the area equation:
4



The area is largest when $x \approx 1.41$.



d. Graphing the perimeter equation:
10

9. a. In Quadrant I, $x + y = 4 \rightarrow y = 4 - x$

$$A(x) = (2x)(2y) = 4x\sqrt{4 - x^2}$$



b. $p(x) = 2(2x) + 2(2y) = 4x + 4\sqrt{4 - x^2}$

c. Graphing the area equation:
10



The area is largest when $x \approx 1.41$.

d. Graphing the perimeter equation:
12




The perimeter is largest when $x \approx 1.41$.

10. a. $A(r) = (2r)(2r) = 4r^2$

b. $p(r) = 4(2r) = 8r$

Chapter 2: Functions and Their Graphs



Section 2.6: Mathematical Models: Building Functions

11. a. C = circumference, A = total area,

r = radius, x = side of square

$$C = 2\pi r = 10 - 4x \Rightarrow r = \frac{5-2x}{\pi}$$



$$\text{Total Area} = \text{area}_{\text{square}} + \text{area}_{\text{circle}} = x^2 + \pi r^2$$

$$A(x) = x^2 + \pi \left(\frac{5-2x}{\pi} \right)^2 = x^2 + \frac{25-20x+4x^2}{\pi}$$

- b. Since the lengths must be positive, we have:
 $10 - 4x > 0$ and $x > 0$
 $-4x > -10$ and $x > 0$
 $x < 2.5$ and $x > 0$

Domain: $\{x \mid 0 < x < 2.5\}$

- c. The total area is smallest when $x \approx 1.40$ meters.



12. a. C = circumference, A = total area,
 r = radius, x = side of equilateral triangle


$$C = 2\pi r = 10 - 3x \Rightarrow r = \frac{10-3x}{2\pi}$$

The height of the equilateral triangle is $\frac{\sqrt{3}}{2}x$.


$$\begin{aligned} \text{Total Area} &= \text{area}_{\text{triangle}} + \text{area}_{\text{circle}} \\ &= \frac{1}{2}x \left(\frac{\sqrt{3}}{2}x \right) + \pi r^2 \end{aligned}$$

$$A(x) = \frac{\sqrt{3}}{4}x^2 + \pi \left(\frac{10-3x}{2\pi} \right)^2$$

- c. The area is smallest when $x \approx 2.08$ meters.



0 3.33



13. a. Since the wire of length x is bent into a circle, the circumference is x . Therefore, $C(x) = x$.

- b. Since $C = x = 2\pi r$, $r = \frac{x}{2\pi}$.

$$A(x) = \pi r^2 = \pi \left(\frac{x}{2\pi} \right)^2 = \frac{x^2}{4\pi}$$

14. a. Since the wire of length x is bent into a square, the perimeter is x . Therefore, $p(x) = x$.

$$\frac{1}{4}$$

- b. Since $P = x = 4s$, $s = \frac{1}{4}x$, we have

$$A(x) = s^2 = \left(\frac{1}{4}x \right)^2 = \frac{1}{16}x^2$$


15. a. A = area, r = radius; diameter = $2r$

$$A(r) = (2r)(r) = 2r^2$$

- b. p = perimeter

$$p(r) = 2(2r) + 2r = 6r$$

16. C = circumference, r



Chapter 2: Functions and Their Graphs

r = radius;

$$= \frac{\sqrt{3}}{4}x^2 + \frac{100 - 60x + 9x}{4\pi}$$

b. Since the lengths must be positive, we have:

$$10 - 3x > 0 \quad \text{and} \quad x > 0$$

$$-3x > -10 \quad \text{and} \quad x > 0$$

$$x < \frac{10}{3} \quad \text{and} \quad x > 0$$

$$\text{Domain: } \left\{ x \mid 0 < x < \frac{10}{3} \right\}$$

Section 2.6: Mathematical Models: Building Functions

x = length of a side of the triangle

Since $\triangle ABC$ is equilateral, $EM = \frac{\sqrt{3}x}{2}$.

Therefore, $OM = \frac{\sqrt{3}x}{2} - OE = \frac{\sqrt{3}x}{2} - r$

In ΔOAM , $r^2 = \left(\frac{x}{2}\right)^2 + \left(\frac{\sqrt{3}x}{2} - r\right)^2$

$$r^2 = \frac{x^2}{4} + \frac{3}{4}x^2 - \sqrt{3}rx + r^2$$

$$\sqrt{3}rx = x^2$$

$$r = \frac{x}{\sqrt{3}}$$

Therefore, the circumference of the circle is

$$C(x) = 2\pi r = 2\pi\left(\frac{x}{\sqrt{3}}\right) = \frac{2\pi\sqrt{3}}{3}x$$

17. Area of the equilateral triangle

$$A = \frac{1}{2} \cdot x \cdot \frac{\sqrt{3}}{2}x = \frac{\sqrt{3}}{4}x^2$$

From problem 16, we have $r^2 = \frac{x^2}{3}$.

Area inside the circle, but outside the triangle:



$$A(x) = \pi r^2 - \frac{\sqrt{3}}{4}x^2$$

$$= \pi \frac{x^2}{3} - \frac{\sqrt{3}}{4}x^2 = \left(\frac{\pi}{3} - \frac{\sqrt{3}}{4}\right)x^2$$


18. $d^2 = d_1^2 + d_2^2$

$$d^2 = (30t)^2 + (40t)^2$$

$$d(t) = \sqrt{900t^2 + 1600t^2} = \sqrt{2500t^2} = 50t$$



b. The distance is smallest at $t \approx 0.07$ hours.



20. r = radius of cylinder, h = height of cylinder,

V = volume of cylinder

$$r^2 + \left(\frac{h}{2}\right)^2 = R^2 \Rightarrow r^2 + \frac{h^2}{4} = R^2 \Rightarrow r^2 = R^2 - \frac{h^2}{4}$$

$$V = \pi r^2 h$$

$$V(h) = \pi \left(R^2 - \frac{h^2}{4}\right)h = \pi h \left(R^2 - \frac{h^2}{4}\right)$$

21. r = radius of cylinder, h = height of cylinder,

V = volume of cylinder

By similar triangles: $\frac{H}{R} = \frac{H-h}{r}$

$$Hr = R(H-h)$$

$$Hr = RH - Rh$$

$$Rh = RH - Hr$$

$$h = \frac{RH - Hr}{R} = \frac{H(R-r)}{R}$$


$$V = \pi r^2 h = \pi r^2 \left(\frac{H(R-r)}{R}\right) = \frac{\pi H(R-r)r^2}{R}$$

Chapter 2: Functions and Their Graphs

19. a. $d^2 = d_1^2 + d_2^2$
 $d^2 = (2 - 30t)^2 + (3 - 40t)^2$
 $d(t) = \sqrt{(2 - 30t)^2 + (3 - 40t)^2}$
 $= \sqrt{4 - 120t + 900t^2 + 9 - 240t + 1600t^2}$
 $= \sqrt{2500t^2 - 360t + 13}$

Section 2.6: Mathematical Models: Building Functions

22. a. The total cost of installing the cable along the road is $500x$. If cable is installed x miles along the road, there are $5 - x$ miles between the road to the house and where the cable ends along the road.



$$d = \sqrt{(5-x)^2 + 2^2}$$


$$= \sqrt{25 - 10x + x^2 + 4} = \sqrt{x^2 - 10x + 29}$$

The total cost of installing the cable is:


$$C(x) = 500x + 700\sqrt{x^2 - 10x + 29}$$

Domain: $\{x \mid 0 \leq x \leq 5\}$


- b. $C(1) = 500(1) + 700\sqrt{1^2 - 10(1) + 29}$
 $= 500 + 700\sqrt{20} = \$3630.50$
- c. $C(3) = 500(3) + 700\sqrt{3^2 - 10(3) + 29}$
 $= 1500 + 700\sqrt{8} = \$3479.90$



- e. Using MINIMUM, the graph indicates that $x \approx 2.96$ miles results in the least cost.



- 23. a. The time on the boat is given by $\frac{d_1}{3}$. The time on land is given by $\frac{12-x}{5}$.



$$d_1 = \sqrt{x^2 + 2^2} = \sqrt{x^2 + 4}$$

The total time for the trip is:



$$T(x) = \frac{12-x}{5} + \frac{d_1}{3} = \frac{12-x}{5} + \frac{\sqrt{x^2 + 4}}{3}$$

- b. Domain: $\{x \mid 0 \leq x \leq 12\}$

- c. $T(4) = \frac{12-4}{5} + \frac{\sqrt{4^2 + 4}}{3}$
 $= \frac{8}{5} + \frac{\sqrt{20}}{3} \approx 3.09$ hours

- d. $T(8) = \frac{12-8}{5} + \frac{\sqrt{8^2 + 4}}{3}$
 $= \frac{4}{5} + \frac{\sqrt{68}}{3} \approx 3.55$ hours

- 24. Consider the diagrams shown below.




There is a pair of similar triangles in the diagram. Since the smaller triangle is similar to the larger triangle, we have the proportion

$$\frac{r}{h} = \frac{4}{16} \Rightarrow r = \frac{1}{4}h \Rightarrow r = \frac{1}{4}h$$


Substituting into the volume formula for the conical portion of water gives

$$V(h) = \frac{1}{3}\pi r^2 h = \frac{1}{3}\pi \left(\frac{1}{4}h\right)^2 h = \frac{\pi}{48}h^3.$$

25. a. length = $24 - 2x$; width = $24 - 2x$; height = x
 $V(x) = x(24 - 2x)(24 - 2x) = x(24 - 2x)^2$
- b. $V(3) = 3(24 - 2(3))^2 = 3(18)^2 = 3(324) = 972 \text{ in}^3.$
- c. $V(10) = 10(24 - 2(10))^2 = 10(4)^2 = 10(16) = 160 \text{ in}^3.$
- d. $y_1 = x(24 - 2x)^2$



Use MAXIMUM.



The volume is largest when $x = 4$ inches.



26. a. Let A = amount of material,
 x = length of the base, h = height, and
 V = volume.

$$A(1) = 1 + \frac{40}{1} = 1 + 40 = 41 \text{ ft}$$

b. $()^2 = 1$

c. $A(2) = 2^2 + \frac{40}{2} = 4 + 20 = 24 \text{ ft}^2$

d. $y_1 = x^2 + \frac{40}{x}$



The amount of material is least when $x = 2.71$ ft.

Chapter 2 Review Exercises

- This relation represents a function. Domain = $\{-1, 2, 4\}$; Range = $\{0, 3\}$.
- This relation does not represent a function, since 4 is paired with two different values.

3. $f(x) = \frac{3x}{x^2 - 1}$

a. $f(2) = \frac{3(2)}{(2)^2 - 1} = \frac{6}{4 - 1} = \frac{6}{3} = 2$

b. $f(-2) = \frac{3(-2)}{(-2)^2 - 1} = \frac{-6}{4 - 1} = \frac{-6}{3} = -2$
 $\frac{3(-x)}{-3x}$

Chapter 2: Functions and Their Graphs

$$V = x^2h = 10 \Rightarrow h = \frac{10}{x^2}$$

$$\text{Total Area } A = (\text{Area}_{\text{base}}) + (4)(\text{Area}_{\text{side}})$$

$$= x^2 + 4xh$$

$$= x^2 + 4x \left(\frac{10}{x^2} \right)$$

$$= x^2 + \frac{40}{x}$$

$$A(x) = x^2 + \frac{40}{x}$$

Chapter 2 Review Exercises

c. $f(-x) = \frac{1}{(-x)^2 - 1} = \frac{1}{x^2 - 1}$

d. $-f(x) = -\left(\frac{3x}{x^2 - 1} \right) = \frac{-3x}{x^2 - 1}$

e. $f(x-2) = \frac{3(x-2)}{(x-2)^2 - 1}$
 $= \frac{3x-6}{x^2-4x+4-1} = \frac{3(x-2)}{x^2-4x+3}$

f. $f(2x) = \frac{3(2x)}{(2x)^2 - 1} = \frac{6x}{4x^2 - 1}$

4. $f(x) = \sqrt{x^2 - 4}$

a. $f(2) = \sqrt{2^2 - 4} = \sqrt{4 - 4} = \sqrt{0} = 0$

b. $f(-2) = \sqrt{(-2)^2 - 4} = \sqrt{4 - 4} = \sqrt{0} = 0$

c. $f(-x) = \sqrt{(-x)^2 - 4} = \sqrt{x^2 - 4}$

d. $-f(x) = -\sqrt{x^2 - 4}$

e. $f(x-2) = \sqrt{(x-2)^2 - 4}$
 $= \sqrt{x^2 - 4x + 4 - 4}$
 $= \sqrt{x^2 - 4x}$

f. $f(2x) = \sqrt{(2x)^2 - 4} = \sqrt{4x^2 - 4}$
 $= \sqrt{4(x^2 - 1)} = 2\sqrt{x^2 - 1}$

5. $f(x) = \frac{x^2 - 4}{x^2}$

a. $f(2) = \frac{2^2 - 4}{2^2} = \frac{4 - 4}{4} = \frac{0}{4} = 0$

b. $f(-2) = \frac{(-2)^2 - 4}{(-2)^2} = \frac{4 - 4}{4} = \frac{0}{4} = 0$

c. $f(-x) = \frac{(-x)^2 - 4}{(-x)^2} = \frac{x^2 - 4}{x^2}$

d. $-f(x) = -\left(\frac{x^2 - 4}{x^2}\right) = \frac{4 - x^2}{x^2} = -\frac{x^2 - 4}{x^2}$

e. $f(x-2) = \frac{(x-2)^2 - 4}{(x-2)^2} = \frac{x^2 - 4x + 4 - 4}{(x-2)^2}$

6. $f(x) = \frac{x}{x^2 - 9}$

The denominator cannot be zero:

$$x^2 - 9 \neq 0$$

$$(x+3)(x-3) \neq 0$$

$$x \neq -3 \text{ or } 3$$

Domain: $\{x \mid x \neq -3, x \neq 3\}$

7. $f(x) = \sqrt{2-x}$

The radicand must be non-negative:

$$2 - x \geq 0$$

$$x \leq 2$$

Domain: $\{x \mid x \leq 2\}$ or $(-\infty, 2]$

8. $g(x) = \frac{|x|}{x}$

The denominator cannot be zero:

$$x \neq 0$$

Domain: $\{x \mid x \neq 0\}$

9. $f(x) = \frac{x}{x^2 + 2x - 3}$

The denominator cannot be zero:

$$x^2 + 2x - 3 \neq 0$$

$$(x+3)(x-1) \neq 0$$

$$x \neq -3 \text{ or } 1$$

Domain: $\{x \mid x \neq -3, x \neq 1\}$

10. $f(x) = \frac{\sqrt{x+1}}{x-4}$

The denominator cannot be zero:

$$x^2 - 4 \neq 0$$

$$x^2 \neq 4$$

$$x \neq \pm 2$$

The radicand in the numerator must be non-

Chapter 2: Functions and Their Graphs

$$= \frac{x^2 - 4x}{(x-2)^2} = \frac{x(x-4)}{(x-2)^2}$$

f. $f(2x) = \frac{(2x)^2 - 4}{(2x)^2} = \frac{4x^2 - 4}{4x^2}$
 $= \frac{4(x^2 - 1)}{4x^2} = \frac{x^2 - 1}{x^2}$

Chapter 2 Review Exercises

negative:
 $x + 1 \geq 0$
 $x \geq -1$

Domain: $\{x \mid x \geq -1, x \neq 2\}$

11. $f(x) = \frac{x}{\sqrt{x+8}}$

The denominator cannot be zero and the radicand must be non-negative:

$$x + 8 > 0$$

$$x > -8$$

Domain: $\{x \mid x > -8\}$

12. $f(x) = 2 - x \quad g(x) = 3x + 1$

$$(f + g)(x) = f(x) + g(x)$$

$$= 2 - x + 3x + 1 = 2x + 3$$

Domain: $\{x \mid x \text{ is any real number}\}$

$$(f - g)(x) = f(x) - g(x)$$

$$= 2 - x - (3x + 1)$$

$$= 2 - x - 3x - 1$$

$$= -4x + 1$$

Domain: $\{x \mid x \text{ is any real number}\}$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$= (2 - x)(3x + 1)$$

$$= 6x + 2 - 3x^2 - x$$

$$= -3x^2 + 5x + 2$$

Domain: $\{x \mid x \text{ is any real number}\}$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} = \frac{2-x}{3x+1}$$

$$3x + 1 \neq 0$$

$$3x \neq -1 \Rightarrow x \neq -\frac{1}{3}$$

Domain: $\left\{x \mid x \neq -\frac{1}{3}\right\}$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$= (3x^2 + x + 1)(3x)$$

$$= 9x^3 + 3x^2 + 3x$$

Domain: $\{x \mid x \text{ is any real number}\}$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} = \frac{3x^2 + x + 1}{3x}$$

$$3x \neq 0 \Rightarrow x \neq 0$$

Domain: $\{x \mid x \neq 0\}$

14. $f(x) = \frac{x+1}{x-1} \quad g(x) = \frac{1}{x}$

$$(f + g)(x) = f(x) + g(x)$$

$$= \frac{x+1}{x-1} + \frac{1}{x} = \frac{x(x+1) + 1(x-1)}{x(x-1)}$$

$$= \frac{x^2 + x + x - 1}{x(x-1)} = \frac{x^2 + 2x - 1}{x(x-1)}$$

Domain: $\{x \mid x \neq 0, x \neq 1\}$

$$(f - g)(x) = f(x) - g(x)$$

$$= \frac{x+1}{x-1} - \frac{1}{x} = \frac{x(x+1) - 1(x-1)}{x(x-1)}$$

$$= \frac{x^2 + x - x + 1}{x(x-1)} = \frac{x^2 + 1}{x(x-1)}$$

Domain: $\{x \mid x \neq 0, x \neq 1\}$

$$(f \cdot g)(x) = f(x) \cdot g(x) = \left(\frac{x+1}{x-1}\right)\left(\frac{1}{x}\right) = \frac{x+1}{x(x-1)}$$

Domain: $\{x \mid x \neq 0, x \neq 1\}$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} = \frac{x+1}{x-1} \cdot \frac{(x-1)(x)}{1(x-1)} = \frac{x(x+1)}{x-1}$$

Chapter 2: Functions and Their Graphs

13. $f(x) = 3x^2 + x + 1$ $g(x) = 3x$

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\ &= 3x^2 + x + 1 + 3x \\ &= 3x^2 + 4x + 1\end{aligned}$$

Domain: $\{x \mid x \text{ is any real number}\}$

$$\begin{aligned}(f - g)(x) &= f(x) - g(x) \\ &= 3x^2 + x + 1 - 3x \\ &= 3x^2 - 2x + 1\end{aligned}$$

Domain: $\{x \mid x \text{ is any real number}\}$

Chapter 2 Review Exercises

$$\left(\frac{1}{x} \right) \left(\frac{1}{x} \right) \left(\frac{1}{x} \right)$$

Domain: $\{x \mid x \neq 0, x \neq 1\}$

15. $f(x) = -2x^2 + x + 1$

$f(x+h) - f(x)$

$$\begin{aligned} &= \frac{-2(x+h)^2 + (x+h) + 1 - (-2x^2 + x + 1)}{h} \\ &= \frac{-2(x^2 + 2xh + h^2) + x + h + 1 + 2x^2 - x - 1}{h} \\ &= \frac{-2x^2 - 4xh - 2h^2 + x + h + 1 + 2x^2 - x - 1}{h} \\ &= \frac{-4xh - 2h^2 + h}{h} = \frac{h(-4x - 2h + 1)}{h} \\ &= -4x - 2h + 1 \end{aligned}$$

16. a. Domain: $\{x \mid -4 \leq x \leq 3\}$; $[-4, 3]$

Range: $\{y \mid -3 \leq y \leq 3\}$; $[-3, 3]$

b. Intercept: $(0, 0)$


c. $f(-2) = -1$

d. $f(x) = -3$ when $x = -4$

e. $f(x) > 0$ when $0 < x \leq 3$
 $\{x \mid 0 < x \leq 3\}$


f. To graph $y = f(x-3)$, shift the graph of f

horizontally 3 units to the right.




g. To graph $y = f\left(\frac{1}{2}x\right)$, stretch the graph of

f horizontally by a factor of 2.



h. To graph $y = -f(x)$, reflect the graph of f

vertically about the y-axis.



17. a. Domain: $\{x \mid x \text{ is any real number}\}$

18.

Chapter 2: Functions and Their GraphsRange: $\{y \mid y \text{ is any real number}\}$

- b. Increasing: $(-\infty, -2)$ and $(2, \infty)$; Decreasing: $(-2, 2)$
- c. Local minimum is -1 at $x = 2$; Local maximum is 1 at $x = -2$
- d. Absolute minimum is -3 at $x = -4$; Absolute maximum is 3 at $x = 4$
- e. The graph is symmetric with respect to the origin.
- f. The function is odd.
- g. x -intercepts: $-3, 0, 3$;
 y -intercept: 0

$$f(x) = x^3 - 4x$$

$$f(-x) = (-x)^3 - 4(-x) = -x^3 + 4x$$

$$= -(x^3 - 4x) = -f(x)$$

 f is odd.**Chapter 2 Review Exercises**

19. $g(x) = \frac{4+x^2}{1+x^4}$


$$g(-x) = \frac{4+(-x)^2}{1+(-x)^4} = \frac{4+x^2}{1+x^4} = g(x)$$
 g is even.

20. $G(x) = 1 - x + x^3$
 $G(-x) = 1 - (-x) + (-x)^3$
 $= 1 + x - x^3 \neq -G(x)$ or $G(x)$
 G is neither even nor odd.

21. $f(x) = \frac{x}{1+x^2}$

$$f(-x) = \frac{-x}{1+(-x)^2} = \frac{-x}{1+x^2} = -f(x)$$
 f is odd.

22. $f(x) = 2x^3 - 5x + 1$ on the interval $[-3, 3]$
 Use MAXIMUM and MINIMUM on the graph of $y_1 = 2x^3 - 5x + 1$.




local maximum: 4.04 when $x \approx -0.91$

local minimum: -2.04 when $x = 0.91$

f is increasing on: $(-3, -0.91)$ and $(0.91, 3)$;
 f is decreasing on: $(-0.91, 0.91)$.

23. $f(x) = 2x^4 - 5x^3 + 2x + 1$ on the interval $(-2, 3)$
 Use MAXIMUM and MINIMUM on the graph of $y_1 = 2x^4 - 5x^3 + 2x + 1$.



local minima: 0.54 when $x = -0.34$, -3.56 when $x = 1.80$

f is increasing on: $(-0.34, 0.41)$ and $(1.80, 3)$;

f is decreasing on: $(-2, -0.34)$ and $(0.41, 1.80)$.

24. $f(x) = 8x^2 - x$
 a. $\frac{f(2) - f(1)}{2 - 1} = \frac{8(2)^2 - 2 - [8(1)^2 - 1]}{1}$
 $= 32 - 2 - (7) = 23$

b. $\frac{f(1) - f(0)}{1 - 0} = \frac{8(1)^2 - 1 - [8(0)^2 - 0]}{1}$
 $= 8 - 1 - (0) = 7$

c. $\frac{f(4) - f(2)}{4 - 2} = \frac{8(4)^2 - 4 - [8(2)^2 - 2]}{2}$
 $= \frac{128 - 4 - (30)}{2} = \frac{94}{2} = 47$

25. $f(x) = 2 - 5x$

$$\frac{f(3) - f(2)}{3 - 2} = \frac{[2 - 5(3)] - [2 - 5(2)]}{(2 - 15) - (2 - 10)}$$

 $= \frac{-13 - (-8)}{-13 - (-8)} = \frac{-5}{-5} = 1$


26. $f(x) = 3x - 4x^2$

$$\frac{f(3) - f(2)}{3 - 2} = \frac{[3(3) - 4(3)^2] - [3(2) - 4(2)^2]}{9 - 36 - (6 - 16)}$$


 $= \frac{-27 + 10}{-17} = -17$

27. The graph does not pass the Vertical Line Test and is therefore not a function.


29. $f(x) = |x|$



30. $f(x) = \sqrt{x}$




31. $F(x) = |x| - 4$. Using the graph of $y = |x|$, vertically shift the graph downward 4 units.




Intercepts: $(-4, 0), (4, 0), (0, -4)$
 Domain: $\{x \mid x \text{ is any real number}\}$
 Range: $\{y \mid y \geq -4\}$ or $[-4, \infty)$

32. $g(x) = -2|x|$. Reflect the graph of $y = |x|$



Intercepts: $(0, 0)$
 Domain: $\{x \mid x \text{ is any real number}\}$
 Range: $\{y \mid y \leq 0\}$ or $(-\infty, 0]$

33. $h(x) = \sqrt{x-1}$. Using the graph of $y = \sqrt{x}$, horizontally shift the graph to the right 1 unit.




Intercept: $(1, 0)$
 Domain: $\{x \mid x \geq 1\}$ or $[1, \infty)$
 Range: $\{y \mid y \geq 0\}$ or $[0, \infty)$

34. $f(x) = \sqrt{1-x} = \sqrt{-(x-1)}$. Reflect the graph of $y = \sqrt{x}$ about the y-axis and horizontally shift the graph to the right 1 unit.

Chapter 2: Functions and Their Graphs

about the x -axis and vertically stretch the graph by a factor of 2.



Chapter 2 Review Exercises


Intercepts: $(1, 0)$, $(0, 1)$

Domain: $\{x \mid x \leq 1\}$ or $(-\infty, 1]$

Range: $\{y \mid y \geq 0\}$ or $[0, \infty)$

35. $h(x) = (x-1)^2 + 2$. Using the graph of $y = x^2$,

horizontally shift the graph to the right 1 unit and vertically shift the graph up 2 units.



Intercepts: (0, 3)


Domain: $\{x \mid x \text{ is any real number}\}$

Range: $\{y \mid y \geq 2\}$ or $[2, \infty)$

36. $g(x) = -2(x+2)^3 - 8$

Using the graph of $y = x^3$, horizontally shift the

graph to the left 2 units, vertically stretch the graph by a factor of 2, reflect about the x -axis, and vertically shift the graph down 8 units.



Intercepts: $(0, -24), (-2 - \sqrt[3]{4}, 0) \approx (-3.6, 0)$


Domain: $\{x \mid x \text{ is any real number}\}$

Range: $\{y \mid y \text{ is any real number}\}$

37. $f(x) = \begin{cases} 3x & \text{if } -2 < x \leq 1 \\ x+1 & \text{if } x > 1 \end{cases}$

a. Domain: $\{x \mid x > -2\}$ or $(-2, \infty)$

c. Graph:



d. Range: $\{y \mid y > -6\}$ or $(-6, \infty)$

e. There is a jump in the graph at $x = 1$. Therefore, the function is not continuous.


$$\begin{cases} x & \text{if } -4 \leq x < 0 \end{cases}$$

38. $f(x) = \begin{cases} 1 & \text{if } x = 0 \\ 3x & \text{if } x > 0 \end{cases}$

a. Domain: $\{x \mid x \geq -4\}$ or $[-4, \infty)$

b. Intercept: (0, 1)

c. Graph:



d. Range: $\{y \mid y \geq -4, y \neq 0\}$

e. There is a jump at $x = 0$. Therefore, the function is not continuous.

39. $f(x) = \frac{Ax+5}{6x-2}$ and $f(1) = 4$

b. Intercept: (0, 0)

Chapter 2: Functions and Their Graphs

$$\frac{A(1) + 5}{4} = 4$$

Chapter 2 Review Exercises

$$6(1) - 2$$

$$\frac{A + 5}{4} = 4$$

$$A + 5 = 16$$

$$A = 11$$

40. a. $x^2h = 10 \Rightarrow h = \frac{10}{x^2}$

$$A(x) = 2x^2 + 4xh$$



$$= 2x^2 + 4x\left(\frac{10}{x^2}\right)$$

$$= 2x^2 + \frac{40}{x}$$

b. $A(1) = 2 \cdot 1^2 + \frac{40}{1} = 2 + 40 = 42 \text{ ft}^2$


c. $A(2) = 2 \cdot 2^2 + \frac{40}{2} = 8 + 20 = 28 \text{ ft}^2$

d. Graphing:



The area is smallest when $x \approx 2.15$ feet.

41. a. Consider the following diagram:





The area of the rectangle is $A = xy$. Thus,

the area function for the rectangle is:

$$A(x) = x(10 - x^2) = -x^3 + 10x$$

b. The maximum value occurs at the vertex:



Chapter 2 Test

1. a. $\{(2,5), (4,6), (6,7), (8,8)\}$

This relation is a function because there are no ordered pairs that have the same first element and different second elements.

Domain: $\{2, 4, 6, 8\}$

Range: $\{5, 6, 7, 8\}$

b. $\{(1,3), (4,-2), (-3,5), (1,7)\}$

This relation is not a function because there are two ordered pairs that have the same first element but different second elements.

c. This relation is not a function because the graph fails the vertical line test.

d. This relation is a function because it passes the vertical line test.

Domain: $\{x \mid x \text{ is any real number}\}$

Range: $\{y \mid y \geq 2\}$ or $[2, \infty)$

2. $f(x) = \sqrt{4 - 5x}$

The function tells us to take the square root of $4 - 5x$. Only nonnegative numbers have real square roots so we need $4 - 5x \geq 0$.

$$4 - 5x \geq 0$$

$$4 - 5x - 4 \geq 0 - 4$$

$$-5x \geq -4$$

$$\frac{-5x}{-5} \leq \frac{-4}{-5}$$

$$x \leq \frac{4}{5}$$

Domain: $\left\{x \mid x \leq \frac{4}{5}\right\}$ or $\left(-\infty, \frac{4}{5}\right]$

$$f(-1) = \sqrt{4 - 5(-1)} = \sqrt{4 + 5} = \sqrt{9} = 3$$

3. $g(x) = \frac{x+2}{|x+2|}$

The function tells us to divide $x + 2$ by $|x + 2|$.

Chapter 2: Functions and Their Graphs

The maximum area is roughly:

$$A(1.83) = -(1.83)^3 + 10(1.83)$$

$$\approx 12.17 \text{ square units}$$

Chapter 2 Review Exercises

Division by 0 is undefined, so the denominator can never equal 0. This means that $x \neq -2$.

Domain: $\{x \mid x \neq -2\}$

$$g(-1) = \frac{(-1) + 2}{|(-1) + 2|} = \frac{1}{|1|} = 1$$

4. $h(x) = \frac{x-4}{x^2+5x-36}$

The function tells us to divide $x-4$ by $x^2+5x-36$. Since division by 0 is not defined, we need to exclude any values which make the denominator 0.

$$x^2 + 5x - 36 = 0$$

$$(x+9)(x-4) = 0$$

$$x = -9 \text{ or } x = 4$$

$$\text{Domain: } \{x \mid x \neq -9, x \neq 4\}$$

(note: there is a common factor of $x-4$ but we must determine the domain prior to simplifying)

$$h(-1) = \frac{(-1)-4}{(-1)^2+5(-1)-36} = \frac{-5}{-40} = \frac{1}{8}$$

5. a. To find the domain, note that all the points on the graph will have an x -coordinate between -5 and 5 , inclusive. To find the range, note that all the points on the graph will have a y -coordinate between -3 and 3 , inclusive.

$$\text{Domain: } \{x \mid -5 \leq x \leq 5\} \text{ or } [-5, 5]$$

$$\text{Range: } \{y \mid -3 \leq y \leq 3\} \text{ or } [-3, 3]$$

- b. The intercepts are $(0, 2)$, $(-2, 0)$, and $(2, 0)$.

$$x\text{-intercepts: } -2, 2$$

$$y\text{-intercept: } 2$$

- c. $f(1)$ is the value of the function when $x = 1$. According to the graph, $f(1) = 3$.

- d. Since $(-5, -3)$ and $(3, -3)$ are the only points on the graph for which




$$y = f(x) = -3, \text{ we have } f(x) = -3 \text{ when}$$

$$x = -5 \text{ and } x = 3.$$

- e. To solve $f(x) < 0$, we want to find x -

values such that the graph is below the x -axis. The graph is below the x -axis for

Y_{min} and Y_{max} will not be good enough to see the whole picture so some adjustment must be made.



We see that the graph has a local maximum of -0.86 (rounded to two places) when $x = -0.85$ and another local maximum of 15.55 when $x = 2.35$. There is a local minimum of -2 when $x = 0$. Thus, we have

$$\text{Local maxima: } f(-0.85) \approx -0.86$$

$$f(2.35) \approx 15.55$$

$$\text{Local minima: } f(0) = -2$$

The function is increasing on the intervals $(-5, -0.85)$ and $(0, 2.35)$ and decreasing on the intervals $(-0.85, 0)$ and $(2.35, 5)$.


7. a. $f(x) = \begin{cases} 2x+1 & x < -1 \\ x-4 & x \geq -1 \end{cases}$

To graph the function, we graph each "piece". First we graph the line $y = 2x + 1$

but only keep the part for which $x < -1$.

Then we plot the line $y = x - 4$ but only

keep the part for which $x \geq -1$.



Chapter 2: Functions and Their Graphs

values in the domain that are less than -2 and greater than 2 . Therefore, the solution set is $\{x \mid -5 \leq x < -2 \text{ or } 2 < x \leq 5\}$. In interval notation we would write the solution set as $[-5, -2) \cup (2, 5]$.

6. $f(x) = -x^4 + 2x^3 + 4x^2 - 2$

We set $X_{\min} = -5$ and $X_{\max} = 5$. The standard

Chapter 2 Chapter Test

- b. To find the intercepts, notice that the only piece that hits either axis is $y = x - 4$.

$$\begin{aligned} y &= x - 4 & y &= x - 4 \\ y &= 0 - 4 & 0 &= x - 4 \\ y &= -4 & 4 &= x \end{aligned}$$

The intercepts are $(0, -4)$ and $(4, 0)$.

- c. To find $g(-5)$ we first note that $x = -5$ so we must use the first “piece” because $-5 < -1$.

$$g(-5) = 2(-5) + 1 = -10 + 1 = -9$$

- d. To find $g(2)$ we first note that $x = 2$ so we must use the second “piece” because $2 \geq -1$.

$$g(2) = 2 - 4 = -2$$



8. The average rate of change from 3 to 4 is given by

$$\begin{aligned} \frac{\Delta y}{\Delta x} &= \frac{f(4) - f(3)}{4 - 3} \\ &= \frac{3(4)^2 - 2(4) + 4}{4 - 3} - \frac{3(3)^2 - 2(3) + 4}{4 - 3} \\ &= \frac{44 - 25}{4 - 3} = \frac{19}{1} = 19 \end{aligned}$$


9. a. $f - g = (2x^2 + 1) - (3x - 2)$
 $= 2x^2 + 1 - 3x + 2 = 2x^2 - 3x + 3$
 b. $f \cdot g = (2x^2 + 1)(3x - 2) = 6x^3 - 4x^2 + 3x - 2$

c. $f(x+h) - f(x)$
 $= (2(x+h)^2 + 1) - (2x^2 + 1)$
 $= (2(x^2 + 2xh + h^2) + 1) - (2x^2 + 1)$
 $= 2x^2 + 4xh + 2h^2 + 1 - 2x^2 - 1$
 $= 4xh + 2h^2$

10. a. The basic function is $y = x^3$ so we start with the graph of this function.




Next we reflect this graph about the x-axis to obtain the graph of $y = -(x+1)^3$.



Next we stretch this graph vertically by a factor of 2 to obtain the graph of


$$y = -2(x+1)^3$$




$$y = -2(x+1)^3$$

The last step is to shift this graph up 3 units

to obtain the graph of $y = -2(x+1)^3 + 3$.





-10

Next we shift this graph 1 unit to the left to


obtain the graph of $y = (x + 1)^3$.




-10

$$y = -2(x + 1)^3 + 3$$


- b. The basic function is $y = |x|$ so we start with the graph of this function.



Next we shift this graph 4 units to the left to obtain the graph of $y = |x + 4|$.




Next we shift this graph up 2 units to obtain the graph of $y = |x + 4| + 2$.




11. a. $r(x) = -0.115x^2 + 1.183x + 5.623$

For the years 1992 to 2004, we have values of x between 0 and 12. Therefore, we can let $X_{min} = 0$ and $X_{max} = 12$. Since r is the interest rate as a percent, we can try letting $Y_{min} = 0$ and $Y_{max} = 10$.



b. For 2010, we have $x = 2010 - 1992 = 18$.

$$r(18) = -0.115(18)^2 + 1.183(18) + 5.623 = -10.343$$



The model predicts that the interest rate will be -10.343% . This is not a reasonable

value since it implies that the bank would be paying interest to the borrower.

12. a. Let $x =$ width of the rink in feet. Then the length of the rectangular portion is given by $2x - 20$. The radius of the semicircular

portions is half the width, or $r = \frac{x}{2}$.

To find the volume, we first find the area of the surface and multiply by the thickness of the ice. The two semicircles can be combined to form a complete circle, so the area is given by

$$A = l \cdot w + \pi r^2 = (2x - 20)(x) + \pi \left(\frac{x}{2}\right)^2 = 2x^2 - 20x + \frac{\pi x^2}{4}$$


We have expressed our measures in feet so we need to convert the thickness to feet as well.

$$0.75 \text{ in} \cdot \frac{1 \text{ ft}}{12 \text{ in}} = \frac{0.75}{12} \text{ ft} = \frac{1}{16} \text{ ft}$$

Now we multiply this by the area to obtain the volume. That is,

$$V(x) = \frac{1}{16} \left(2x^2 - 20x + \frac{\pi x^2}{4} \right) = \frac{x^2}{8} - \frac{5x}{4} + \frac{\pi x^2}{64}$$

b. If the rink is 90 feet wide, then we have $x = 90$.



The highest rate during this period appears to be 8.67%, occurring in 1997 ($x \approx 5$).

$$V(90) = \frac{90^2}{8} - \frac{5(90)}{4} + \frac{\pi(90)^2}{64} \approx 1297.61$$

The volume of ice is roughly 1297.61 ft³.

Chapter 2 Cumulative Review

1. $3x - 8 = 10$
 $3x - 8 + 8 = 10 + 8$
 $3x = 18$

$$\frac{3x}{3} = \frac{18}{3}$$

$$x = 6$$

The solution set is $\{6\}$.

2. $3x^2 - x = 0$
 $x(3x - 1) = 0$

$$x = 0 \text{ or } 3x - 1 = 0$$

$$3x = 1$$

$$x = \frac{1}{3}$$

The solution set is $\left\{0, \frac{1}{3}\right\}$.

3. $x^2 - 8x - 9 = 0$
 $(x - 9)(x + 1) = 0$
 $x - 9 = 0 \text{ or } x + 1 = 0$
 $x = 9 \quad x = -1$

The solution set is $\{-1, 9\}$.

4. $6x^2 - 5x + 1 = 0$

5. $|2x + 3| = 4$
 $2x + 3 = -4 \text{ or } 2x + 3 = 4$
 $2x = -7 \quad 2x = 1$
 $x = -\frac{7}{2} \quad x = \frac{1}{2}$

The solution set is $\left\{-\frac{7}{2}, \frac{1}{2}\right\}$.

6. $\sqrt{2x + 3} = 2$
 $\left(\sqrt{2x + 3}\right)^2 = 2^2$
 $2x + 3 = 4$
 $2x = 1$
 $x = \frac{1}{2}$

Check:

$$\sqrt{2\left(\frac{1}{2}\right) + 3} = 2$$

$$\sqrt{1 + 3} = 2$$

$$\sqrt{4} = 2$$

$$2 = 2 \text{ T}$$

The solution set is $\left\{\frac{1}{2}\right\}$.

7. $2 - 3x > 6$
 $-3x > 4$
 $x < -\frac{4}{3}$
 $\left\{x \mid x < -\frac{4}{3}\right\}$
 Solution set: $\left\{x \mid x < -\frac{4}{3}\right\}$
 Interval notation: $\left(-\infty, -\frac{4}{3}\right)$

Chapter 2: Functions and Their Graphs

Chapter 2 Chapter Test


$$(3x-1)(2x-1) = 0$$

$$3x-1=0 \quad \text{or} \quad 2x-1=0$$

$$3x=1 \quad 2x=1$$

$$x = \frac{1}{3} \quad x = \frac{1}{2}$$

The solution set is $\left\{\frac{1}{3}, \frac{1}{2}\right\}$.



8. $|2x-5| < 3$

$$-3 < 2x-5 < 3$$

$$2 < 2x < 8$$

$$1 < x < 4$$

Solution set: $\{x | 1 < x < 4\}$

Interval notation: $(1, 4)$



9. $|4x + 1| \geq 7$


$4x + 1 \leq -7$ or $4x + 1 \geq 7$

$4x \leq -8$ $4x \geq 6$

$x \leq -2$ $x \geq \frac{3}{2}$

Solution set: $\left\{ x \mid x \leq -2 \text{ or } x \geq \frac{3}{2} \right\}$

Interval notation: $(-\infty, -2] \cup \left[\frac{3}{2}, \infty \right)$



10. a. $d = \sqrt{\underset{2}{(x-x)}^2 + \underset{1}{(y-y)}^2}$
 $= \sqrt{\underset{2}{(3-(-2))}^2 + \underset{1}{(-5-(-3))}^2}$
 $= \sqrt{(3+2)^2 + (-5+3)^2}$
 $= \sqrt{5^2 + (-2)^2} = \sqrt{25+4}$
 $= \sqrt{29}$

b. $M = \left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2} \right)$
 $= \left(\frac{-2+3}{2}, \frac{-3+(-5)}{2} \right)$
 $= \left(\frac{1}{2}, -4 \right)$

c. $m = \frac{y_2 - y_1}{x_2 - x_1} = \frac{-5 - (-3)}{3 - (-2)} = \frac{-2}{5} = -\frac{2}{5}$


y-intercept:

$3(0) - 2y = 12$

$-2y = 12$


$y = -6$

The point $(0, -6)$ is on the graph.



12. $x = y^2$

y	$x = y^2$	(x, y)
-2	$x = (-2)^2 = 4$	$(4, -2)$
-1	$x = (-1)^2 = 1$	$(1, -1)$
0	$x = 0^2 = 0$	$(0, 0)$
1	$x = 1^2 = 1$	$(1, 1)$
2	$x = 2^2 = 4$	$(4, 2)$



13. $x^2 + (y-3)^2 = 16$

This is the equation of a circle with radius

$\sqrt{\quad}$

Chapter 2: Functions and Their Graphs

11. $3x - 2y = 12$

x -intercept:

$$3x - 2(0) = 12$$

$$3x = 12$$


$$x = 4$$

The point $(4, 0)$ is on the graph.

Chapter 2 Cumulative Review


$r = \sqrt{16} = 4$ and center at $(0, 3)$. Starting at the center we can obtain some points on the graph by moving 4 units up, down, left, and right. The corresponding points are $(0, 7)$, $(0, -1)$,

$(-4, 3)$, and $(4, 3)$, respectively.



14. $y = \sqrt{x}$

x	$y = \sqrt{x}$	(x, y)
0	$y = \sqrt{0} = 0$	$(0, 0)$
1	$y = \sqrt{1} = 1$	$(1, 1)$
4	$y = \sqrt{4} = 2$	$(4, 2)$



15. $3x^2 - 4y = 12$

x -intercepts:

$$3x^2 - 4(0) = 12$$

$$3x^2 = 12$$

$$x^2 = 4$$

$$x = \pm 2$$

y -intercept:

$$3(0)^2 - 4y = 12$$

$$-4y = 12$$

$$y = -3$$

The intercepts are $(-2, 0)$, $(2, 0)$, and $(0, -3)$.

Check x -axis symmetry:

$$3x^2 - 4(-y) = 12$$

$$3x^2 + 4y = 12 \text{ different}$$

Check y -axis symmetry:

$$3(-x)^2 - 4y = 12$$

$$3x^2 - 4y = 12 \text{ same}$$

Check origin symmetry:

$$3(-x)^2 - 4(-y) = 12$$

$$3x^2 + 4y = 12 \text{ different}$$

The graph of the equation has y -axis symmetry.

16. First we find the slope:

$$m = \frac{8 - 4}{6 - (-2)} = \frac{4}{8} = \frac{1}{2}$$

Next we use the slope and the given point $(6, 8)$ in the point-slope form of the equation of a line:

$$y - y_1 = m(x - x_1)$$

$$y - 8 = \frac{1}{2}(x - 6)$$

$$y - 8 = \frac{1}{2}x - 3$$


$$y = \frac{1}{2}x + 5$$

17. $f(x) = (x + 2)^2 - 3$

Starting with the graph of $y = x^2$, shift the graph


2 units to the left [$y = (x + 2)^2$] and down 3

units [$y = (x + 2)^2 - 3$].




18. $f(x) = \frac{1}{x}$

x	$y = \frac{1}{x}$	(x, y)
-1	$y = \frac{1}{-1} = -1$	$(-1, -1)$
1	$y = \frac{1}{1} = 1$	$(1, 1)$
2	$y = \frac{1}{2}$	$(2, \frac{1}{2})$



19. $f(x) = \begin{cases} 2-x & \text{if } x \leq 2 \\ |x| & \text{if } x > 2 \end{cases}$

Graph the line $y = 2 - x$ for $x \leq 2$. Two points on the graph are $(0, 2)$ and $(2, 0)$. Graph the line $y = x$ for $x > 2$. There is a hole in the graph at $x = 2$.



Chapter 2 Projects

Project I – Internet Based Project – Answers will vary

Project II

1. Silver: $C(x) = 20 + 0.16(x - 200) = 0.16x - 12$


$$C(x) = \begin{cases} 20 & 0 \leq x \leq 200 \\ 0.16x - 12 & x > 200 \end{cases}$$

Gold: $C(x) = 50 + 0.08(x - 1000) = 0.08x - 30$

$$C(x) = \begin{cases} 50.00 & 0 \leq x \leq 1000 \\ 0.08x - 30 & x > 1000 \end{cases}$$

Platinum: $C(x) = 100 + 0.04(x - 3000) = 0.04x - 20$

$$C(x) = \begin{cases} 100.00 & 0 \leq x \leq 3000 \\ 0.04x - 20 & x > 3000 \end{cases}$$



3. Let $y = \#K\text{-bytes of service over the plan minimum}$.

Silver: $20 + 0.16y \leq 50$

$$0.16y \leq 30$$

$$y \leq 187.5$$

Silver is the best up to $187.5 + 200 = 387.5$ K-bytes of service.

Gold: $50 + 0.08y \leq 100$

$$0.08y \leq 50$$

$$y \leq 625$$


Gold is the best from 387.5 K-bytes to 625 K-bytes to service.

Platinum: Platinum will be the best if more than 625 K-bytes is needed.


4. Answers will vary.

Project III

1.



2.



$$C(x) = 100x + 140L$$


$$C(x) = 100x + 140\sqrt{4 + (5 - x)^2}$$

3.

x	$C(x)$
0	$100(0) + 140\sqrt{4 + 25} \approx \753.92
1	$100(1) + 140\sqrt{4 + 16} \approx \726.10
2	$100(2) + 140\sqrt{4 + 9} \approx \704.78
3	$100(3) + 140\sqrt{4 + 4} \approx \695.98
4	$100(4) + 140\sqrt{4 + 1} \approx \713.05
5	$100(5) + 140(4 + 0) = \$780.00$

The choice where the cable goes 3 miles down the road then cutting up to the house seems to yield the lowest cost.

4. Since all of the costs are less than \$800, there would be a profit made with any of the plans.



Using the MINIMUM function on a graphing calculator, the minimum occurs at $x \approx 2.96$.

$C(x)$ dollars
800

6. $C(4.5) = 100(4.5) + 140\sqrt{4 + (5 - 4.5)^2}$
 $\approx \$738.62$

The cost for the Steven's cable would be

7. $5000(738.62) = \$3,693,100$ State legislated
 $5000(695.96) = \$3,479,800$ cheapest cost

Project IV

1. $A = \pi r^2$

2. $r = 2.2t$

3. $r = 2.2(2) = 4.4$ ft

$r = 2.2(2.5) = 5.5$ ft

4. $A = \pi(4.4)^2 = 60.82$ ft²

$A = \pi(5.5)^2 = 95.03$ ft²

5. $A = \pi(2.2t)^2 = 4.84\pi t^2$

6. $A = 4.84\pi(2)^2 = 60.82$ ft²

$A = 4.84\pi(2.5)^2 = 95.03$ ft²

7. $\frac{A(2.5) - A(2)}{2.5 - 2} = \frac{95.03 - 60.82}{0.5} = 68.42$ ft/hr

8. $\frac{A(3.5) - A(3)}{3.5 - 3} = \frac{186.27 - 136.85}{0.5} = 98.84$ ft/hr

9. The average rate of change is increasing.

10. 150 yds = 450 ft


$r = 2.2t$

$t = \frac{450}{2.2} = 204.5$ hours

11. 6 miles = 31680 ft

Therefore, we need a radius of 15,840 ft.

$t = \frac{15,840}{2.2} = 7200$ hours



The minimum cost occurs when the cable runs for 2.96 mile along the road.