Test Bank for Principles and Practice of Physics 1st Edition Eric Mazur 032194920X 9780321949202

Full link download:

Test Bank:

https://testbankpack.com/p/test-bank-for-principles-

and-practice-of-physics-1st-edition-eric-mazur-

032194920x-9780321949202/


Solution Manual:

https://testbankpack.com/p/solution-manual-for-principles-and-practice-of-physics-1st-edition-eric-mazur-032194920x-9780321949202/

Principles & Practice of Physics (Mazur) Chapter 2 Motion in One Dimension

2.1 Conceptual Questions


1) The motion of a particle is described in the velocity versus time graph shown in the figure. We can say that its speed



- A) increases.
- B) decreases.
- C) increases and then decreases.
- D) decreases and then increases.

Answer: D Var: 1

2) The motions of a car and a truck along a straight road are represented by the velocity-time graphs in the figure. The two vehicles are initially alongside each other at time t = 0. At time T, what is true about these two vehicles since time t = 0?




- A) The truck will have traveled further than the car.
- B) The car will have traveled further than the truck.
- C) The truck and the car will have traveled the same distance.
- D) The car will be traveling faster than the truck.

Answer: A


3) The graph in the figure shows the position of an object as a function of time. The letters H-L represent particular moments of time. At which moments shown (H, I, etc.) is the speed of the object


- (a) the greatest?
- (b) the smallest?




Answer: (a) J (b) I


4) The figure shows the graph of the position *x* as a function of time for an object moving in the straight line (the *x*-axis). Which of the following graphs best describes the velocity along the *x*-axis as a function of time for this object?










5) An object starts its motion with a constant velocity of 2.0 m/s toward the east. After 3.0 s, the object stops for 1.0 s. The object then moves toward the west a distance of 2.0 m in 3.0 s. The object continues traveling in the same direction, but increases its speed by 1.0 m/s for the next 2.0 s. Which graph below could represent the motion of this object?









Answer: D Var: 1


2.2 Problems

- 1) A cat runs along a straight line (the x-axis) from point A to point B to point C, as shown in the figure. The distance between points A and C is 5.00 m, the distance between points B and C is 10.0 m, and the positive direction of the x-axis points to the right. The time to run from A to B is 20.0 s, and the time from B to C is 8.00 s. As the cat runs along the x-axis between points A and C
- (a) what is the magnitude of its average velocity?
- (b) what is its average speed?



Answer: (a) 0.179 m/s (b) 0.893 m/s

- 2) The figure shows the position of an object as a function of time. During the time interval from time t = 0.0 s and time t = 9.0 s
- (a) what is the length of the path the object followed?
- (b) what is the displacement of the object?




Answer: (a) 5.0 m (b) 1.0 m

Var: 1

- 3) As part of an exercise program, a woman walks south at a speed of 2.00 m/s for 60.0 minutes. She then turns around and walks north a distance 3000 m in 25.0 minutes
- (a) What is the woman's average velocity during her entire motion?
 - A) 0.824 m/s south
 - B) 1.93 m/s south
 - C) 2.00 m/s south
 - D) 1.79 m/s south
 - E) 800 m/s south
- (b) What is the woman's average speed during her entire motion?
 - A) 0.824 m/s
 - B) 1.93 m/s
 - C) 2.00 m/s
 - D) 1.79 m/s
 - E) 800 m/s
- Answer: (a) A (b) C

Var: 1

- 4) The figure shows the position of an object as a function of time, with all numbers accurate to two significant figures. Between time t = 0.0 s and time t = 9.0 s
- (a) what is the average speed of the object?
- (b) what is the average velocity of the object?



Answer: (a) 0.56 m/s (b) 0.11 m/s


Var: 1

- 5) If the fastest you can safely drive is 65 mi/h, what is the longest time you can stop for dinner if you must travel 541 mi in 9.6 h total?
- A) 1.0 h
- B) 1.3 h
- C) 1.4 h
- D) You can't stop at all.

Answer: B

- Var: 50+
- 6) Arthur and Betty start walking toward each other when they are 100 m apart. Arthur has a speed of 3.0 m/s and Betty has a speed of 2.0 m/s. Their dog, Spot, starts by Arthur's side at the same time and runs back and forth between them at 5.0 m/s. By the time Arthur and Betty meet, what distance has Spot run? Answer: 100 m

7) The figure represents the position of a particle as it travels along the x-axis. Between t = 2 s and t = 4 s, what is (a) the average speed of the particle and (b) the average velocity of the particle?




Answer: (a) 1.0 m/s (b) 0.00 m/s

Var: 1

8) The figure shows a graph of the velocity as a function of time for a basketball player traveling up and down the court in a straight-line path. For the 10 s shown on the graph, find

(a) the net displacement of the player.

(b) the total distance run by the player.



Answer: (a) 18 m (b) 20 m

Var: 1

9) The position of an object as a function of time is given by $x = bt^2 - ct$, where $b = 2.0 \text{ m/s}^2$ and c = 6.7 m/s and t are in SI units. What is the instantaneous velocity of the object when t = 2.2?

A) 1.7 m/s

B) 2.1 m/s

C) 2.3 m/s

D) 2.7 m/s

Answer: B

- 10) The velocity of an object is given by the expression $v(t) = 3.00 \text{ m/s} + (4.00 \text{ m/s}^3)t^2$, where t is in seconds. Determine the position of the object as a function of time if it is located at x = 1.00 m at time t = 0.000 s.
- A) (4.00 m/s)t + 1.00 m
- B) $(3.00 \text{ m/s})t + (1.33 \text{ m/s}3)t^3$
- C) (4.00 m/s)t
- D) 1.33 m
- E) $1.00 \text{ m} + (3.00 \text{ m/s})t + (1.33 \text{ m/s}^3)t^3$

Answer: E Var: 5