Test Bank for Applied Statistics in Business and Economics 4th Edition Doane Seward 0073521485

9780073521480

Full Link

Download: Test

Bank:

https://testbankpack.com/p/test-bank-for-applied-statistics-in-businessand-economics-4th-edition-doane-seward-0073521485-9780073521480/

Solution Manual:

https://testbankpack.com/p/solution-manual-for-applied-statistics-in-businessand-economics-4th-edition-doane-seward-0073521485-9780073521480/

Chapter 02

Data Collection

True / False Questions

1. Categorical data have values that are described by words rather than numbers.

True False

2. Numerical data can be either discrete or continuous.

True False

3. Categorical data are also referred to as nominal or qualitative data.

True False

4. The number of checks processed at a bank in a day is an example of categorical data.

5. The number of planes per day that land at an airport is an example of discrete data.

True False

6. The weight of a bag of dog food is an example of discrete data.

True False

7. In last year's annual report, Thompson Distributors indicated that it had 12 regional warehouses. This is an example of ordinal level data.

True False

8. Nominal data refer to data that can be ordered in a natural way.

True False

9. This year, Oxnard University produced two football All-Americans. This is an example of continuous data.

True False

10. The type of statistical test that we can perform is independent of the level of measurement of the variable of interest.

11. Your weight recorded at your annual physical would not be ratio data because you cannot have zero weight.

True False

12. The level of measurement for categorical data is nominal.

True False

13. Temperature measured in degrees Fahrenheit is an example of interval data.

True False

14. The closing price of a stock is an example of ratio data.

True False

15. The Statistical Abstract of the United States is a huge annual compendium of data for the United States and it is available online free of charge.

True False

16. Ordinal data can be treated as if it were nominal data but not vice versa.

17. Responses on a seven-point Likert scale are usually treated as ratio data.

True False

18. Likert scales are especially important in opinion polls and marketing surveys.

True False

19. Ordinal data are data that can be ranked based on some natural characteristic of the items.

True False

20. Ratio data are distinguished from interval data by the presence of a zero reference point.

True False

21. It is better to attempt a census of a large population instead of relying on a sample.

True False

22. Judgment sampling and convenience sampling are non-random sampling techniques.

23. A problem with judgment sampling is that the sample may not reflect the population.

True False

24. When the population is large, a sample estimate is usually preferable to a census.

True False

25. Sampling error is avoidable by choosing the sample scientifically.

True False

26. A sampling frame is used to identify the target population in a statistical study.

True False

27. By taking a systematic sample, in which we select every 50th shopper arriving at a specific store, we are approximating a random sample of shoppers.

True False

28. A worker collecting data from every other shopper who leaves a store is taking a simple random sample of customer opinion.

29. Creating a list of people by taking the third name listed on every 10th page of the phone book is an example of convenience sampling.

True False

30. Internet surveys posted on popular websites have no bias since anyone can reply.

True False

31. Analysis of month-by-month changes in stock market prices during the most recent recession would require the use of time series data.

True False

32. A cluster sample is a type of stratified sample that is based on geographical location.

True False

33. An advantage of a systematic sample is that no list of enumerated data items is required.

34. Telephone surveys often have a low response rate and fail to reach the desired population.

True False

35. Mail surveys are attractive because of their high response rates.

True False

36. A problem with convenience sampling is that the target population is not well defined.

True False

37. If you randomly sample 50 students about their favorite places to eat, the data collected would be referred to as cross-sectional data.

True False

38. The number of FedEx shipping centers in each of 50 cities would be ordinal level data.

39. Internet surveys posted on popular websites such as MSN.com suffer from nonresponse bias.

True False

Multiple Choice Questions

- 40. An investment firm rates bonds for AardCo Inc. as "B+" while bonds of Deva Corp. are rated "AA." Which level of measurement would be appropriate for such data?
 - A. Nominal
 - B. Ordinal
 - C. Interval
 - D. Ratio
- 41. Which variable is least likely to be regarded as ratio data?
 - A. Length of time required for a randomly chosen vehicle to cross a toll bridge (minutes)
 - B. Weight of a randomly chosen student (pounds)
 - C. Number of fatalities in a randomly chosen traffic disaster (persons)
 - D. Student's evaluation of a professor's teaching (Likert scale)

- 42. Which of the following is numerical data?
 - A. Your gender
 - B. The brand of cell phone you own
 - C. Whether you have an American Express card
 - D. The fuel economy (MPG) of your car
- 43. Measurements from a sample are called:
 - A. statistics.
 - B. inferences.
 - C. parameters.
 - D. variables.
- 44. Quantitative variables use which two levels of measurement?
 - A. Ordinal and ratio
 - B. Interval and ordinal
 - C. Nominal and ordinal
 - D. Interval and ratio

45. Temperature in degrees Fahrenheit is an example of a(n) ______ variable.

- A. nominal
- B. ordinal
- C. interval
- D. ratio

46. Using a sample to make generalizations about an aspect of a population is called:

- A. data mining.
- B. descriptive statistics.
- C. random sampling.
- D. statistical inference.

47. Your telephone area code is an example of a(n) ______ variable.

- A. nominal
- B. ordinal
- C. interval
- D. ratio

48. Which is least likely to be regarded as a ratio variable?

- A. A critic's rating of a restaurant on a 1 to 4 scale
- B. Automobile exhaust emission of nitrogen dioxide (milligrams per mile)
- C. Number of customer complaints per day at a cable TV company office
- D. Cost of an e-Bay purchase

49. Automobile exhaust emission of CO₂ (milligrams per mile) is ______ data.

- A. nominal
- B. ordinal
- C. interval
- D. ratio
- 50. Your rating of the food served at a local restaurant using a three-point scale of 0
 - = gross, 1 = decent, 2 = yummy is _____ data.
 - A. nominal
 - B. ordinal
 - C. interval
 - D. ratio

- 51. The number of passengers "bumped" on a particular airline flight is ______ data.
 - A. nominal
 - B. ordinal
 - C. interval
 - D. ratio
- 52. Which should not be regarded as a continuous random variable?
 - A. Tonnage carried by a randomly chosen oil tanker at sea
 - B. Wind velocity at 7 o'clock this morning
 - C. Number of personal fouls by the Miami Heat in a game
 - D. Length of time to play a Wimbledon tennis match
- 53. Which of the following is not true?
 - A. Categorical data have values that are described by words rather than numbers.
 - B. Categorical data are also referred to as nominal or qualitative data.
 - C. The number of checks processed at a bank in a day is categorical data.
 - D. Numerical data can be either discrete or continuous.

- 54. Which of the following is true?
 - A. The type of charge card used by a customer (Visa, MasterCard, AmEx) is ordinal data.
 - B. The duration (minutes) of a flight from Boston to Minneapolis is ratio data.
 - C. The number of Nobel Prize-winning faculty at Oxnard University is continuous data.
 - D. The number of regional warehouses owned by Jankord Industries is ordinal data.
- 55. Which statement is correct?
 - A. Judgment sampling is preferred to systematic sampling.
 - B. Sampling without replacement introduces bias in our estimates of parameters.
 - C. Cluster sampling is useful when strata characteristics are unknown.
 - D. Focus groups usually work best without a moderator.

56. A Likert scale:

- A. yields interval data if scale distances are equal.
- B. must have an odd number of scale points.
- C. must have a verbal label on each scale point.
- D. is rarely used in marketing surveys.

57. Which is most nearly correct regarding sampling error?

A. It can be eliminated by increasing the sample size.

- B. It cannot be eliminated by any statistical sampling method.
- C. It can be eliminated by using Excel's =RANDBETWEEN() function.
- D. It can be eliminated by utilizing systematic random sampling.

58. Which statement is false?

- A. Random dialing phone surveys have low response and are poorly targeted.
- B. Selection bias means that many respondents dislike the interviewer.
- C. Simple random sampling requires a list of the population.
- D. Web surveys are economical but suffer from nonresponse bias.
- 59. Judgment sampling is sometimes preferred over random sampling, for example, when:
 - A. the desired sample size is much larger than the population.
 - B. the sampling budget is large and the population is conveniently located.
 - C. time is short and the sampling budget is limited.
 - D. the population is readily accessible and sampling is nondestructive.

60. An advantage of convenience samples is that:

- A. the required sample size is easier to calculate.
- B. sampling error can be reduced.
- C. computation of statistics is easier.
- D. they are often quicker and cheaper.
- 61. Before deciding whether to assess heavy fines against noisy airlines, which sampling method would the Federal Aviation Administration probably use to measure the peak noise from departing jets as measured by a ground-level observer at a point one mile from the end of the departure runway?
 - A. Radio survey of pilots.
 - B. Simple random sample.
 - C. Judgment sample.
 - D. Stratified sample using.

- 62. Professor Hardtack chose a sample of 7 students from his statistics class of 35 students by picking every student who was wearing red that day. Which kind of sample is this?
 - A. Simple random sample
 - B. Judgment sample
 - C. Systematic sample
 - D. Convenience sample
- 63. Thirty work orders are selected from a filing cabinet containing 500 work order folders by choosing every 15th folder. Which sampling method is this?
 - A. Simple random sample
 - B. Systematic sample
 - C. Stratified sample
 - D. Cluster sample
- 64. Which of the following is not a likely reason for sampling?
 - A. The destructive nature of certain tests
 - B. The physical impossibility of checking all the items in the population
 - C. Prohibitive cost of studying the entire population
 - D. The expense of obtaining random numbers

- 65. Comparing a census of a large population to a sample drawn from it, we expect that the:
 - A. sample is usually a more practical method of obtaining the desired information.
 - B. accuracy of the observations in the census is surely higher than in the sample.
 - C. sample must be a large fraction of the population to be accurate.

66. A stratified sample is sometimes recommended when:

- A. the sample size is very large.
- B. the population is small compared to the sample.
- C. distinguishable strata can be identified in the populations.
- D. the population is spread out geographically.
- 67. A random sample is one in which the:
 - A. probability that an item is selected for the sample is the same for all population items.
 - B. population items are selected haphazardly by experienced workers.
 - C. items to be selected from the population are specified based on expert judgment.
 - D. probability of selecting a population item depends on the item's data value.

68. An advantage of convenience samples over random samples is that:

A. they are easy to analyze.

- B. it is easier to determine the sample size needed.
- C. it is easier to calculate the sampling errors involved.
- D. data collection cost is reduced.
- 69. To measure satisfaction with its cell phone service, AT&T takes a stratified sample of its customers by age, gender, and location. Which is an advantage of this type of sampling, as opposed to other sampling methods?
 - A. It is less intrusive on customers' privacy.
 - B. It does not require random numbers.
 - C. It gives faster results.
 - D. It can give more accurate results.
- 70. An accounting professor wishing to know how many MBA students would take a summer elective in international accounting did a survey of the class she was teaching. Which kind of sample is this?
 - A. Simple random sample
 - B. Cluster sample
 - C. Systematic sample
 - D. Convenience sample

- 71. A binary variable (also called a dichotomous variable or dummy variable) has:
 - A. only two possible values.
 - B. continuous scale values.
 - C. rounded data values.
 - D. ordinal or interval values.
- 72. A population has groups that have a small amount of variation within them, but large variation among or between the groups themselves. The proper sampling technique is:
 - A. simple random.
 - B. stratified.
 - C. cluster.
 - D. judgment.
- 73. A manager chose two people from his team of eight to give an oral presentation because she felt they were representative of the whole team's views. What sampling technique did she use in choosing these two people?
 - A. Convenience
 - B. Simple random
 - C. Judgment
 - D. Cluster

- 74. Sampling bias can best be reduced by:
 - A. using appropriate data coding.
 - B. having a computer tabulate the results.
 - C. utilizing random sampling.
 - D. taking a judgment sample.
- 75. A sampling technique used when groups are defined by their geographical location is:
 - A. cluster sampling.
 - B. convenience sampling.
 - C. judgment sampling.
 - D. random sampling.
- 76. If we choose 500 random numbers using Excel's function =RANDBETWEEN(1,99), we would most likely find that:
 - A. numbers near the mean (50) would tend to occur more frequently.
 - B. numbers near 1 and 99 would tend to occur less frequently.
 - C. some numbers would occur more than once.
 - D. the numbers would have a clear pattern.

- 77. A problem with nonrandom sampling is that:
 - A. larger samples need to be taken to reduce the sampling error inherent in this approach.
 - B. not every item in the population has the same chance of being selected, as it should.
 - C. it is usually more expensive than random sampling.
 - D. it generally provides lower response rates than random sampling.
- 78. From its 32 regions, the FAA selects 6 regions, and then randomly audits 25 departing commercial flights in each region for compliance with legal fuel and weight requirements. This is an example of:
 - A. simple random sampling.
 - B. stratified random sampling.
 - C. cluster sampling.
 - D. judgment sampling.

- 79. Which of the following is a correct statement?
 - A. Choosing the third person listed on every fifth page of the phone book is stratified sampling.
 - B. An advantage of a systematic sample is that no list of enumerated data items is required.
 - C. Convenience sampling is used to study shoppers in convenience stores.
 - D. Judgment sampling is an example of true random sampling.
- 80. Which of the following is false?
 - A. Sampling error is the difference between the true parameter and a random number in Excel.
 - B. Sampling error is a result of unavoidable random variation in a sample.
 - C. A sampling frame is used to help identify the target population in a statistical study.
 - D. The target population must first be defined by a full list or data file of individuals.

- 81. When we are choosing a random sample and we do not place chosen units back into the population, we are:
 - A. sampling with replacement.
 - B. sampling without replacement.
 - C. using a systematic sample.
 - D. using a voluntary sample.
- 82. Which method is likely to be used by a journalism student who is casually surveying opinions of students about the university's cafeteria food for an article that she is writing?
 - A. Simple random sample
 - B. Systematic random sample
 - C. Cluster sample
 - D. Convenience sample
- 83. Which of the following is false?
 - A. Mail surveys are cheap but have low response rates.
 - B. Coverage error is when respondents give untruthful answers.
 - C. Focus groups are nonrandom but can probe issues more deeply.
 - D. Surveys posted on popular websites suffer from selection bias.

84. Which is a time series variable?

A. VISA balances of 30 students on December 31 of this year

- B. Net earnings reported by Xena Corp. for the last 10 quarters
- C. Dollar exchange rates yesterday against 10 other world currencies
- D. Titles of the top 10 movies in total revenue last week

85. An observation in a data set would refer to:

A. only a variable whose value is recorded by visual inspection.

- B. a data item whose value is numerical (as opposed to categorical).
- C. a single row that contains one or more observed variables.
- D. the values of all the variables in the entire data set.

86. A multivariate data set contains:

- A. more than two observations.
- B. more than two categorical variables.
- C. more than two variables.
- D. more than two levels of measurement.

- 87. The Centers for Disease Control and Prevention (CDC) wants to estimate the average extra hospital stay that occurs when heart surgery patients experience postoperative atrial fibrillation. They divide the United States into nine regions. In each region, hospitals are selected at random within each hospital size group (small, medium, large). In each hospital, heart surgery patients are sampled according to known percentages by age group (under 50, 50 to 64, 65 and over) and gender (male, female). This procedure combines which sampling methods?
 - A. Systematic, simple random, and convenience
 - B. Convenience, systematic, and judgment
 - C. Cluster, stratified, and simple random
 - D. Judgment, systematic, and simple random
- 88. Which statement is correct?
 - A. Selecting every fifth shopper arriving at a store will approximate a random sample of shoppers.
 - B. Selecting only shoppers who drive SUVs is a stratified sampling method.
 - C. A census is preferable to a sample for most business problems.
 - D. Stratified samples are usually cheaper than other methods.

89. Which is a categorical variable?

- A. The brand of jeans you usually wear
- B. The price you paid for your last pair of jeans
- C. The distance to the store where you purchased your last pair of jeans
- D. The number of pairs of jeans that you own

90. Which is a discrete variable?

- A. The time it takes to put on a pair of jeans
- B. The price you paid for your last pair of jeans
- C. The distance to the store where you purchased your last pair of jeans
- D. The number of pairs of jeans that you own
- 91. A section of the population we have targeted for analysis is:
 - A. a statistic.
 - B. a frame.
 - C. a sample.
 - D. a coven.

Short Answer Questions

92. Which survey method would you recommend to survey opinions of airline passengers about the cleanliness of the restrooms in the Detroit airport? Why not the others?

93. What kind of sampling method would you suggest in order to tabulate the number of formulas on a typical page of the Doane-Seward textbook? Defend your choice.

94. How would you design a study to see whether drivers using hands-free cell phones are distracted enough to slow their reactions to emergency situations? How would you collect data?

95. Explain the concept of a focus group. In what ways does a focus group resemble a survey? Why is a moderator desirable? What else is required to make a successful focus group?

Chapter 02 Data Collection Answer Key

True / False Questions

1. Categorical data have values that are described by words rather than numbers.

TRUE

Categories are nominal data but could also be ranked (e.g., sophomore, junior, senior).

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

2. Numerical data can be either discrete or continuous.

TRUE

Numerical data can be counts (e.g., cars owned) or continuous scales (e.g., height).

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

3. Categorical data are also referred to as nominal or qualitative data.

TRUE

Categories are nominal data (nonnumerical), sometimes called qualitative data.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

4. The number of checks processed at a bank in a day is an example of categorical data.

FALSE

Integers are actually numerical data.

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions 5. The number of planes per day that land at an airport is an example of discrete data.

TRUE

Integers are discrete numerical data.

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

6. The weight of a bag of dog food is an example of discrete data.

FALSE

Weight is measured on a continuous scale.

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions 7. In last year's annual report, Thompson Distributors indicated that it had 12 regional warehouses. This is an example of ordinal level data.

FALSE

"Number of" is a count, which is ratio data because a zero exists (better than ordinal).

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

8. Nominal data refer to data that can be ordered in a natural way.

FALSE

Nominal (categorical) data would be called ordinal only if categories can be ranked.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement 9. This year, Oxnard University produced two football All-Americans. This is an example of continuous data.

FALSE

The "number of" anything is discrete.

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

10. The type of statistical test that we can perform is independent of the level of measurement of the variable of interest.

FALSE

Some statistical operations are restricted unless you have ratio or interval rata.

AACSB: Analytic Blooms: Understand Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement 11. Your weight recorded at your annual physical would not be ratio data because you cannot have zero weight.

FALSE

Zero is only a reference point, not necessarily an observable data value.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

12. The level of measurement for categorical data is nominal.

TRUE

Categorical and nominal are equivalent terms.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement 13. Temperature measured in degrees Fahrenheit is an example of interval data.

TRUE

For temperature, scale distances are meaningful (20 to 25 is the same as 50 to 55 degrees) and 0 degrees Fahrenheit does not mean the absence of heat so it is not a ratio measurement.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

14. The closing price of a stock is an example of ratio data.

TRUE

True zero exists as a reference, whether or not it is observed.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement
15. The Statistical Abstract of the United States is a huge annual compendium of data for the United States and it is available online free of charge.

TRUE

A useful reference for business (e.g., for marketing, economics, or finance).

AACSB: Technology Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-08 Find everyday print or electronic data sources. Topic: Data Sources

16. Ordinal data can be treated as if it were nominal data but not vice versa.

TRUE

You can always go back to a lower level of measurement (but not vice versa).

AACSB: Analytic Blooms: Understand Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement 17. Responses on a seven-point Likert scale are usually treated as ratio data.

FALSE

No true zero point exists.

AACSB: Analytic Blooms: Understand Difficulty: 1 Easy Learning Objective: 02-05 Recognize a Likert scale and know how to use it. Topic: Level of Measurement

18. Likert scales are especially important in opinion polls and marketing surveys.

TRUE

Likert scales are used in all business areas.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-05 Recognize a Likert scale and know how to use it. Topic: Level of Measurement 19. Ordinal data are data that can be ranked based on some natural characteristic of the items.

TRUE

For example, the eras Jurassic, Paleozoic, and Mesozoic can be ranked in time.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

20. Ratio data are distinguished from interval data by the presence of a zero reference point.

TRUE

The true zero is a reference that need not be observable.

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement 21. It is better to attempt a census of a large population instead of relying on a sample.

FALSE

A census may founder on cost and time, while samples can be quick and accurate.

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts

22. Judgment sampling and convenience sampling are non-random sampling techniques.

TRUE

To be random, every item must have the same chance of being chosen.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods 23. A problem with judgment sampling is that the sample may not reflect the population.

TRUE

While better than mere convenience, judgment may still have flaws.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

24. When the population is large, a sample estimate is usually preferable to a census.

TRUE

A census may founder on cost and time, while samples can be quick and accurate.

AACSB: Analytic Blooms: Understand Difficulty: 1 Easy Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts 25. Sampling error is avoidable by choosing the sample scientifically.

FALSE

Sampling error is unavoidable, though it can be reduced by careful sampling.

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

26. A sampling frame is used to identify the target population in a statistical study.

TRUE

Only some portion of the population may be targeted (e.g., independent voters).

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts 27. By taking a systematic sample, in which we select every 50th shopper arriving at a specific store, we are approximating a random sample of shoppers.

TRUE

There is no bias if this method is implemented correctly.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

28. A worker collecting data from every other shopper who leaves a store is taking a simple random sample of customer opinion.

FALSE

Not unless the target population is customers who shopped today (cf., all customers).

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods 29. Creating a list of people by taking the third name listed on every 10th page of the phone book is an example of convenience sampling.

FALSE

This resembles two-stage cluster sampling combined with systematic sampling.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

 Internet surveys posted on popular websites have no bias since anyone can reply.

FALSE

Self-selection bias exists (respondents may be atypical).

AACSB: Technology Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-09 Describe basic elements of survey design; survey types; and sources of error. Topic: Surveys 31. Analysis of month-by-month changes in stock market prices during the most recent recession would require the use of time series data.

TRUE

Data collected over time is a time series.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-03 Explain the difference between time series and cross-sectional data. Topic: Definitions

32. A cluster sample is a type of stratified sample that is based on geographical location.

TRUE

For example, sampling voters randomly within random zip codes.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods 33. An advantage of a systematic sample is that no list of enumerated data items is required.

TRUE

Systematic sampling works with a list (like random sampling) but also without one.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

34. Telephone surveys often have a low response rate and fail to reach the desired population.

TRUE

Phone surveys are cheaper, but it is hard to avoid these problems.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-09 Describe basic elements of survey design; survey types; and sources of error. Topic: Surveys 35. Mail surveys are attractive because of their high response rates.

FALSE

Mail surveys have low response rates and invite self-selection bias.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-09 Describe basic elements of survey design; survey types; and sources of error. Topic: Surveys

36. A problem with convenience sampling is that the target population is not well defined.

TRUE

Convenience sampling is quick but not random and target population is unclear.

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods 37. If you randomly sample 50 students about their favorite places to eat, the data collected would be referred to as cross-sectional data.

TRUE

Data for individuals would be a cross section (not a time series).

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-03 Explain the difference between time series and cross-sectional data. Topic: Definitions

 The number of FedEx shipping centers in each of 50 cities would be ordinal level data.

FALSE

The "number of" anything is ratio data because a true zero reference point exists.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement 39. Internet surveys posted on popular websites such as MSN.com suffer from nonresponse bias.

TRUE

Nonresponse or self-selection bias is rampant in such surveys.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-09 Describe basic elements of survey design; survey types; and sources of error. Topic: Surveys

Multiple Choice Questions

- 40. An investment firm rates bonds for AardCo Inc. as "B+" while bonds of Deva Corp. are rated "AA." Which level of measurement would be appropriate for such data?
 - A. Nominal
 - <u>B.</u> Ordinal
 - C. Interval
 - D. Ratio

Ranks are clear, but interval would require assumed equal scale distances (doubtful).

AACSB: Analytic Blooms: Evaluate Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

- 41. Which variable is least likely to be regarded as ratio data?
 - A. Length of time required for a randomly chosen vehicle to cross a toll bridge (minutes)
 - B. Weight of a randomly chosen student (pounds)
 - C. Number of fatalities in a randomly chosen traffic disaster (persons)
 - D. Student's evaluation of a professor's teaching (Likert scale)

Likert scales have no true zero.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

- 42. Which of the following is numerical data?
 - A. Your gender
 - B. The brand of cell phone you own
 - C. Whether you have an American Express card
 - D. The fuel economy (MPG) of your car

Fuel economy is numerical.

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

- 43. Measurements from a sample are called:
 - <u>A.</u> statistics.
 - B. inferences.
 - C. parameters.
 - D. variables.

A measurement calculated from a sample is a statistic.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Methods

44. Quantitative variables use which two levels of measurement?

- A. Ordinal and ratio
- B. Interval and ordinal
- C. Nominal and ordinal
- D. Interval and ratio

Numerical (quantitative) data can be interval or ratio.

Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

45. Temperature in degrees Fahrenheit is an example of a(n) ______ variable.

- A. nominal
- B. ordinal
- <u>C.</u> interval
- D. ratio

No true zero exists except on the Kelvin scale.

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

- 46. Using a sample to make generalizations about an aspect of a population is called:
 - A. data mining.
 - B. descriptive statistics.
 - C. random sampling.
 - D. statistical inference.

Generalizing from a sample to a population is an inference.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts

47. Your telephone area code is an example of a(n) variable.

<u>A.</u> nominal

- B. ordinal
- C. interval
- D. ratio

Area codes are not even ranked, so just nominal.

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

- 48. Which is least likely to be regarded as a ratio variable?
 - A. A critic's rating of a restaurant on a 1 to 4 scale
 - B. Automobile exhaust emission of nitrogen dioxide (milligrams per mile)
 - C. Number of customer complaints per day at a cable TV company office
 - D. Cost of an e-Bay purchase

Ratings on a Likert scale have no meaningful zero.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

49. Automobile exhaust emission of CO₂ (milligrams per mile) is ______ data.

- A. nominal
- B. ordinal
- C. interval
- <u>D.</u> ratio

True zero exists.

Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

50. Your rating of the food served at a local restaurant using a three-point scale of

0 = gross, 1 = decent, 2 = yummy is _____ data.

A. nominal

<u>B.</u> ordinal

C. interval

D. ratio

Only rankings implied (not equal scale distances).

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

- 51. The number of passengers "bumped" on a particular airline flight is ______ data.
 - A. nominal
 - B. ordinal
 - C. interval
 - <u>D.</u> ratio

True zero exists.

AACSB: Analytic Blooms: Apply Difficulty: 1 Easy Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

- 52. Which should not be regarded as a continuous random variable?
 - A. Tonnage carried by a randomly chosen oil tanker at sea
 - B. Wind velocity at 7 o'clock this morning
 - C. Number of personal fouls by the Miami Heat in a game
 - D. Length of time to play a Wimbledon tennis match

Counting things yields integer (discrete) data.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

- 53. Which of the following is not true?
 - A. Categorical data have values that are described by words rather than numbers.
 - B. Categorical data are also referred to as nominal or qualitative data.
 - <u>C.</u> The number of checks processed at a bank in a day is categorical data.
 - D. Numerical data can be either discrete or continuous.

The "number of" anything is a discrete numerical variable.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

- 54. Which of the following is true?
 - A. The type of charge card used by a customer (Visa, MasterCard, AmEx) is ordinal data.
 - B. The duration (minutes) of a flight from Boston to Minneapolis is ratio data.
 - C. The number of Nobel Prize-winning faculty at Oxnard University is continuous data.
 - D. The number of regional warehouses owned by Jankord Industries is ordinal data.

True zero exists (not observable, but as a reference point) so ratios have meaning.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-04 Recognize levels of measurement in data and ways of coding data. Topic: Level of Measurement

- 55. Which statement is correct?
 - A. Judgment sampling is preferred to systematic sampling.
 - B. Sampling without replacement introduces bias in our estimates of parameters.
 - C. Cluster sampling is useful when strata characteristics are unknown.
 - D. Focus groups usually work best without a moderator.

Review the characteristics of each sampling method.

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

56. A Likert scale:

A. yields interval data if scale distances are equal.

- B. must have an odd number of scale points.
- C. must have a verbal label on each scale point.
- D. is rarely used in marketing surveys.

Marketers use Likert scales and try to make scales with meaningful intervals.

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium

- 57. Which is most nearly correct regarding sampling error?
 - A. It can be eliminated by increasing the sample size.
 - <u>B.</u> It cannot be eliminated by any statistical sampling method.
 - C. It can be eliminated by using Excel's =RANDBETWEEN() function.
 - D. It can be eliminated by utilizing systematic random sampling.

Sampling involves error, though it can be minimized by proper methodology.

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts

58. Which statement is false?

- A. Random dialing phone surveys have low response and are poorly targeted.
- B. Selection bias means that many respondents dislike the interviewer.
- C. Simple random sampling requires a list of the population.
- D. Web surveys are economical but suffer from nonresponse bias.

Selection bias occurs when respondents are atypical.

- 59. Judgment sampling is sometimes preferred over random sampling, for example, when:
 - A. the desired sample size is much larger than the population.
 - B. the sampling budget is large and the population is conveniently located.
 - C. time is short and the sampling budget is limited.
 - D. the population is readily accessible and sampling is nondestructive.

Judgment sampling can save time and may be better than mere convenience.

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 60. An advantage of convenience samples is that:
 - A. the required sample size is easier to calculate.
 - B. sampling error can be reduced.
 - C. computation of statistics is easier.
 - D. they are often quicker and cheaper.

Convenience samples are quick, with a possible trade-off of accuracy.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 61. Before deciding whether to assess heavy fines against noisy airlines, which sampling method would the Federal Aviation Administration probably use to measure the peak noise from departing jets as measured by a ground-level observer at a point one mile from the end of the departure runway?
 - A. Radio survey of pilots.
 - B. Simple random sample.
 - C. Judgment sample.
 - D. Stratified sample using.

Pilots can't assess noise (the measurements must be taken from the ground). No list is available for the unpredictable mix of departing flights, so we can't use a simple random sample. A judgment sample would not provide an objective basis for assessing heavy fines. A raeasonable option would be for ground observers to record the aircraft size, type, and carrier (airline) for each departing flight for a week and use this information to construct a stratified sample.

> AACSB: Analytic Blooms: Evaluate Difficulty: 3 Hard Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 62. Professor Hardtack chose a sample of 7 students from his statistics class of 35 students by picking every student who was wearing red that day. Which kind of sample is this?
 - A. Simple random sample
 - B. Judgment sample
 - C. Systematic sample
 - D. Convenience sample

Quick but may not be representative of all students.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 63. Thirty work orders are selected from a filing cabinet containing 500 work order folders by choosing every 15th folder. Which sampling method is this?
 - A. Simple random sample
 - <u>B.</u> Systematic sample
 - C. Stratified sample
 - D. Cluster sample

Classic systematic sample from an accessible but unlisted population.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 64. Which of the following is not a likely reason for sampling?
 - A. The destructive nature of certain tests
 - B. The physical impossibility of checking all the items in the population
 - C. Prohibitive cost of studying the entire population
 - D. The expense of obtaining random numbers

Random numbers are cheap (e.g., Excel).

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts

- 65. Comparing a census of a large population to a sample drawn from it, we expect that the:
 - <u>A.</u> sample is usually a more practical method of obtaining the desired information.
 - B. accuracy of the observations in the census is surely higher than in the sample.
 - C. sample must be a large fraction of the population to be accurate.

Census is often impractical, while samples can be extremely accurate.

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts

- 66. A stratified sample is sometimes recommended when:
 - A. the sample size is very large.
 - B. the population is small compared to the sample.
 - C. distinguishable strata can be identified in the populations.
 - D. the population is spread out geographically.

Identifiable strata such as gender, ethnicity, or region can be used.

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 67. A random sample is one in which the:
 - <u>A.</u> probability that an item is selected for the sample is the same for all population items.
 - B. population items are selected haphazardly by experienced workers.
 - C. items to be selected from the population are specified based on expert judgment.
 - D. probability of selecting a population item depends on the item's data value.

Each item must have the same chance of being picked if the sample is random.

Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

68. An advantage of convenience samples over random samples is that:

- A. they are easy to analyze.
- B. it is easier to determine the sample size needed.
- C. it is easier to calculate the sampling errors involved.
- D. data collection cost is reduced.

Convenience samples are often used because they are quick (but maybe not accurate).

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 69. To measure satisfaction with its cell phone service, AT&T takes a stratified sample of its customers by age, gender, and location. Which is an advantage of this type of sampling, as opposed to other sampling methods?
 - A. It is less intrusive on customers' privacy.
 - B. It does not require random numbers.
 - C. It gives faster results.
 - D. It can give more accurate results.

Stratified sampling can yield more complete and accurate information.

AACSB: Diversity Blooms: Analyze Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 70. An accounting professor wishing to know how many MBA students would take a summer elective in international accounting did a survey of the class she was teaching. Which kind of sample is this?
 - A. Simple random sample
 - B. Cluster sample
 - C. Systematic sample
 - D. Convenience sample

She may bias the estimate because only accounting students were surveyed.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 71. A binary variable (also called a dichotomous variable or dummy variable) has:
 - <u>A.</u> only two possible values.
 - B. continuous scale values.
 - C. rounded data values.
 - D. ordinal or interval values.

Binary variables are used in every field of business to code qualitative (nominal) data.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions
- 72. A population has groups that have a small amount of variation within them, but large variation among or between the groups themselves. The proper sampling technique is:
 - A. simple random.
 - <u>B.</u> stratified.
 - C. cluster.
 - D. judgment.

Identifiable strata call for stratified sampling if you can afford the extra time and cost.

AACSB: Analytic Blooms: Apply Difficulty: 3 Hard Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 73. A manager chose two people from his team of eight to give an oral presentation because she felt they were representative of the whole team's views. What sampling technique did she use in choosing these two people?
 - A. Convenience
 - B. Simple random
 - C. Judgment
 - D. Cluster

Expert judgment may be better than just pointing a finger (we hope).

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 74. Sampling bias can best be reduced by:
 - A. using appropriate data coding.
 - B. having a computer tabulate the results.
 - C. utilizing random sampling.
 - D. taking a judgment sample.

Sampling error can't be eliminated, but sampling bias can be avoided.

Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

75. A sampling technique used when groups are defined by their geographical location is:

A. cluster sampling.

- B. convenience sampling.
- C. judgment sampling.
- D. random sampling.

Strata based on location can be targeted through cluster sampling.

AACSB: Analytic Blooms: Remember Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 76. If we choose 500 random numbers using Excel's function=RANDBETWEEN(1,99), we would most likely find that:
 - A. numbers near the mean (50) would tend to occur more frequently.
 - B. numbers near 1 and 99 would tend to occur less frequently.
 - <u>C.</u> some numbers would occur more than once.
 - D. the numbers would have a clear pattern.

On average, we'd expect each number to occur around five times.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 77. A problem with nonrandom sampling is that:
 - A. larger samples need to be taken to reduce the sampling error inherent in this approach.
 - <u>B.</u> not every item in the population has the same chance of being selected, as it should.
 - C. it is usually more expensive than random sampling.
 - D. it generally provides lower response rates than random sampling.

Only random sampling gives every item the same chance to be picked.

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 78. From its 32 regions, the FAA selects 6 regions, and then randomly audits 25 departing commercial flights in each region for compliance with legal fuel and weight requirements. This is an example of:
 - A. simple random sampling.
 - B. stratified random sampling.
 - C. cluster sampling.
 - D. judgment sampling.

Two-stage cluster sampling is being used (a special form of stratification).

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 79. Which of the following is a correct statement?
 - A. Choosing the third person listed on every fifth page of the phone book is stratified sampling.
 - <u>B.</u> An advantage of a systematic sample is that no list of enumerated data items is required.
 - C. Convenience sampling is used to study shoppers in convenience stores.
 - D. Judgment sampling is an example of true random sampling.

Review the sampling methods and their characteristics.

AACSB: Analytic Blooms: Analyze Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 80. Which of the following is false?
 - A. Sampling error is the difference between the true parameter and a random number in Excel.
 - B. Sampling error is a result of unavoidable random variation in a sample.
 - C. A sampling frame is used to help identify the target population in a statistical study.
 - <u>D.</u> The target population must first be defined by a full list or data file of individuals.

Review the terminology of sampling.

AACSB: Analytic Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts

- 81. When we are choosing a random sample and we do not place chosen units back into the population, we are:
 - A. sampling with replacement.
 - B. sampling without replacement.
 - C. using a systematic sample.
 - D. using a voluntary sample.

Sampling without replacement avoids redundancy, yet nonreplacement is biased.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 82. Which method is likely to be used by a journalism student who is casually surveying opinions of students about the university's cafeteria food for an article that she is writing?
 - A. Simple random sample
 - B. Systematic random sample
 - C. Cluster sample
 - D. Convenience sample

Quick and easy may trump true random sampling for a busy journalist.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 83. Which of the following is false?
 - A. Mail surveys are cheap but have low response rates.
 - <u>B.</u> Coverage error is when respondents give untruthful answers.
 - C. Focus groups are nonrandom but can probe issues more deeply.
 - D. Surveys posted on popular websites suffer from selection bias.

Coverage error is when you miss some segment of the target population.

Blooms: Understand Difficulty: 2 Medium Learning Objective: 02-09 Describe basic elements of survey design; survey types; and sources of error. Topic: Surveys

- 84. Which is a time series variable?
 - A. VISA balances of 30 students on December 31 of this year
 - B. Net earnings reported by Xena Corp. for the last 10 quarters
 - C. Dollar exchange rates yesterday against 10 other world currencies
 - D. Titles of the top 10 movies in total revenue last week
 - If x_1, x_2, \ldots, x_n do not refer to n time periods, it isn't a time series.

AACSB: Analytic Blooms: Apply Difficulty: 2 Medium Learning Objective: 02-03 Explain the difference between time series and cross-sectional data. Topic: Level of Measurement

- 85. An observation in a data set would refer to:
 - A. only a variable whose value is recorded by visual inspection.
 - B. a data item whose value is numerical (as opposed to categorical).
 - <u>C.</u> a single row that contains one or more observed variables.
 - D. the values of all the variables in the entire data set.

We usually put observations in rows on a spreadsheet, while each column is a variable.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-01 Use basic terminology for describing data and samples. Topic: Definitions

- 86. A multivariate data set contains:
 - A. more than two observations.
 - B. more than two categorical variables.
 - C. more than two variables.
 - D. more than two levels of measurement.

When you have more than two variables it is multivariate data.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-01 Use basic terminology for describing data and samples. Topic: Definitions

- 87. The Centers for Disease Control and Prevention (CDC) wants to estimate the average extra hospital stay that occurs when heart surgery patients experience postoperative atrial fibrillation. They divide the United States into nine regions. In each region, hospitals are selected at random within each hospital size group (small, medium, large). In each hospital, heart surgery patients are sampled according to known percentages by age group (under 50, 50 to 64, 65 and over) and gender (male, female). This procedure combines which sampling methods?
 - A. Systematic, simple random, and convenience
 - B. Convenience, systematic, and judgment
 - C. Cluster, stratified, and simple random
 - D. Judgment, systematic, and simple random

Identifiable strata were sampled, but also random within strata and regional clusters.

AACSB: Analytic Blooms: Evaluate Difficulty: 3 Hard Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 88. Which statement is correct?
 - <u>A.</u> Selecting every fifth shopper arriving at a store will approximate a random sample of shoppers.
 - B. Selecting only shoppers who drive SUVs is a stratified sampling method.
 - C. A census is preferable to a sample for most business problems.
 - D. Stratified samples are usually cheaper than other methods.

Done carefully, systematic sampling is close to random when there is no list.

AACSB: Analytic Blooms: Evaluate Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

- 89. Which is a categorical variable?
 - A. The brand of jeans you usually wear
 - B. The price you paid for your last pair of jeans
 - C. The distance to the store where you purchased your last pair of jeans
 - D. The number of pairs of jeans that you own

Categories have only names (e.g., Calvin Klein).

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

90. Which is a discrete variable?

- A. The time it takes to put on a pair of jeans
- B. The price you paid for your last pair of jeans
- C. The distance to the store where you purchased your last pair of jeans
- D. The number of pairs of jeans that you own

The "number of" anything is discrete numerical data.

AACSB: Analytic Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-02 Explain the distinction between numerical and categorical data. Topic: Definitions

- 91. A section of the population we have targeted for analysis is:
 - A. a statistic.
 - <u>B.</u> a frame.
 - C. a sample.
 - D. a coven.

We must define the segment we want to look at (e.g., independent voters).

AACSB: Analytic

Blooms: Remember Difficulty: 1 Easy Learning Objective: 02-06 Use the correct terminology for samples and populations. Topic: Sampling Concepts

Short Answer Questions

92. Which survey method would you recommend to survey opinions of airline passengers about the cleanliness of the restrooms in the Detroit airport? Why not the others?

Restroom users are usually in a hurry and do not wish to talk to anyone while entering or leaving the facility, so direct observation or handout surveys would not work. A questionnaire could be e-mailed or mailed to a sample of frequent flyers. No telephone surveys because people would distrust the call. A web survey would have nonresponse bias.

Feedback: This is a difficult sampling problem. Restroom users are usually in a hurry and do not wish to talk to anyone while entering or leaving the facility. Thus, direct observation or handout surveys would not work. Perhaps a questionnaire could be e-mailed or mailed to a sample of frequent flyers who departed or arrived at this airport in a recent month, if a major airline were willing to cooperate. This assumes that frequent flyers are a reasonable target population. You would probably not use a telephone survey because people would distrust the call. A web survey would have nonresponse bias (i.e., respondents would probably be those with a gripe). Students may digress into sampling methods (random, systematic, cluster, stratified) instead of survey types. Ingenious students may propose other novel methods of sampling opinions.

AACSB: Reflective Thinking Blooms: Evaluate Difficulty: 3 Hard Learning Objective: 02-09 Describe basic elements of survey design; survey types; and sources of error. 93. What kind of sampling method would you suggest in order to tabulate the number of formulas on a typical page of the Doane-Seward textbook? Defend your choice.

Pages are numbered so a simple random sample would be easy, or a systematic sample (e.g., every 20th page starting at page 17). Case could be made for cluster or stratified samples (e.g., by chapter or topic).

Feedback: Since the pages are numbered, a simple random sample would be quite easy. Have Excel print n random integers between 001 and 773 (or whatever the length of the book is). You would want to exclude the table of contents, appendixes, indexes, and so on). Another good choice would be a systematic sample (e.g., every 20th page starting at page 17) or a variation such as every page divisible by 20. These methods would be unbiased. Possible cases could be made for cluster or stratified samples (e.g., by chapter or topical area), but these might offer little gain.

> AACSB: Reflective Thinking Blooms: Evaluate Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods

94. How would you design a study to see whether drivers using hands-free cell phones are distracted enough to slow their reactions to emergency situations? How would you collect data?

No observation of drivers (too dangerous). Tests using a simulator would permit data to be collected automatically and would permit stratified sampling by driver characteristics (e.g., age group, gender, cell phone type).

Feedback: Don't use direct observation of drivers (too dangerous). Tests using a simulator would permit data to be collected automatically on reaction times to emergencies. This would permit stratified sampling by driver characteristics (e.g., age group, gender, cell phone type). Many answers are possible. The emphasis should be on how carefully the student has thought about the question. Beware of simplistic answers or impossible data collection schemes.

AACSB: Reflective Thinking Blooms: Evaluate Difficulty: 3 Hard Learning Objective: 02-09 Describe basic elements of survey design; survey types; and sources of error. Topic: Surveys 95. Explain the concept of a focus group. In what ways does a focus group resemble a survey? Why is a moderator desirable? What else is required to make a successful focus group?

Data collected in a focus group are richer in qualitative details and may contain information that would be missed in a survey. A trained moderator can help keep the group on track and manage interpersonal issues that may arise.

Feedback: Data collected in a focus group are richer in qualitative details, and may contain information that would be missed in a survey. Yet a focus group is like a survey in that it seeks to extract useful information and patterns from individuals. Participants are not chosen completely at random, but rather are selected to represent different backgrounds and diverse viewpoints of interest to the researchers. A well-trained moderator can help keep the group on track and manage interpersonal issues that may arise.

> AACSB: Reflective Thinking Blooms: Evaluate Difficulty: 2 Medium Learning Objective: 02-07 Explain the common sampling methods and how to implement them. Topic: Sampling Methods